

NIES-Collection

LIST OF STRAINS

Sixth Edition

2000

Microalgae

and

Protozoa

Edited by

Makoto M. Watanabe, Masanobu Kawachi,

Mikiya Hiroki and Fumie Kasai

Supervised by

Committee for Evaluating Microbial Culture Strains

National Institute for Environmental Studies

Environment Agency

JAPAN

NIES-Collection. List of Strains
Sixth Edition
Microalgae and Protozoa
March 1, 2000

Contributors:

Makoto M. Watanabe, Mikiya Hiroki, Fumie Kasai,
Masanobu Kawachi, Akira Shimizu,
Mayumi Erata, Fumi Mori and Kosei Yumoto

Acknowledgment

Sincere thanks are expressed to Dr. H. Nozaki, University of Tokyo, Dr. S. Suda, *Marine Biotechnology Institute Co., Ltd.*, and Ms. T. Suzuki, *National Institute for Environmental Studies*, for their helpful advice and technical assistance about the taxonomy and the maintenance of some strains.

Cover design : Mayumi Erata

第六版の序

国立環境研究所微生物系統保存施設が、1997年に保存株リスト第五版を発行してから、3年を迎えることとなった。初版～五版に関して、国内国外の各方面から多くの建設的意見や激励が寄せられたことには非常に勇気づけられたと共に、我々の事業が環境科学分野のみならず、基礎生物学、農学、水産学、食品学、医学等の分野でも注目され、重要視されていることを知り、責任の重さを痛感したものである。

この第六版は、初版～五版と同様に微生物系統保存株評価委員会の監修を受け、微細藻類 752 株、原生動物 2 株を掲載した。特に保存株の分類、保存株特性については注意深い検討がなされたが、不備な点をご指摘願えれば幸いである。

本施設に保存されている微生物株の殆どは、わが国の藻類学者によって分離培養されたものであり、他の微生物保存機関には保存されていないものである。今後、貴重な微生物株については、国内国外の微生物保存機関と密接な連携・協力関係を組み、共通のルールで共有していくことを考えている。また、本施設の事業は、微生物株の収集・保存・分譲にとどまらず、分類学的研究、保存技術の開発、株情報の収集およびその電算機管理システムの開発等多岐に亘っているが、これらの事業が益々充実し成果をあげるために、施設・要員の充実と拡充をはかっていく所存である。今後とも一層のご批判とご支援を賜ることができれば幸いである。

最後に、寄託依頼された藻類株の評価並びに本リスト刊行に際して様々なご指導とご助言をいただいた評価委員会委員に深甚な謝意を表するとともに、微生物系統保存施設のスタッフ一同の熱意に満腔の敬意を表したい。

平成12年3月

国立環境研究所微生物系統保存株評価委員会委員長
国立環境研究所生物圏環境部長

渡 辺 信

保存株リスト第一版発刊に寄せて

国立環境研究所に我が国最初の環境微生物の系統保存施設が設置されたのは、昭和58年1月であったが、その後約2年間にわたって、同研究所の関係者の並々ならぬ努力によって、微生物保存事業に関する周到なる準備作業が繰り展げられ、ようやくここにその成果を保存株リストとして集大成されたことは、環境科学にたずさわる多くの研究者にとって、これ程慶ばしいことはない。ここに関係者各位に対して満腔の敬意を表明したい。

今回刊行された保存株リストは、当面環境生物学上重要な生産者である微細藻類に的を絞ったものであるが、これは我が国の現行微生物系統保存事業のうちで最も弱点とされていた分野であり、学界・産業界からもその実現が強く要望されていたところである。微細藻類の系統保存は、長年にわたり活発に研究されてきた細菌類や菌類の系統保存とは異なり、その分離、培養、保存等の条件が極めて複雑で、技術的に多くの困難な作業を伴うものである。本研究所においてはその性格上多角的研究に取り組んでいるが、その特徴を生かして所内の衆知を結集してこの点を克服し、世界的に通用する信頼度の高い系統保存事業を軌道に載せることに成功した。本施設の保存する微生物株は、その特性が科学的に実証されているために、これを実験的に使用する研究者、あるいはそれら微生物株データの利用者にとって、高い信頼感をもって利用することができる。しかも本施設では、保存微生物株に関する独自の電算機管理システムを開発したので、その保存株データを環境生物学に関するデータベースの一環として利用することが可能となった。このことによって、とかく遅れがちであった我が国環境生物学の近代化が著しく促進されるものと信ずる。

本施設の当初の目標は環境問題に関係ある多種多様の微生物株を総合的に収集保存することであったが、現状ではようやく微細藻類についての系統保存体制が確立されたに止まっている。今後益々施設設備の充実をはかって、微細藻類のみならず、環境生物学の調査研究上欠かすことのできないその他の微生物の系統保存をも実施し、名実ともにそなわった世界的な環境微生物株保存センターの一つとして発展されることを期待したい。

昭和60年2月

元富山大学長
東京大学名誉教授
柳 田 友 道

目 次

第六版の序.....	渡 辺 信
保存株リスト第一版発刊に寄せて.....	柳 田 友 道
I. はじめに.....	1
II. 培養株の寄託.....	3
1. 寄託条件.....	3
2. 寄託の手続き.....	3
III. 保存株の分譲.....	6
1. 所内研究者への分譲.....	6
2. 所外への分譲.....	6
3. "Untransportable" 株の分譲について.....	6
4. 凍結保存株の分譲について.....	6
IV. 分譲株の培養保存法.....	9
V. 藻類培地作成の基本手法.....	9
1. 保存試薬液.....	9
2. 培地作成.....	11
3. 寒天斜面培地.....	12
VI. 培地 (Media).....	30
1. 藻類の保存培地 (Stock media for algae).....	30
2. 無菌検査培地 (Bacteria-free check media).....	36
3. 微量金属混液・ビタミン混液・土壌浸出液 (Trace metals, vitamin mixes and soil extract).....	37
4. 原生動物の保存培地 (Stock medium for protozoa).....	38
VII. 保存株データの利用法.....	39
VIII. 保存株データ (Strain data).....	43
IX. 索引 (Indexes).....	117
1. 株番号索引 (Numerical index).....	117
2. 分類群別索引 (Systematic index).....	125
X. 文献 (References).....	135

I. はじめに

国立環境研究所微生物系統保存施設は、昭和58年に環境微生物の系統保存を行なうことを目的として設立された。この施設は、当研究所で遂行されている微生物学的研究で使用されている微生物の培養を、所内研究者の要望に応じて保存し、分譲することを目的としていたが、環境科学に携る微生物学者からの強い要望を配慮して、所内に止らず、広く他機関からも微生物株とそれらの株データの収集および分譲を積極的に行なうこととし、将来的には国際的な環境微生物のカルチャーコレクションセンターとして国内外の環境関連研究機関および研究者と密接なネットワーク体制を構築し、環境微生物研究の推進を支える役割を担っていくことを計画している。

本施設で保存される微生物の培養株は、表1に記されているように微細藻類、原生動物および特殊な浄化能を有する細菌類が対象となっているが、現状ではこれらすべてを同時に保存できる体制の整備が不十分であることおよび環境科学の分野では水域の汚染と浄化に微細藻類が密接に関連していることから、微細藻類株が積極的に収集・保存されている。収集されたすべての株について、その種名、培養条件、保存法、形態学的特徴、生理生態学的特徴、環境科学との関連性に関する株特性の検査や情報収集が行なわれ、更にそれらの株データ管理のパーソナルコンピューターによるシステム化が行なわれている。

表1 本施設に保存される対象となる微生物株

環境問題との関連性での類型	対象となる微生物株
環境汚染の原因となる微生物	赤潮形成藻類、水の華形成藻類、有毒藻類、水道水の異味異臭をもたらす藻類または放線菌類、硫酸還元細菌等
環境汚染の指標となる微生物	AGP供試藻類、重金属耐性微生物、水質の富栄養化の指標となる細菌類、微細藻類、原生動物等
自浄作用、廃水および廃棄物処理に関する微生物	光合成細菌、脱窒菌、硝化細菌、汚染原因微生物を捕食または溶解する微生物、活性汚泥および生物膜処理の原生動物および細菌類、嫌気性処理にかかわる嫌気性細菌、生物学的処理の障害となる微生物等
有機合成化合物の分解に関する微生物	PCB、フェノール、各種除草剤および農薬等の分解に関与する細菌類
金属の酸化・還元作用に関連する微生物	塩化水銀(HgCl_2)やシアン化水銀の還元に関与する細菌類、亜硫酸の酸化に関与する細菌類、重金属のバクテリアリーチングに関与する細菌類等

本施設に保存された環境微生物培養株の最初のリストには、施設、組織、基本業務の概要説明とともに、微細藻類262株が掲載された(文献395, 396)。それ以降、施設、組織、基本業務の大きな変化はないが、寄託された株、安定した増殖が得られた株および株データの変更を行なった株があり、それらは追補株リストおよび第2版、第3版、第4版、第5版として掲載された(文献397, 398, 400, 409, 408, 390)。現在、微細藻類752、原生動物2株が保存されるに至っている。第6版は、これらの保存株すべてを再整理し、新たなデータを加えて、掲載したものである。

II. 培養株の寄託

1. 寄託条件

微生物の培養株の本施設への保存寄託は、以下の条件を満たしている培養株で、微生物系統保存株評価委員会の審査を経たものとする。

- (1) 寄託の対象となる微生物は原則として以下のいずれかにあてはまることとする。
 - (i)環境汚染の原因となる微生物、(ii)環境汚染の指標となる微生物、(iii)自浄作用、廃水及び廃棄物処理に関係する微生物、(iv)有機合成化合物の分解に関係する微生物、(v)金属の酸化・還元作用に関係する微生物。
- (2) 種名及び履歴が明らかである培養株であることを原則とするが、既に多くの調査研究において属名をもって使用されている微生物株については例外として受け入れる。
- (3) 寄託対象保存株は、保存条件が確立している培養株、すなわち保存中の状態が安定しており、次のいずれかにあてはまる培養株であることとする。
 - (i)微細藻類ではクローン培養株か単藻培養株であり、無菌培養株であることが望ましい、(ii)原生動物では無菌培養株か餌料としての他の微生物のみが混入している単一種培養株であること、(iii)細菌類はすべて純粋培養株であること。
- (4) 寄託された培養株は原則としてすべて分譲対象として扱う。
- (5) その他、特に微生物系統保存株評価委員会が必要と認めたもの。

2. 寄託の手続き

- (1) 寄託者は様式-1の書類に所定事項を記入の上、下記の寄託先へ申し込むこととする。

〒305-0053 茨城県つくば市小野川16-2 国立環境研究所 微生物系統保存施設
電話 0298(50)2556 FAX 0298(50)2587
- (2) 受託可否は寄託依頼があった日から1ヶ月以内に行う。
- (3) 寄託者は受託の回答があった日から1ヶ月以内に、微生物株を本施設に寄託するものとする。
- (4) 寄託書類の記載事項と寄託された微生物の状態が一致せず、前述した寄託条件より逸脱した場合には、寄託のあった日より1ヶ月以内に受託の取り消しを寄託者へ知らせることとする。

微生物株寄託依頼書

国立環境研究所
微生物系統保存施設 殿

国環研記入
受付日 _____
受付担当者 _____
受付番号 _____
受託 可 否

年 月 日
(フリガナ) 依頼者名 _____
所属機関 (日本語名) _____
(英語名) _____
所属機関住所 〒□□□□-□□□□
電話 () (内線)
FAX ()
Eメールアドレス

下記微生物の寄託を依頼します。

寄託理由

① 学名及び命名者名 _____
② 株番号又はシンボル _____
③ 履歴 1. 採集場所: _____ 2. 生息環境 (25ページより番号で記入してください。): _____ 3. 採集年月日: 年 月 日 (フリガナ) 4. 採集者名: _____ 5. 分離年月日: 年 月 日 (フリガナ) 6. 分離者名: _____ 7. 分離試料源: <input type="checkbox"/> 土, <input type="checkbox"/> 底泥, <input type="checkbox"/> 水, <input type="checkbox"/> 植物 (), <input type="checkbox"/> 動物 (), <input type="checkbox"/> 雪または氷, <input type="checkbox"/> その他 () 8. 分離時の生物の状態: <input type="checkbox"/> 運動性栄養細胞, <input type="checkbox"/> 非運動性栄養細胞, <input type="checkbox"/> 休眠細胞, <input type="checkbox"/> 孢子, <input type="checkbox"/> その他 () 9. 分離方法: <input type="checkbox"/> ピペット洗浄法, <input type="checkbox"/> 希釈法, <input type="checkbox"/> 寒天平板法, <input type="checkbox"/> 走性, <input type="checkbox"/> その他 () 10. 分離時の処理: <input type="checkbox"/> 無処理, <input type="checkbox"/> 抗生物質, <input type="checkbox"/> 紫外線照射, <input type="checkbox"/> 化学物質, <input type="checkbox"/> 熱処理, <input type="checkbox"/> 超音波処理, <input type="checkbox"/> 集積(強化)培養, <input type="checkbox"/> その他 () (フリガナ) 11. 同定者名: _____ (フリガナ) 12. 無菌化者名: _____ (フリガナ) 13. クローン化者名: _____

様式-1 (2)

④ 株の状態

1. 微細藻類 無菌, 単藻, クローン, 二種混合
2. 細菌類 純粹, 非クローン
3. 原生動物 無菌, 単一種混菌, 二種混菌, 混合
4. その他 (_____)

⑤ 培地

1. 培地名及び出典: _____
 2. 培地組成及び作成上の注意
(通常よく使用されている培地の場合、原典を記すだけでよい。)
- _____

⑥ 培養条件

1. 温度: _____ 2. 照度: _____
3. 光源種類: _____ 4. 明暗周期: _____

⑦ 保存条件

- 継代培養条件
1. 温度: _____ 2. 照度: _____
 3. 光源種類: _____ 4. 明暗周期: _____
 5. 継代周期: _____

凍結保存

- 可 否
1. 凍害防御物質: なし, DMSO, PVP, グリセロール,
メタノール, その他 (_____)
濃度 (_____ %)
 2. 凍結速度: _____
 3. 融解条件: 40℃ウォーターバス中
その他 (_____)

4. 保存温度: 液体窒素, ディープフリーザー (_____ °C)
その他 (_____)

- 凍結乾燥保存
可 否

- L-乾燥保存
可 否

⑧ 株特性

1. 環境上問題となる特性 (25ページより番号で記入してください。)
2. 生理生態的特性 (25ページより番号で記入してください。)
3. その他の特性 (25ページより番号で記入してください。)

⑨ 遺伝子データ

1. 遺伝子名: _____
2. 登録番号: _____
(フリガナ)
3. 登録者名: _____
4. 登録年月日: _____ 年 _____ 月 _____ 日

⑩ その他の情報

- ⑪ この株に関する文献がある場合は、別刷りまたはコピーを2部ずつ添付してください。

III. 保存株の分譲

1. 所内研究者への分譲

(1) 分譲条件

- i) 分譲された株を使った研究成果を論文として発表する場合は、NIES株番号（例：“NIES-125”）と本施設から分譲を受けたことを明記し、別刷りまたはコピーを2部ずつ本施設に送ることとする。
- ii) 分譲された株を第三者に分譲することを禁止する。
- iii) 株データの分譲については、保存株の分譲に準じて行われる。

(2) 分譲依頼の手続き

- i) 分譲希望者は様式-2の書類に所定事項を記入の上、本施設へ申し込むこととする。
- ii) 分譲を受けた者は受領後直ちに培養株の状態について、様式-3の書類に所定事項を記入の上、本施設へ提出するものとする。

2. 所外への分譲

本施設に保存されている微生物株の所外への分譲は、(財)地球・人間環境フォーラムで行われている。分譲依頼等はフォーラム発行のカタログを参照されたい。

3. “Untransportable”株の分譲について

保存株リストの「株の性質」の項において“Untransportable”と記載されている株（40頁を参照のこと）についての分譲依頼は季節や株の生育状態等により受け付けられないことがあるので、これらの株の分譲依頼にあたっては必ず事前に本施設へ問い合わせるものとする。

また当該株の海外への分譲は、持ち帰りの場合を除き、原則として行わないものとする。

4. 凍結保存株の分譲について

保存株リストの「培養条件」の項において“[Cryopreserved]”と記載されている株（40頁を参照のこと）は現在凍結保存のみで維持されている。これらの株の分譲については、依頼を受領した時点で解凍・再培養を開始するため引渡し（発送）までに最低3～4週間を要する。分譲依頼にあたってはあらかじめこの点を考慮されたい。

様式-2

微生物株分譲依頼書

国立環境研究所
微生物系統保存施設 殿

国環研記入
受付日_____
受付者_____
受付番号_____

研究目的（具体的に）

年 月 日

（フリガナ）
依頼者名 _____

所属機関（日本語名） _____

（英語名） _____

所属機関住所
〒□□□-□□□□

電話 () (内線)

FAX ()

Eメールアドレス

株データ

要 (株番号)

不要

下記微生物についての分譲を依頼します。

微生物学名及び株番号

国環研担当者記入

微生物株の受領と受領時の状態についての報告

国立環境研究所
微生物系統保存施設 殿

国環研記入
受付日 _____
受付者 _____
受付番号 _____

年 月 日

(フリガナ)
依頼者名 _____

所属機関 (日本語名) _____

(英語名) _____

所属機関住所

〒□□□-□□□□

電話 () (内線)

FAX ()

Eメールアドレス

年 月 日に分譲されました微生物株の受領と分譲時の株
の状態について下記のように報告いたします。

分譲株 (微生物学名及び株番号)

株の受領時の状態

良好株

不良株

その他

当施設についての意見と要望

国環研担当者記入

IV. 分譲株の培養保存法

微生物株は、ねじ口試験管に培養された状態で郵送される。株の分譲を受けた場合、株を絶やさないために下記の点に留意する必要がある。

- i) 培地は株を受け取る前に作成しておく。
- ii) 株を受領後速やかに荷をとき、新鮮な培地に植え継ぎ、当方で指示した温度と照度下（第Ⅷ章参照）で培養する。その場合明暗サイクルは12時間明期12時間暗期とし、ねじ口試験管のねじ蓋をゆるくする。
- iii) 良好な増殖が確認された後に、更に株を保存する場合には、当方で指示した期間毎に新鮮な培地に移植する必要がある（第Ⅷ章参照）。

V. 藻類培地作成の基本手法

藻類株の保存には、数多くの培地を必要とする。それぞれの培地は次章に掲載した処方せんに従って作成されるが、正確かつ簡便に培地を作成するために、本施設で採用している基本手法について述べておきたい。

1. 保存試薬液

培地は一般に多量栄養素、微量金属、およびビタミン類(表2)で構成されている。これらの諸成分の保存試薬液を作成しておくことが、培地作成の簡便さをもたらす。このうち微量金属やビタミン類の保存液の濃度は非常に低いので、保存試薬液作成時には、より濃度の高い原液を作成する必要がある。以下、各々について保存試薬液の濃度と作成方法について述べる。

A 多量栄養素：各栄養素につき、10mg/mlの濃度の保存試薬液を別々に作成し、冷蔵庫(5℃)で保管する。

B 微量金属：これらの成分は、各種の保存試薬液として別々に作成され保管される場合と、混液で保管される場合がある。

(1) 各種保存試薬液

- i) 10-100mg/mlの濃度で各種金属の原液を作成する。
- ii) 各原液を1mg/mlの濃度に希釈し冷蔵庫(5℃)に保管する。

IV. 分譲株の培養保存法

微生物株は、ねじ口試験管に培養された状態で郵送される。株の分譲を受けた場合、株を絶やさないために下記の点に留意する必要がある。

- i) 培地は株を受け取る前に作成しておく。
- ii) 株を受領後速やかに荷をとき、新鮮な培地に植え継ぎ、当方で指示した温度と照度下（第Ⅷ章参照）で培養する。その場合明暗サイクルは12時間明期12時間暗期とし、ねじ口試験管のねじ蓋をゆるくする。
- iii) 良好な増殖が確認された後に、更に株を保存する場合には、当方で指示した期間毎に新鮮な培地に移植する必要がある（第Ⅷ章参照）。

V. 藻類培地作成の基本手法

藻類株の保存には、数多くの培地を必要とする。それぞれの培地は次章に掲載した処方せんに従って作成されるが、正確かつ簡便に培地を作成するために、本施設で採用している基本手法について述べておきたい。

1. 保存試薬液

培地は一般に多量栄養素、微量金属、およびビタミン類(表2)で構成されている。これらの諸成分の保存試薬液を作成しておくことが、培地作成の簡便さをもたらす。このうち微量金属やビタミン類の保存液の濃度は非常に低いので、保存試薬液作成時には、より濃度の高い原液を作成する必要がある。以下、各々について保存試薬液の濃度と作成方法について述べる。

A 多量栄養素：各栄養素につき、10mg/mlの濃度の保存試薬液を別々に作成し、冷蔵庫(5℃)で保管する。

B 微量金属：これらの成分は、各種の保存試薬液として別々に作成され保管される場合と、混液で保管される場合がある。

(1) 各種保存試薬液

- i) 10-100mg/mlの濃度で各種金属の原液を作成する。
- ii) 各原液を1mg/mlの濃度に希釈し冷蔵庫(5℃)に保管する。

表2. 培地に使われる各種栄養素

多量栄養素	微量金属
NaCl	H ₃ BO ₃
KCl	MnCl ₂ · 4H ₂ O
CaCl ₂ · 2H ₂ O	MnSO ₄ · 7H ₂ O
MgCl ₂ · 6H ₂ O	FeCl ₃ · 6H ₂ O
Na ₂ SO ₄	FeSO ₄ · 7H ₂ O
K ₂ SO ₄	CoCl ₂ · 6H ₂ O
MgSO ₄ · 7H ₂ O	ZnSO ₄ · 7H ₂ O
NaNO ₃	CuSO ₄ · 5H ₂ O
KNO ₃	Na ₂ MoO ₄ · 2H ₂ O
Ca(NO ₃) ₂ · 4H ₂ O	ビタミン類
NH ₄ NO ₃	Vitamin B ₁₂
NaH ₂ PO ₄ · 2H ₂ O	Biotin
β -Na ₂ glycerophosphate · 5H ₂ O	Thiamine HCl
KH ₂ PO ₄	Nicotinic acid
K ₂ HPO ₄	Calcium panthothenate
Na ₂ CO ₃	<i>p</i> -Aminobenzoic acid
NaHCO ₃	Inositol
Na ₂ SiO ₃ · 9H ₂ O	Folic acid
	Thymine

(2) 混液

- i) (1)-i)と同様の操作を行う。
- ii) 必要量の80%の蒸留水をビーカーに加える。
- iii) 十分に攪拌しながら必要量のNa₂EDTAを溶解する。
- iv) 十分に攪拌しながら各種微量金属原液を必要量添加する。
- v) 蒸留水を加え、最終量に調整し、冷蔵庫(5℃)に保管する。

C ビタミン類：ビタミンB₁₂、ビオチン、チアミンの3種のビタミンだけで多くの藻類が増殖するので、殆どの培地はこれら3種のビタミン類だけが添加されている。しかし、いくつかの培地では、他のビタミン類が添加されている。

(1) ビタミンB₁₂、ビオチン、チアミン

- i) ビタミンB₁₂とビオチンについては、各々0.1mg/mlの原液を作成し、チアミンについては10mg/mlの原液を作成する。

- ii) これらの原液を多数の試験管に1mlずつ分注し、オートクレーブ滅菌(121°C, 20min)後、-20°Cのフリーザーに保管する。
 - iii) 各ビタミンについて、保存原液の1mlを融解し、蒸留水で1/100に希釈してビタミンB₁₂、ビオチンについては1μg/mlの保存試薬液、チアミンについては、100μg/mlの保存試薬液を作成し、冷蔵庫に保管しながら使用する。
- (2) 他のビタミン類：ある培地では、多種のビタミン類が混液の形で添加される(第VI章-56参照)。大量に作成しておくことをすすめる。
- i) 各種のビタミンについて0.1-1mg/mlの原液を作成する。
 - ii) 必要量の80%の蒸留水をビーカーに加える。
 - iii) 十分に攪拌しながら各種ビタミンを必要量加える。
 - iv) 蒸留水で最終量に調整する。
 - v) ミリポアフィルター(0.22μm)でろ過滅菌したのち、滅菌された試薬瓶に100mlずつ分注し、-20°Cのフリーザーで保管する。一部を融解し、冷蔵庫(5°C)に保管しながら使用する。

2. 培地作成

培地は、合成培地と強化培地に大別される。すべての淡水藻や一部の海産藻は合成培地で、殆どの海産藻は強化培地で保存されている。

(1) 合成培地(淡水)

- i) 必要量の80-90%の蒸留水をビーカーに加える。
- ii) 十分に攪拌しながら、Tris、glycylglycine、HEPES、TAPS、Bicine、MES等の緩衝剤(必要とされる場合)を必要量天秤で秤量し、添加する。
- iii) 各種栄養塩を各々の保存液から必要量添加する。
- iv) 蒸留水で最終量に調整する。
- v) 緩衝剤が使用されている場合、1N HClあるいは、1N NaOHで、使用されていない場合は各々1/10の濃度でpHを調整する。
- vi) 培地10mlずつ試験管(18×150mm)に分注し、オートクレーブで滅菌する(121°C, 20min)。

(2) 合成培地(海水)

- i) 必要量の80%の蒸留水をビーカーに加える。
- ii) 十分に攪拌しながら、緩衝剤(Tris, NTA等)および多量栄養塩類(NaCl, MgSO₄·7H₂O, KCl, CaCl₂·2H₂O)を必要量天秤で秤量し、添加する。
- iii) 他の各種栄養塩を各々の保存液から必要量添加する。
- iv) 蒸留水で最終量に調整する。
- v) 1N HClでpHを調整する(通常8.0)。
- vi) 培地10mlずつ試験管に分注し、オートクレーブで滅菌する(121°C, 20min)。

(3) 強化海水培地

- i) 汚染のない外洋海水を採取し、ワットマンGF/Cフィルターでろ過し、粒子を除く。
- ii) 塩分を調べる。通常の外洋海水の塩分は約35%である。
- iii) 必要量の80-90%の蒸留水をビーカーに加える。
- iv) 必要量のTris等の緩衝剤を天秤で秤量し、溶解する(必要とされる場合)。
- v) 他の栄養塩類を各々の保存液から必要量添加する。
- vi) 海水で最終量に調整する。
- vii) pHを測定する。指示されている場合は1N HClで調整する(通常8.0)。
- viii) 培地10mlずつ試験管に分注し、オートクレーブで滅菌する(121℃, 20min)。

3. 寒天斜面培地

通常寒天は1.5%の濃度で滅菌する前に液体培地に加えられる。

- i) 寒天を必要量天秤で秤量し、液体培地に添加し、オートクレーブで121℃に熱し、溶解する。
- ii) 溶解後、速やかに10mlずつ試験管に分注し、オートクレーブで滅菌する(121℃, 20min)。
- iii) 滅菌後、試験管上部に直径1cmの枕木をして寝かせ、放冷して培地を斜面状に固まらせる。

PREFACE TO THE SIXTH EDITION

Three years have past since we published the fifth edition of the list. During this period a considerable number of new cultures have been added to the NIES-Collection. We appreciate the many comments and words of encouragement about the publications from people in diverse places. These have led us to recognize more than ever the value of the NIES-Collection for research and development. Its use extends not only to environmental science, but also to basic biology and microbiology-related fields such as agriculture, fisheries, food manufacture and medical science.

The sixth edition lists 752 strains of microalgae and 2 strains of protozoa. These have been evaluated by the Committee for Evaluating Microbial Culture Strains, which is composed of microbiologists from this institute and authorities from other organizations. Although special care has been exercised to ascertain that the taxonomy and characteristics of all strains are clear and precise, we are always grateful for further advice and criticism.

Most of the strains in the NIES-Collection were isolated originally by phycologists in our country and do not exist in other collections. We plan to share responsibility for preservation of the important strains by keeping close contacts with other culture collections.

The NIES-Collection carries out such wide-ranging activities as collection, preservation, distribution, taxonomy, preservation technology, and development of a microbial strain data processing system. We hope to make steady progress in these various activities by expansion of facilities and staff. We would much appreciate your advice, criticism and cooperation concerning the performance of the NIES-Collection.

I wish to express my sincere thanks to all of the members of the committee for their effort devoted to the evaluation of microbial strains for deposit, and their numerous considerations and suggestions for this publication. I would also like to pay my respect to the staffs of the NIES-Collection for their enthusiasm for culture collection.

March 1, 2000



Makoto M. Watanabe, D.Sc.

Chairman for the Committee for
Evaluating Microbial Culture Strains
Director of Environmental Biology Division

PREFACE TO THE FIRST EDITION

In January 1983, the first culture collection of environmental microorganisms in Japan was established at the National Institute for Environmental Studies. In the two years since that time, many dedicated people have collaborated in the collection of microorganisms for the institute. The fruits of their efforts have culminated in a "List of Strains," which I feel will be highly praised by environmental scientists. I would like to extend to all who were involved, my most sincere thanks and gratitude.

The list published herein focuses on microalgae which are important primary producers in the environment. Notwithstanding the fact that there has been a high demand for microalgal collections by both the academic and industrial worlds, until the establishment of the NIES-Collection, no microalgal culture collection for environmental studies *per se* existed in Japan. Unlike the culture collection of bacteria and fungi, organisms which have been actively studied for a long time, the isolation, cultivation, and preservation of microalgae are technically much more complex. Since this institution has characteristically performed interdisciplinary studies, it was possible to conquer these difficulties, and set the culture collection of microalgae on the right path by utilizing the knowledge of its many experts.

Users of the microbial strains of the NIES-Collection will find both their quality and the data maintained about them, highly reliable because the characteristics of the microalgae have been carefully examined and re-examined. Due to the development of the strain computer data processing system, strain data have added to the general data base of environmental biology. Collectively, these developments will contribute to the rapid growth of environmental microbiology, and allow it to catch up with microbiological research in other fields.

Although the ultimate objective of the NIES-Collection is to collect and preserve a great variety of microorganisms related to environmental problems, at present only the collection of microalgal cultures has been established. I hope that in the future the NIES-Collection will preserve not only microalgae, but also other microorganisms which are indispensable to environmental biology. By planning expansion of the facilities and the staff, the NIES-Collection should develop as an international culture collection center, truly worthy of the name.

September 1, 1985



Tomomichi Yanagita, D. Sci.

Professor Emeritus of the University of Tokyo

CONTENTS

Preface to the Sixth Edition	Makoto M. Watanabe
Preface to the First Edition	Tomomichi Yanagita
I. Introduction	17
II. Deposition of Strains	18
1. Condition for deposit	18
2. Procedure for deposit	18
III. Ordering and Distribution of Strains	22
1. Distribution to researchers of this institute	22
2. Distribution to people of other organizations, both academic and commercial.	22
3. Special warning for distribution of "Untransportable" and "[Cryopreserved]" strains	22
IV. Establishment of Fresh Cultures	26
V. Basic Methods for Preparation of Algal Culture Media	26
1. Stock solutions	26
2. Media	28
3. Agar slant	29
VI. Media	30
1. Stock media for algae	30
2. Bacteria-free check media	36
3. Trace metals, vitamin mixes and soil extract	37
4. Stock medium for protozoa	38
VII. Explanatory Notes about Strain Data	41
VIII. Strain Data	43
IX. Indexes	117
1. Numerical index	117
2. Systematic index	125
X. References	135

I. INTRODUCTION

Microbial Culture Collection at the National Institute for Environmental Studies (NIES-Collection) was founded in 1983 as the first collection center of environmental microorganisms.

Although microalgae, bacteria and protozoa related to environmental problems will be preserved in this collection in the future (Table 1), microalgae associated with water pollution and cleaning have been collected and preserved the most actively as the first step. The scientific names, sources, conditions of cultivation and preservation, purity, morphological and physiological characteristics, and environmental characteristics, of all the strains collected have been re-examined, and the revised data processed using a personal computer.

The first list of environmental microorganisms preserved in the NIES-Collection (Ref. 395, 396) offered 262 strains of microalgae, together with the examples of NIES-Collection facilities, organization and fundamental pattern of research. Since then, as the result of our studies on many strains isolated by us and deposited by many other microbiologists, a considerable number of new algal strains have been added and the supplementary, the second, third, fourth and the fifth lists published (Ref. 397, 398, 400, 409, 408, 390). The total numbers of strains of algae and protozoa in the NIES-Collection are now 752 and 2, respectively.

The sixth edition lists all the strains of algae and protozoa preserved in the collection together with new or revised data.

Table 1. Microbial culture strains preserved in the NIES-Collection

Environmental Microorganisms	Examples
Causative microorganisms of environmental pollution.	Causative algae of red tide or water bloom, toxic algae, microorganisms causing the offensive odor or taste of tap water, and sulfate-reducing bacteria.
Microbial indicators of environmental pollution.	Algae used in bioassay studies of water pollution, metal resistant microorganisms, and microbial indicators of eutrophication.
Microorganisms associated with environmental self-cleaning and waste water treatment	Photosynthetic bacteria, denitrifying bacteria, microorganisms which prey upon or lyse causative organisms of environmental pollution, and bacteria and protozoa associated with activated sludge, microbial film processing or anaerobic processing.
Microorganisms associated with biodegradation of synthetic organic compounds	Bacteria associated with biodegradation of PCB, phenol and agricultural chemicals.
Microorganisms associated with oxidation and reduction of metals.	Bacteria associated with reduction of mercury chloride (II) or mercury cyanide (II), oxidation of arsenious acid, or bacterial leaching of heavy metals.

II. DEPOSITION OF STRAINS

1. Condition for deposit

The decision to accept the deposit of a strain is made by the Committee for Evaluating Microbial Culture Strains. A strain for deposit in the NIES-Collection should fit the following criteria.

- (1) It must be at least one of the following microorganism types:
 - i) Causative microorganism of environmental pollution.
 - ii) Microbial indicator of environmental pollution.
 - iii) Microorganism related to waste water treatment of self-cleaning by the environment.
 - iv) Microorganism related to the biodegradation of synthetic organic compounds.
 - v) Microorganism capable of oxidation or reduction of metals.
- (2) The source of the strain and the specific name should be established, though strains which have been used in number of studies may be accepted even if only the generic name is known.
- (3) It should be a stable culture under defined conditions and fit one of the following states:
 - i) Microalgae : clonal or unialgal strain.
 - ii) Protozoa : axenic or xenic strain with supplementary microorganisms added as food.
 - iii) Bacteria : pure strain.
- (4) As a rule, deposited strains are available to the general public.
- (5) At the discretion of the Committee for Evaluating Microbial Culture Strains, some microorganisms may be accepted for deposit, even if they do not meet the above criteria.

2. Procedure for deposit

- (1) The depositor should complete the Strain Deposit Request Form (p. 20) and send it to the following address:

Microbial Culture Collection
National Institute for Environmental Studies,
16-2 Onogawa, Tsukuba, Ibaraki 305-0053, Japan
Phone : +81-298-50-2556
Fax : +81-298-50-2587

- (2) The decision for the deposit of the strain is given within one month from the date of receipt of the Strain Deposit Request Form.
- (3) The depositor should send an actively growing or lyophilized sample of the strain with two copies of relevant reprint(s) if available within one month of the date of the acceptance.
- (4) If the state of the strain sent does not coincide with the description of the Strain Deposit Request Form, or do not meet any of the rules described above, the acceptance for deposit is canceled. (The NIES reserves the right to refuse any deposit at its discretion.)

Strain Deposit Request Form

Director,
Microbial Culture Collection,
National Institute for Environmental Studies

Date:

Depositor's full name (underline the family name):

NIES use only

Date

Name

Number

Acceptance YES NO

Depositor's affiliation and address:

TEL:

FAX:

E-mail:

I wish to contribute the following microbial culture strain to the NIES-Collection.

Reason for deposit:

1. Scientific name with citation of authority

2. Strain designation or symbol and other collection number

3. History

a. Locality:

b. Habitat (select from Nos. in page 25): _____

c. Collection date:

d. Collector's full name (underline the family name):

e. Isolation date:

f. Isolator's full name (underline the family name):

g. Source of isolation: soil, sediment, water, animal (),
 plant (), snow or ice, others ()

h. Isolation objective: motile vegetative cell, nonmotile vegetative cell, dormant cell,
 spore, others ()

i. Physical separation: pipette washing, dilution, agar plating, taxis,
 others ()

j. Isolation treatment: none, antibiotics, ultra-violet irradiation, enrichment culture,
 chemicals (), heat, ultra-sonic, others ()

k. Identified by (write full name with underlined family name):

l. Axenified by (write full name with underlined family name):

m. Clonized by (write full name with underlined family name):

4. Status

- a. Microalgae: axenic, unialgal, clonal, mixed
b. Bacteria: pure, non-clonal
c. Protozoa: axenic, monoxenic, dixenic, mixed

5. Medium

- a. Designation and references:

b. Composition and notes for preparation of medium:

6. Experimental culture conditions

- a. Temperature:
b. Light intensity:
c. Light quality:
d. L/D cycle:

7. Stock-culture conditions

- a. Maintenance by sub-culturing
i. Temperature:
ii. Light intensity:
iii. Light quality:
iv. L/D cycle:
v. Duration:
b. Preservation in freezing: yes no
i. Cryoprotectant:
ii. Freezing rate:
iii. Thawing condition: 40°C waterbath others ()
iv. Preservation: liquid nitrogen Deep freezer (°C) others ()
c. Preservation in freeze-drying: yes no
d. Preservation in L-drying: yes no

8. Strain characteristics

- a. Environmental characteristics (select from Nos. in page 25): _____
b. Physiological and ecological characteristics (select from Nos. in page 25): _____
c. Miscellaneous characteristics (select from Nos. in page 25): _____

9. Gene data

- a. Gene name and accession No:
b. Registrar's full name (underline the family name):
c. Registration date:

10. Other information

11. References

Two copies of relevant reprint(s) should be accompanied with this form.

III. ORDERING AND DISTRIBUTION OF STRAINS

1. Distribution to researchers of this institute

- (1) Rules on distribution
 - i) Anyone who uses a NIES-Collection strain in a paper which is subsequently published, is requested to give the full number of the strain, e.g. NIES-125, and to send two copies of the reprint(s) or Xerox copies to the NIES-Collection.
 - ii) In order to prevent trouble, confusion, or difficulty in the collection, accumulation and processing of strain information and data, the distribution of any NIES-Collection strain to a third party is strictly prohibited.
- (2) Procedure for ordering strains
 - i) All orders for strains must be requested to the NIES-Collection by completing the Strain Ordering Form (p. 23).
 - ii) Upon receipt of a strain, the Strain Receipt Form (p. 24) should be completed and returned to the NIES-Collection as soon as possible.

2. Distribution to people of other organizations, both academic and commercial

The distribution of the strains is made through the Global Environmental Form (GEF), and the ordering procedure is shown in the GEF Catalogue (April, 2000).

3. Special warning for distribution of "Untransportable" and "[Cryopreserved]" strains

Orders for the strains designated as "Untransportable" in the strain description (see p. 42) may not be accepted, depending on the season or condition of the cultures.

In principle, such strains must be personally carried (e.g. as hand luggage) in order to be transported overseas. Such transport must be arranged by individual requestors.

And for the strains designated as "[Cryopreserved]" in the strain description (see p. 42), frozen cells are thawed and inoculated into the fresh medium just after the order is accepted. As a result, it takes at least one month for the overseas shipping of these strains.

Strain Ordering Form

NIES use only
Date
Name
Number

Director,
Microbial Culture Collection,
National Institute for Environmental Studies

Date:

Requestor's full name (underline the family name):

Requestor's affiliation and address:

TEL:

FAX:

E-mail:

The following microbial culture strains are requested.

Scientific names and strain numbers:

Object of use (in detail):

Strain data

Needed (strain number)

Not needed

Strain Receipt Form

Director,
Microbial Culture Collection,
National Institute for Environmental Studies

NIES use only

Date

Name

Number

Date:

Recipient's full name (underline the family name):

Recipient's affiliation and address:

TEL:

FAX:

Date of strain receipt :

I received the following culture strains.

Scientific names and strain numbers:

States of strains received:

Good (strain number)

Poor (strain number)

Others (strain number)

Comments:

--- **Habitat** (生息環境) ---

- 1) Freshwater (淡水)
- 1-a) Oligotrophic (貧栄養) 1-b) Mesotrophic (中栄養) 1-c) Eutrophic (富栄養) 1-d) Dystrophic (腐植栄養)
- 2) Marine (海水)
- 2-a) Coastal (沿岸) 2-b) Oceanic (外洋) 2-c) Surface (表層) 2-d) Deep (深層, 採水深度をお書きください)
- 3) Brackish (汽水) 4) Tide pool (潮だまり) 5) Tideland (干潟)
- 6) Salt water (塩水) 7) Soil (土) 8) Hot spring (温泉)
- 9) Cold spring (鉱泉) 10) Snow or ice (雪または氷) 11) Wetland (湿原, 湿地)
- 12) Rice field (水田) 13) Lotic (流水) 14) Lentic (止水)
- 15) Others (その他) {write details (お書きください)}

--- **Environmental Characteristics** (環境上問題となる特性) ---

- 1) Red tide (赤潮) 2) Water bloom (水の華)
- 3) AGP (藻類生長試験) 4) Oxidation pond (酸化池)
- 5) Biodegradation (生分解) 6) Activated sludge (活性汚泥)
- 7) Microbial film process (生物膜処理) 8) Indicator (指標)
- 9) Predator (捕食性) 10) Offensive taste and odor (異味異臭)
- 11) Toxic (有毒) 12) Inhibition of purification (浄水障害)
- 13) Corrosion (腐食性) 14) Others (その他) {write details (お書きください)}

--- **Physiological and Ecological Characteristics** (生理生態的特性) ---

- 1) Autotrophic (独立栄養) 2) Mixotrophic (混合栄養)
- 3) Phagotrophic (摂食栄養) 4) Heterotrophic (従属栄養)
- 5) Planktonic (浮遊性) 6) Benthic (底生)
- 7) Symbiotic (共生) 8) Parasitic (寄生)
- 9) Saprophytic (腐生) 10) Endophytic (内生)
- 11) Epiphytic (植物着生) 12) Epilithic (岩石着生)
- 13) Eurythermal (広温性) 14) Stenothermal (狭温性)
- 15) Thermophilic (好熱性) 16) Psychrophilic (好冷性)
- 17) Euryhaline (広塩性) 18) Stenohaline (狭塩性)
- 19) Halophilic (好塩性) 20) Acidophilic (好酸性)
- 21) Sun-type (陽生型) 22) Shade-type (陰生型)
- 23) Nitrogen fixation (窒素固定) 24) Fermentation (発酵)
- 25) Bioluminescence (生物発光) 26) Phototaxis (走光性)
- 27) Hydrogen evolution (水素発生) 28) Aerobic (好気性)
- 29) Anaerobic (嫌気性) 30) Gram positive (グラム陽性)
- 31) Gram negative (グラム陰性) 32) Others (その他) {write details (お書きください)}

--- **Miscellaneous Characteristics** (その他の特性) ---

- 1) Mutant (突然変異株) 2) Type strain (タイプ株)
- 3) Heterothallic (雌雄異株) 4) Homothallic (雌雄同株)
- 5) Dioecious (雌雄異体) 6) Monoecious (雌雄同体)
- 7) Isogamy (同型配偶) 8) Anisogamy (異型配偶)
- 9) Oogamy (卵生殖) 10) H, h type (H, h型生活環)
- 11) H, d type (H, d型生活環) 12) D, d+h type (D, d+h型生活環)
- 13) Polyploidy (倍数性株) 14) Chromatic adaptation (色順応)
- 15) Rod (桿菌) 16) Coccus (球菌)
- 17) Coryne form (コリネ形菌) 18) Spiral (らせん菌)
- 19) Motile (運動性) 20) Immotile (非運動性)
- 21) Resting spore forming (休眠孢子形成) 22) Resting spore not forming (休眠孢子非形成)
- 23) Mating type + (交配型 +) 24) Mating type - (交配型 -)
- 25) Mating type female (交配型 雌) 26) Mating type male (交配型 雄)
- 27) Others (その他) {write details (お書きください)}

IV. ESTABLISHMENT OF FRESH CULTURES

When investigators are to receive culture strains, the following steps should be carried out to establish fresh cultures.

- i) Appropriate culture media should be prepared before receipt of the strains according to the recipes given in Chap. VI and with reference to the basic methods given in Chap. V.
- ii) Immediately after receipt, cultures should be unpacked, transferred to new media and grown at the temperature and light intensity directed by the Collection (cf. Chap. VIII); the light-dark cycle should be 12 hours light : 12 hours dark, and the screw-cap on the tube should be loosened.
- iii) After detecting good growth, further maintenance of cultures requires transfer into new media at intervals suggested by the Collection (cf. Chap. VIII).

V. BASIC METHODS FOR PREPARATION OF ALGAL CULTURE MEDIA

A number of media are used for maintenance of algal cultures and prepared according to the recipes given in the next chapter. The present chapter introduces the basic methods for preparation adopted in the Global Environmental Forum.

1. Stock solutions

Media are generally composed of three components, macronutrients, trace metals and vitamins (cf. Table 2) and prepared from stock solutions of these components. The concentration of stock solutions of trace metals and vitamins is very low and primary stock solutions are prepared for dilution to obtain the stock solutions.

A. Macronutrients : Separate stock solutions with a concentration of 10 mg/ml of each macronutrient are prepared and stored in a refrigerator (5°C).

B. Trace metals : These elements are prepared either as separate stock solutions or mixed stock solutions.

(1) Separate stock solutions

- i) Prepare a separate primary solution with a concentration of 10-100 mg/ml.
- ii) Dilute each primary solution to prepare the stock solution with a concentration of 1 mg/ml and store in a refrigerator (5°C).

Table 2. Chemicals employed for culture media

Macronutrients	Trace metals
NaCl	H ₃ BO ₃
KCl	MnCl ₂ ·4H ₂ O
CaCl ₂ ·2H ₂ O	MnSO ₄ ·7H ₂ O
MgCl ₂ ·6H ₂ O	FeCl ₃ ·6H ₂ O
Na ₂ SO ₄	FeSO ₄ ·7H ₂ O
K ₂ SO ₄	CoCl ₂ ·6H ₂ O
MgSO ₄ ·7H ₂ O	ZnSO ₄ ·7H ₂ O
NaNO ₃	CuSO ₄ ·5H ₂ O
KNO ₃	Na ₂ MoO ₄ ·2H ₂ O
Ca(NO ₃) ₂ ·4H ₂ O	Vitamins
NH ₄ NO ₃	Vitamin B ₁₂
NaH ₂ PO ₄ ·2H ₂ O	Biotin
β -Na ₂ glycerophosphate·5H ₂ O	Thiamine HCl
KH ₂ PO ₄	Nicotinic acid
K ₂ HPO ₄	Calcium panthothenate
Na ₂ CO ₃	<i>p</i> -Aminobenzoic acid
NaHCO ₃	Inositol
Na ₂ SiO ₃ ·9H ₂ O	Folic acid
	Thymine

(2) Mixed stock solution

- i) Same as (1)-i)
- ii) Add approximately 80 % of the required volume of distilled water to a beaker.
- iii) Dissolve the required amount of Na₂EDTA, while stirring continuously.
- iv) Dispense the required volume of each trace metal from primary solution, while stirring continuously.
- v) Dilute to final volume with distilled water and store in a refrigerator (5°C).

C. Vitamins : Only three vitamins, vitamin B₁₂, biotin, and thiamine HCl have been found necessary for growth of many microalgae and are added to most media. Some media, in addition, contain other vitamins

(1) Vitamin B₁₂, biotin and thiamine HCl

- i) Prepare separate primary stock solution with a concentration of 0.1 mg/ml of vitamin B₁₂ and biotin and 10 mg/ml of thiamine HCl.
- ii) After dispersing 1 ml of these primary stock solution into each of a number of test tubes and autoclaving (121°C, 20 min), store in a freezer at -20°C.
- iii) Thaw and dilute 1 ml of primary stock solution of each vitamins to prepare the working stock solution with a concentration of 1 µg/ml of vitamin B₁₂ and biotin or of 100 µg/ml of thiamine HCl, and store in a refrigerator (5°C).

(2) Other vitamins: Additional vitamins are added to some media in the forms of mixes (cf. Chap. VI-56). It is recommended to prepare a large volume of mixed stock solution.

- i) Prepare a separate primary solution with a concentration of 0.1-1.0 mg/ml.
- ii) Add approximately 80 % of the required volume of distilled water to a beaker.
- iii) Dispense the required volume of each vitamin from the primary solution, while stirring continuously.
- iv) After sterilization by passing through a Millipore filter (0.22 μm), aseptically dispense 100 ml of the mixed stock solution into each of a number of vessels and store in a freezer at -20°C .

2. Media

Media are divided broadly into two categories, synthetic and enriched. The former are used for maintenance of all freshwater algal cultures and some marine ones and the latter for most marine ones.

(1) Synthetic medium (freshwater)

- i) Add approximately 80-90% of the required volume of distilled water to a beaker.
- ii) Dissolve appropriate quantities of weighed buffer such as Tris (hydroxymethyl) aminomethane (known as Tris), glycylglycine, HEPES, TAPS, Bicine, MES or 1, 2, 3, 4-cyclopentan tetracarboxylic acid (if required), while stirring continuously. These buffers are easily soluble with stirring.
- iii) Dispense the appropriate nutrients from previously prepared stock solutions, while stirring continuously.
- iv) Dilute to final volume with distilled water.
- v) Check the pH and make any adjustments with either 1N HCl or 1N NaOH (if buffers required) or with either 0.1N HCl or 0.1N NaOH (if no buffers required).
- vi) Dispense 10 ml of medium into each of the test tube (18 \times 150mm) and sterilize by autoclaving (121°C , 20 min).

(2) Synthetic medium (marine)

- i) Add approximately 80% of the required volume of distilled water to a beaker.
- ii) Dissolve appropriate quantities of weighed Tris, Nitritotriacetic acid (known as NTA) and major salts such as NaCl, $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, KCl and $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$, while stirring continuously.
- iii) Dispense the other nutrients from previously prepared stock solutions.
- iv) Dilute to the final volume with the distilled water.
- v) Check the pH, which is usually adjusted to 8.0 with 1N HCl.
- vi) Dispense 10 ml of medium into each of the test tubes and sterilize by autoclaving (121°C , 20 min).

(3) Enriched seawater medium

- i) Collect offshore water free from gross pollution and remove particulate matter with Whatman GF/C filters.
- ii) Check the salinity. A salinity of 35‰ is considered normal seawater.
- iii) Add approximately 80-90% of the required volume of seawater to a beaker.
- iv) Dissolve appropriate quantities of weighed Tris (if required).
- v) Dispense the appropriate nutrients from previously prepared stock solutions.
- vi) Dilute to the final volume with seawater.
- vii) Check the pH and adjust to 8.0 with 1N HCl if necessary.
- viii) Dispense 10 ml of medium into each test tube and sterilize by autoclaving (121°C, 20 min).

3. Agar slant

Agar is added usually at concentrations of 1.5% after liquid medium has been prepared, prior to autoclaving.

- i) Add the appropriate quantities of weighed agar to liquid medium and heat to 121°C by autoclaving to melt all the agar.
- ii) After melting, quickly dispense 10 ml of agar medium into each test tube and sterilize by autoclaving (121°C, 20 min).
- iii) After sterilization, lay the upper part of the test-tube on a rod (1 cmφ) and cool to form an agar slant.

VI. MEDIA

1) Stock media for algae

1)-1. For freshwater algae

1. AF-6 (130)¹⁾

NaNO ₃	14	mg
NH ₄ NO ₃	2.2	mg
MgSO ₄ •7H ₂ O	3	mg
KH ₂ PO ₄	1	mg
K ₂ HPO ₄	0.5	mg
CaCl ₂ •2H ₂ O	1	mg
CaCO ₃ ²⁾	1	mg
Fe-citrate	0.2	mg
Citric acid	0.2	mg
Biotin	0.2	µg
Thiamine HCl	1	µg
Vitamin B ₆	0.1	µg
Vitamin B ₁₂	0.1	µg
Trace metals ²⁾	0.5	ml
Distilled water	99.5	ml
pH 6.6 ³⁾		

1) Reference number in parentheses.

2) In the NIES-Collection, CaCO₃ is removed and PIV metals are used instead of trace metals.

3) pH is adjusted to 6.6 by buffering with 40 mg MES in the NIES-Collection.

2. AF-6 / 2

AF-6 medium is diluted with distilled water to 1 / 2.

3. Allen (1)

(NH ₄) ₂ SO ₄	132	mg
KH ₂ PO ₄	27.2	mg
MgSO ₄ •7H ₂ O	24.6	mg
CaCl ₂ •2H ₂ O	7.4	mg
Allen Metals ¹⁾	0.01	ml
Distilled water	99.9	ml
pH 2.5 ²⁾		

1) See 48

2) pH is adjusted to 2.5 with 1 N H₂SO₄.

4. C (75)

Ca(NO ₃) ₂ •4H ₂ O	15	mg
KNO ₃	10	mg
β-Na ₂ glycerophosphate•5H ₂ O	5	mg
MgSO ₄ •7H ₂ O	4	mg
Vitamin B ₁₂	0.01	µg
Biotin	0.01	µg
Thiamine HCl	1	µg
PIV metals ¹⁾	0.3	ml
Tris (hydroxymethyl) aminomethane	50	mg
Distilled water	99.7	ml
pH 7.5		

1) See 54

5. CA (86)

Ca(NO ₃) ₂ •4H ₂ O	2	mg
KNO ₃	10	mg
NH ₄ NO ₃	5	mg
β-Na ₂ glycerophosphate•5H ₂ O	3	mg
MgSO ₄ •7H ₂ O	2	mg
Vitamin B ₁₂	0.01	µg
Biotin	0.01	µg
Thiamine HCl	1	µg
PIV metals ¹⁾	0.1	ml
Fe (as EDTA; 1:1 molar) ²⁾	0.1	mg
HEPES	40	mg
Distilled water	99.9	ml

pH 7.2

1) See 54

2) See 50

6. CAM

CA medium with pH adjusted to 6.5 by buffering with MES instead of HEPES.

7. Carefoot (13)

NaNO ₃	24.7	mg
CaCl ₂ •2H ₂ O	1.1	mg
MgSO ₄ •7H ₂ O	4.7	mg
K ₂ HPO ₄	0.9	mg
KH ₂ PO ₄	2.3	mg
NaCl	1.5	mg
PIV metals ¹⁾	0.5	ml
Distilled water	99.5	ml

pH 7.5

* In the NIES-Collection, 0.02 µg Vitamin B₁₂, 0.02 µg Biotin and 2 µg Thiamine HCl are added to this medium.

1) See 54

8. CB

C medium with pH adjusted to 9.0 by buffering with Bicine instead of Tris (hydroxymethyl) aminomethane.

9. CC (80)

C medium with pH adjusted to 3.0 by buffering with 1, 2, 3, 4 - cyclopentan tetracarboxylic acid instead of Tris (hydroxymethyl) aminomethane.

10. CSi

C medium with pH adjusted to 7.0 by buffering with 50 mg HEPES instead of Tris (hydroxymethyl) aminomethane. Thereafter, 10 mg Na₂SiO₃•9H₂O is added.

11. CSi+Cu

0.25 mg CuSO₄•5H₂O is added to CSi medium.

12. CT (391)

C medium with pH adjusted to 8.2 by buffering with 40 mg TAPS instead of Tris (hydroxymethyl) aminomethane.

13. CYT

10 mg Yeast extract and 20 mg Tryptone are added to C medium.

14. HUT (74)

KH ₂ PO ₄	2	mg
MgSO ₄ ·7H ₂ O	2.5	mg
Sodium acetate	40	mg
Potassium citrate	4	mg
Polypeptone	60	mg
Yeast extract	40	mg
Vitamin B ₁₂	0.05	µg
Thiamine HCl	0.04	mg
Distilled water	100	ml
pH 6.4		

* Add 150 mg agar to 100 ml of the medium for semi-solid medium.

15. M-11 (51), (421)

NaNO ₃	10	mg
K ₂ HPO ₄	1	mg
MgSO ₄ ·7H ₂ O	7.5	mg
CaCl ₂ ·2H ₂ O	4	mg
Na ₂ CO ₃	3	mg
FeSO ₄ ·7H ₂ O	0.1	mg
Na ₂ EDTA·2H ₂ O	0.1	mg
Distilled water	100	ml
pH 8.0		

16. MA (77)

Ca(NO ₃) ₂ ·4H ₂ O	5	mg
KNO ₃	10	mg
NaNO ₃	5	mg
Na ₂ SO ₄	4	mg
MgCl ₂ ·6H ₂ O	5	mg
β-Na ₂ glycerophosphate·5H ₂ O	10	mg
Na ₂ EDTA	0.5	mg
FeCl ₃ ·6H ₂ O	0.05	mg
MnCl ₂ ·4H ₂ O	0.5	mg
ZnCl ₂	0.05	mg
CoCl ₂ ·6H ₂ O	0.5	mg
Na ₂ MoO ₄ ·2H ₂ O	0.08	mg
H ₃ BO ₃	2	mg
Bicine	50	mg
Distilled water	100	ml
pH 8.6		

17. MAF-6

10 mg glucose and 10 mg sodium acetate are added to AF-6 medium.

18. M Chu No. 10 (15)

Ca(NO ₃) ₂ ·4H ₂ O	2.0	mg
KH ₂ PO ₄	0.62	mg
MgSO ₄ ·7H ₂ O	2.5	mg
Na ₂ CO ₃	2	mg
Na ₂ SiO ₃ ·9H ₂ O	2.5	mg
HCl (1N) ¹⁾	0.025	ml
Na ₂ EDTA·2H ₂ O	0.2	mg
FeCl ₃ ·6H ₂ O	0.1	mg
H ₃ BO ₃	0.248	mg
MnCl ₂ ·4H ₂ O	0.139	mg
(NH ₄) ₆ Mo ₇ O ₂₄ ·4H ₂ O	0.1	mg
Vitamin B ₁₂	1	µg
Thiamine HCl	0.1	µg
Biotin	0.1	µg
Distilled water	100	ml

1) In the NIES-Collection, pH is adjusted to 7.6 with respective volume of 1 N HCl.

19. MDM (366)

KNO ₃	100	mg
MgSO ₄ ·7H ₂ O	25	mg
K ₂ HPO ₄	25	mg
NaCl	10	mg
CaCl ₂ ·2H ₂ O	1	mg
Fe solution ¹⁾	0.1	ml
A ₅ solution ²⁾	0.1	ml
Agar	1.5	g
Distilled water	99.8	ml
pH 8.0		

1) See 51

2) See 47

20. MG (76)

Ca(NO ₃) ₂ •4H ₂ O	2	mg
KNO ₃	10	mg
β-Na ₂ glycerophosphate•5H ₂ O	3	mg
MgSO ₄ •7H ₂ O	2	mg
Vitamin B ₁₂	0.01	µg
Biotin	0.01	µg
Thiamine HCl	1	µg
PIV metals ¹⁾	0.1	ml
Fe (as EDTA; 1:1 molar) ²⁾	0.1	ml
HEPES	40	mg
Distilled water	99.9	ml
pH 7.2		

1) See 54

2) See 50

21. MGM

MG medium with pH adjusted to 6.5 by buffering with MES instead of HEPES.

22. P 35 (77)

NH ₄ NO ₃	10	mg
MgSO ₄ •7H ₂ O	4	mg
KCl	5	mg
CaCl ₂ •2H ₂ O	7.4	mg
β-Na ₂ glycerophosphate•5H ₂ O	5	mg
Sodium acetate	100	mg
Vitamin B ₁₂	0.01	µg
Biotin	0.01	µg
Thiamine HCl	1	µg
PIV metals ¹⁾	0.3	ml
Tris (hydroxymethyl) aminomethane	50	mg
Distilled water	99.7	ml
pH 8.0		

1) See 54

23. MW (289)

Urea	0.85	mg
NaNO ₃	0.17	mg
NH ₄ Cl	0.042	mg
Ca(NO ₃) ₂ •4H ₂ O	10	mg
CaCO ₃	1	mg
CaCl ₂ •2H ₂ O	1.4	mg
KNO ₃	1	mg
KHCO ₃	0.9	mg
β-Na ₂ glycerophosphate•5H ₂ O	2	mg
MgSO ₄ •7H ₂ O	1.5	mg
PIV metals ¹⁾	0.05	ml
Vitamin B ₁₂	0.02	µg
Thiamine HCl	2	µg
Biotin	0.02	µg
Glycylglycine	10	mg
Distilled water	99.95	ml
pH 7.2		

1) See 54

24. MW / 5

MW medium is diluted with distilled water to 1 / 5.

25. URO (136), (196)

NH ₄ NO ₃	0.5	mg
β-Na ₂ glycerophosphate•5H ₂ O	0.4	mg
MgSO ₄ •7H ₂ O	1	mg
CaCl ₂ •2H ₂ O	1	mg
KCl	0.1	mg
Thiamine HCl	1	µg
Vitamin B ₁₂	0.01	µg
Biotin	0.01	µg
Fe-EDTA	0.05	mg
PIV metals ¹⁾	0.1	ml
Distilled water	99.9	ml
pH 7.5 ²⁾		

1) See 54

2) pH is adjusted to 7.5 with 0.1 N HCl.

26. VT (286)

Ca(NO ₃) ₂ •4H ₂ O	11.78	mg
β-Na ₂ glycerophosphate•5H ₂ O	5	mg
MgSO ₄ •7H ₂ O	4	mg
KCl	5	mg
Vitamin B ₁₂	0.01	μg
Biotin	0.01	μg
Thiamine HCl	1	μg
PIV metals ¹⁾	0.3	ml
Glycylglycine	50	mg
Distilled water	99.7	ml
pH 7.5		

1) See 54

27. VTAC (250)

20 mg sodium acetate is added to VT medium.

28. VTYT (80)

10 mg yeast extract and 20 mg tryptone are added to VT medium.

29. W (388)

Ca(NO ₃) ₂ •4H ₂ O	10	mg
KNO ₃	1	mg
MgSO ₄ •7H ₂ O	1.5	mg
β-Na ₂ glycerophosphate•5H ₂ O	2	mg
Urea	1.7	mg
Thiamine HCl	0.2	μg
Vitamin B ₁₂	0.002	μg
Biotin	0.002	μg
PIV metals ¹⁾	0.05	ml
Glycylglycine	10	mg
Distilled water	99.95	ml
pH 7.5		

1) See 54

30. SW (282)

A small amount of dried soil is put into a test tube, and 20 ml distilled water is added.

31. SOT (258)

NaHCO ₃	1.68	g
K ₂ HPO ₄	50	mg
NaNO ₃	250	mg
K ₂ SO ₄	100	mg
NaCl	100	mg
MgSO ₄ •7H ₂ O	20	mg
CaCl ₂ •2H ₂ O	4	mg
FeSO ₄ •7H ₂ O	1	mg
Na ₂ EDTA	8	mg
A ₅ solution ¹⁾	0.1	ml
Distilled water	99.9	ml

1) See 47

1)-2. For marine algae

32. ESM (264)

NaNO ₃	12	mg
K ₂ HPO ₄	0.5	mg
Vitamin B ₁₂	0.1	µg
Biotin	0.1	µg
Thiamine HCl	10	µg
Fe-EDTA	25.9	µg
Mn-EDTA	33.2	µg
Tris (hydroxymethyl) aminomethane	100	mg
Soil extract ¹⁾	5	ml
Sea water	95	ml
pH 8.0		

1) See 57

33. f / 2 (50)

NaNO ₃	7.5	mg
NaH ₂ PO ₄ · 2H ₂ O	0.6	mg
Vitamin B ₁₂	0.05	µg
Biotin	0.05	µg
Thiamine HCl	10	µg
Na ₂ SiO ₃ · 9H ₂ O	1	mg
f / 2 metals ¹⁾	0.1	ml
Sea water	99.9	ml

1) See 52

34. M-ASP7 (409)

NaCl	2.5	g
MgSO ₄ · 7H ₂ O	900	mg
KCl	70	mg
CaCl ₂ · 2H ₂ O	30	mg
NaNO ₃	5	mg
NaH ₂ PO ₄ · 2H ₂ O	2	mg
Vitamin B ₁₂	0.1	µg
Vitamin mix S ₃ ¹⁾	1	ml
Na ₂ SiO ₃ · 9H ₂ O	1	mg
P _N metals ²⁾	3	ml
Tris (hydroxymethyl) aminomethane	100	mg
NTA	7	mg
Distilled water	96	ml
pH 8.0		

1) See 56

2) See 55

35. MF

f / 2 medium with Na₂SiO₃ · 9H₂O replaced by 1.0ml soil extract¹⁾ and adjusted to pH 8.0 by buffering with 100mg Tris (hydroxymethyl) aminomethane.

1) See 57

36. MKM (366)

KNO ₃	75	mg
KH ₂ PO ₄	2.5	mg
MgSO ₄ · 7H ₂ O	2	mg
Fe-citrate	250	µg
Agar	1.5	g
Sea water	50	ml
Distilled water	50	ml

37. WESM

ESM medium with 95 ml sea water replaced by 85 ml sea water and 10 ml distilled water.

2) Bacteria-free check media

2)-1. For freshwater algae

38. YT (80)

Stock medium	100	ml
Yeast extract	100	mg
Tryptone	200	mg

39. B - I (87)

Stock medium	100	ml
Proteose peptone	100	mg

40. B - II (87)

Stock medium	100	ml
Yeast extract	500	mg

41. B - III (87)

Stock medium	100	ml
Peptone	500	mg
Beef extract	300	mg

42. B - IV (87)

Stock medium	100	ml
Glucose	100	mg
Peptone	100	mg

43. B - V (87)

Stock medium	100	ml
Sodium acetate	50	mg
Glucose	50	mg
Tryptone	50	mg
Yeast extract	30	mg

2)-2. For marine algae

44. STP (285)

NaNO ₃	20	mg
K ₂ HPO ₄	1	mg
Sodium glutamate	50	mg
Glucose	20	mg
Glycine	10	mg
D, L - Alanine	10	mg
Vitamin mix 8 ¹⁾	0.1	ml
Trypticase	20	mg
Yeast autolysate ²⁾	20	mg
Sucrose	100	mg
Soil extract ³⁾	5	ml
Sea water	80	ml
Distilled water	15	ml

pH 7.5

1) In the NIES-Collection, vitamin mix 8 is replaced by Vitamin mix S₃.

2) In the NIES-Collection, yeast autolysate is replaced by yeast extract.

3) . See 57

45. MM23 (M. Tatewaki, pers. comm.)

NaCl	1.8	g
MgSO ₄ •7H ₂ O	500	mg
KCl	60	mg
NaNO ₃	100	mg
CaCl ₂ •2H ₂ O	36.7	mg
K ₂ HPO ₄	6	mg
Sucrose	400	mg
PII metals ¹⁾	2	ml
FeCl ₃ •6H ₂ O	48	µg
Thiamine HCl	10	µg
Biotin	0.1	µg
Vitamin B ₁₂	0.2	µg
C-Source Mix II ²⁾	1	ml
Tris (hydroxymethyl) aminomethane	100	mg
Distilled water	97	ml

pH 8.0

1) See 53

2) See 49

46. Bf / 2 (437)

ASP7 ¹⁾	100	ml
Trypticase	50	mg
Yeast extract	5	mg

1) In the NIES-Collection, ASP7 is replaced by f / 2 medium.

3) Trace metals, vitamin mixes and soil extract**47. A₅ solution (71)**

H ₃ BO ₃	286	mg
MnSO ₄ ·7H ₂ O	250	mg
ZnSO ₄ ·7H ₂ O	22.2	mg
CuSO ₄ ·5H ₂ O	7.9	mg
Na ₂ MoO ₄ ·2H ₂ O	2.1	mg
Distilled water	100	ml

48. Allen metals (1)

Fe-EDTA	30.16	mg
MnCl ₂ ·4H ₂ O	1.79	mg
H ₃ BO ₃	2.86	mg
ZnSO ₄ ·7H ₂ O	0.22	mg
CuSO ₄ ·5H ₂ O	0.079	mg
(NH ₄) ₆ MoO ₂₄ ·4H ₂ O	0.13	mg
NH ₄ VO ₃	0.023	mg
Distilled water	100	ml

49. C - Source Mix II (M. Tatewaki, pers. comm.)

Glycine	100	mg
D, L - Alanine	100	mg
L - Asparagine	100	mg
Sodium acetate·3H ₂ O	200	mg
Glucose	200	mg
L - Glutamic acid	200	mg
Distilled water	100	ml

50. Fe (as EDTA; 1:1 molar) (284)

Fe(NH ₄) ₂ (SO ₄) ₂ ·6H ₂ O	70.2	mg
Na ₂ EDTA·2H ₂ O	66	mg
Distilled water	100	ml

* 1 ml of this solution contains 0.1 mg Fe.

51. Fe solution (80)

FeSO ₄ ·7H ₂ O	200	mg
Distilled water	100	ml
Conc·H ₂ SO ₄	0.026	ml ¹⁾

1) 2 drops / 500ml (Ref. 80).

52. f / 2 metals (50)

Na ₂ EDTA·2H ₂ O	440	mg
FeCl ₃ ·6H ₂ O	316	mg
CoSO ₄ ·7H ₂ O	1.2	mg
ZnSO ₄ ·7H ₂ O	2.1	mg
MnCl ₂ ·4H ₂ O	18	mg
CuSO ₄ ·5H ₂ O	0.7	mg
Na ₂ MoO ₄ ·2H ₂ O	0.7	mg
Distilled water	100	ml

53. P II metals (283)

H ₃ BO ₃	114	mg
FeCl ₃ ·6H ₂ O	4.9	mg
MnSO ₄ ·4H ₂ O	16.4	mg
ZnSO ₄ ·7H ₂ O	2.2	mg
CoSO ₄ ·7H ₂ O	480	μg
Na ₂ EDTA·2H ₂ O	100	mg
Distilled water	100	ml

54. P IV metals (286)

FeCl ₃ •6H ₂ O	19.6	mg
MnCl ₂ •4H ₂ O	3.6	mg
ZnSO ₄ •7H ₂ O ¹⁾	2.2	mg
CoCl ₂ •6H ₂ O	0.4	mg
Na ₂ MoO ₄ •2H ₂ O	0.25	mg
Na ₂ EDTA•2H ₂ O	100	mg
Distilled water	100	ml

1) In NIES-Collectoin, ZnCl₂ is replaced by ZnSO₄•7H₂O.

55. P_N metals (409)

Na ₂ EDTA•2H ₂ O	100	mg
H ₃ BO ₃	113	mg
FeCl ₃ •6H ₂ O	6.3	mg
CoSO ₄ •7H ₂ O	0.093	mg
ZnSO ₄ •7H ₂ O	4.66	mg
MnCl ₂ •4H ₂ O	3.2	mg
Distilled water	100	ml

56. Vitamine mix S₃ (283)

Thiamine HCl	5	mg
Nicotinic acid	1	mg
Calcium pantothenate	1	mg
<i>p</i> - Aminobenzoic acid	0.1	mg
Biotin	0.01	mg
Inositol	50	mg
Folic acid	0.02	mg
Thymine	30	mg
Distilled water	100	ml

57. Soil extract (285)

100g soil combined with 100ml distilled water is heated for 2h and then cooled. The supernatant is passed through a GF / C filter and then distilled water added until there is a total of 100ml.

4) Stock medium for protozoa

58. LE

L Solution: White part of lettuce is dried at 90 °C for 16 - 18 h without scorching. 300 mg of the dried lettuce is added to 100 ml boiling water (9 : 1 distilled water / tap water) and boiled for 30 minutes, while stirring. The supernatant is passed through cottonwool.

E solution: 300 mg of crushed yolk of hardboiled egg is added to 100ml water (9 : 1 distilled water / tap water) and boiled for 30 minutes, while stirring. The supernatant is passed through cottonwool.

Equal quantities of L and E solutions are mixed. The pH is adjusted to 6.8 - 7.0 with 1 N NaOH. 100 ml of the solution is dispensed into each 200ml-Erlenmayer flasks and sterilized by autoclaving (121°C, 15 min).

VII. 保存株データの利用法

系統保存株の学名はアルファベット順に並べてあり、学名が同じ場合は株番号順に並べてある。同定者が記載されていない限り、学名は原則として分離者によってつけられたものである。また、株番号は、数字の前に NIES- をつけて使用することとし(例:NIES-1)、株の学名が命名法などの変更で変わった場合や、やむをえない理由で株が消失した場合にも変更したり付け変えたりしないものとする。

個々の項目についての説明は下記の例を参照されたい。

Spirulina platensis (Gomont) Geitler¹⁾

Syn. *Arthrospira platensis* Gomont²⁾

45³⁾

Lake Kasumigaura / Ibaraki⁴⁾ (1975-11)⁵⁾

IAM M-184⁶⁾, Unialgal, Clonal⁷⁾, M.M.Watanabe⁸⁾ (1975-11)⁹⁾

Identified by: M.M.Watanabe¹⁰⁾

Culture conditions: MA, 25° C, 24 μE / m² sec, 1M, [Cryopreserved]¹¹⁾

Characteristics: Water bloom, Freshwater,

Forming water bloom in Inbanuma¹²⁾

KAS-6-50¹³⁾

References: 80, 335, 387, 391, 399, 417¹⁴⁾

- 1) 学名と原著者名：原著者名は学名の後に記した。
- 2) 異名。
- 3) 株番号：数字の前にNIES-を付けて使用すること。
- 4) 採集地。
- 5) 採集年月。
- 6) 他の保存機関に保存されている場合の株番号。保存機関名は略号で株番号の前に記されている。
 - IAM : 東京大学分子細胞生物学研究所
 - TAC : 国立科学博物館筑波実験植物園
 - CCAP : 英国 CCAPカルチャーコレクション
 - NIVA : ノルウェー 水界研究所藻類株保存施設
 - SAG : ドイツ ゲッチンゲン大学藻類株保存施設
 - UTEX : 米国 テキサス大学藻類株保存施設
- 7) 株の状態。

Axenic の表示があるものは無菌株である。

- 8) 分離者。
- 9) 分離年月。
- 10) 同定者。
- 11) 保存条件。培地名、保存温度、保存光強度、継代周期の順である。本施設では明暗周期は12時間明期／12時間暗期に設定されている。培地は特に記さない限り液体である。軟寒天培地：SS、寒天斜面培地：Sの場合は略号を（ ）内に記した。また（ ）内の温度および光強度は前培養が必要な場合、その条件である。なお光強度の表記については、本第6版より $\mu\text{E}/\text{m}^2\text{sec}$ を用いるものとする。現在凍結保存中の株については【Cryopreserved】と記した。
- 12) 株の性質。
Unstable; 保存状態が不安定で永続的な維持が困難である株。
Untransportable; 長時間の（航空便での）郵送では、生存状態で受け取るのが困難である株。
- 13) 分離者等の使用している株名。
- 14) 参考文献の番号。

なお、第IX章 2. 分類群別索引（125～134頁）における藻類門・綱の分類および配列は千原光雄（編）「藻類の多様性と系統」（バイオダイバーシティ・シリーズ、裳華房 1999）に掲載された分類表にしたがった。

VII. EXPLANATORY NOTES ABOUT STRAIN DATA

The strains are listed by scientific names in alphabetical order. Strains with the same scientific name are arranged in order of their strain numbers. The scientific name of each strain was designated by the isolator, unless the identifier is described. The number assigned to the given strain remains the same, regardless of any change in nomenclature. The strain number should be used with the initials "NIES-" (e.g. NIES-1). A detailed example of a strain description is presented below.

Spirulina platensis (Gomont) Geitler¹⁾

Syn. *Arthrospira platensis* Gomont²⁾

45³⁾

Lake Kasumigaura / Ibaraki⁴⁾ (1975-11)⁵⁾

IAM M-184⁶⁾, Unialgal, Clonal⁷⁾, M.M.Watanabe⁸⁾ (1975-11)⁹⁾

Identified by: M.M.Watanabe¹⁰⁾

Culture conditions: MA, 25° C, 24 μE/m² sec, 1M, [Cryopreserved]¹¹⁾

Characteristics: Water bloom, Freshwater,

Forming water bloom in Inbanuma¹²⁾

KAS-6-50¹³⁾

References: 80, 335, 387, 391, 399, 417¹⁴⁾

- 1) Scientific name with authority.
- 2) Synonym.
- 3) Strain number (used with the initials "NIES-").
- 4) Collection site.
- 5) Collection date.
- 6) The strain designations in other culture collections or institutions. The following abbreviations are presented before the strain number.
IAM: Institute of Molecular and Cellular Biosciences, University of Tokyo.
TAC: Tsukuba Botanical Garden, National Science Museum.
CCAP: Culture Collection of Algae and Protozoa, U.K.
NIVA: Culture Collection of Algae, Norwegian Institute for Water Research.
SAG: Culture Collection of Algae at the University of Göttingen, Germany.
UTEX: Culture Collection of Algae at the University of Texas at Austin, U.S.A.
- 7) Status of the strain (Unialgal or Axenic, Clonal or Non-clonal).

- 8) Isolator.
- 9) Isolation date.
- 10) Identifier.
- 11) Culture condition for maintenance: medium *, temperature, light intensity * * and duration of subculturing * * *

The light-dark cycle is defined as 12 hours light 12 hours dark.

* Unless otherwise noted the phase of the medium is liquid.

The abbreviations in parentheses are SS for semi-solid and S for solid.

* * Light intensity is indicated as $\mu\text{E}/\text{m}^2\text{sec}$ in this edition.

* * * Preculture temperature and light intensity are given in parentheses when preculture is required.

" [Cryopreserved] " indicates that the strain is preserved as a frozen condition at present.

- 12) Characteristics of the strain.

"Unstable" indicates that the strain probably cannot be maintained indefinitely, for various reasons including unsuccessful induction of auxospore formation and germination in diatom.

"Untransportable" indicates that the strain is not robust enough to be sent by air mail, involving much time.

- 13) Strain designation given by the isolator.

- 14) Reference number. References corresponding to the numbers are listed in pp.135~158.

Special Note. Algal phyla and classes and assignment of strains to each taxon (as shown in Chap. IX. 2. Systematic Index (pp.125~134) are arranged according to the system in Chihara (Ed.), "Diversity and Evolution of Algae" (Shokabo, Tokyo, 1999).

VIII. STRAIN DATA

ALGAE

- Achnanthes longipes* Agardh
330
Kawazu / Shizuoka (1985-05)
Axenic, Clonal, T.Sawaguchi (1985-05)
Identified by: T.Sawaguchi
Culture conditions: f/2, 10° C, 25 μ E/m² sec, 2M
Characteristics: Marine
IMHB-5
Reference: 113
- Achnanthes minutissima* Kützing
71
Kosaka River / Akita (1983-04)
Axenic, Clonal, A.Yuri (1983-09)
Identified by: M.Mizuno
Culture conditions: CSi, 20° C, 40 μ E/m² sec, 1M
Characteristics: Indicator, Freshwater
A15-6
References: 277, 337, 338, 412, 413
- 407
Miyata River / Ibaraki (1987-05)
Unialgal, Non-clonal, F.Kasai (1987-06)
Identified by: N.Takamura
Culture conditions: CSi, 15° C, 15 μ E/m² sec, 2M
Characteristics: Freshwater
4st-0-8
Reference: 338
- 408
Ashio / Gunma (1987-08)
Unialgal, Clonal, F.Kasai (1987-09)
Identified by: M.Idei
Culture conditions: CSi, 15° C, 15 μ E/m² sec, 2M
Characteristics: Freshwater
AT5-23
Reference: 338
- 409
Ashio / Gunma (1987-08)
Unialgal, Clonal, F.Kasai (1987-08)
Identified by: M.Idei
Culture conditions: CSi, 15° C, 15 μ E/m² sec, 2M
Characteristics: Freshwater
- Ast-3-3
Reference: 338
- 410
Ashio / Gunma (1987-08)
Unialgal, Non-clonal, F.Kasai (1987-09)
Identified by: N.Takamura
Culture conditions: CSi, 15° C, 15 μ E/m² sec, 2M
Characteristics: Freshwater
AT4-18
Reference: 338
- 411
Miyata River / Ibaraki (1987-02)
Unialgal, Non-clonal, F.Kasai (1987-03)
Identified by: N.Takamura
Culture conditions: CSi, 15° C, 15 μ E/m² sec, 2M
Characteristics: Freshwater
1st-3-17
References: 337, 338
- 412
Miyata River / Ibaraki (1987-02)
Unialgal, Non-clonal, F.Kasai (1987-03)
Identified by: N.Takamura
Culture conditions: CSi, 15° C, 15 μ E/m² sec, 2M
Characteristics: Freshwater
1St-1-1
References: 337, 338
- 413
Miyata River / Ibaraki (1987-02)
Unialgal, Non-clonal, F.Kasai (1987-03)
Identified by: N.Takamura
Culture conditions: CSi, 15° C, 15 μ E/m² sec, 2M
Characteristics: Freshwater
1st-2-8
References: 337, 338
- 414
Ooe River (Ozegahara) / Fukushima (1987-10)
Unialgal, Non-clonal, F.Kasai (1987-11)
Identified by: N.Takamura
Culture conditions: CSi, 15° C, 15 μ E/m² sec, 2M
Characteristics: Freshwater
0-25
Reference: 338

- Achnanthes minutissima* Kützing
var. *saprophila* Kobayasi et Mayama
372
Lake Kasumigaura / Ibaraki (1985-12)
Axenic, Clonal, T.Sawaguchi (1985-12)
Identified by: T.Sawaguchi
Culture conditions: CSI, M Chu No.10, 20° C,
40 µE/m² sec, 1M
Characteristics: Indicator, Freshwater,
Reidentified by M.Idei
KAAC-6
- Acinetospora crinita* (Carmichael) Sauvageau
548
Tuscan / Italy (1991)
Unialgal, Clonal, T.Hagiwara (1992)
Tentatively reidentified by: G.Sartoni
Culture conditions: f/2, 20° C, 4 µE/m² sec, 3M,
(20° C, 25 µE/m² sec)
Characteristics: Marine, Isolated from mixed culture
with *Tribonema marinum* J.Feldmann,
Formerly identified as *T. marinum*,
COXI gene (AF037996), *tufA* gene (AF038004),
18S rRNA gene (AF038005)
References: 19, 293
- Actinastrum hantzschii* Lagerheim
415
Lake Kasumigaura / Ibaraki (1983-07)
Axenic, Clonal, F.Kasai (1983-07)
Identified by: M.Watanabe
Culture conditions: C(S), 20° C, 4 µE/m² sec, 3M,
(25° C, 30 µE/m² sec)
Characteristics: Indicator, Freshwater,
COXI gene (D63660)
F7-4
References: 66, 399
- Alexandrium affine* (Inoue et Fukuyo) Balech
673
Harima-Nada / Seto Inland Sea (1980-09)
Axenic, Clonal, S.Yoshimatsu (1980-09)
Identified by: S.Yoshimatsu
Culture conditions: ESM, 20° C, 40 µE/m² sec, 1M
Characteristics: Red tide, Marine, Untransportable
- Alexandrium catenella* (Whedon et Kofoid) Balech
Syn. *Protogonyaulax catenella*
(Whedon et Kofoid) Taylor
220
Tsuda Bay / Kagawa (1980-06)
Axenic, Clonal, S.Yoshimatsu
- Culture conditions: ESM, 20° C, 40 µE/m² sec, 2M
Characteristics: Red tide, Marine, Unstable,
Untransportable
KGW-31-1
- 519
Owase Bay / Mie
Axenic, Clonal, T.Okaichi
Culture conditions: ESM, 20° C, 40 µE/m² sec, 2M
Characteristics: Red tide, Marine, Unstable,
Untransportable
KGW-41
- 520
Hachinohe Harbor / Aomori (1988-08)
Unialgal, Clonal, T.Sawaguchi (1988-08)
Identified by: T.Sawaguchi
Culture conditions: ESM, 20° C, 40 µE/m² sec, 1M
Characteristics: Red tide, Marine, Unstable,
Untransportable
88HH-2
- 674
Harima-Nada / Seto Inland Sea (1980-06)
Axenic, Clonal, S.Yoshimatsu (1980-06)
Identified by: S.Yoshimatsu
Culture conditions: ESM, 20° C, 40 µE/m² sec, 1M
Characteristics: Red tide, Marine, Mating type +,
Untransportable
Ac 1
- 675
Harima-Nada / Seto Inland Sea (1980-06)
Axenic, Clonal, S.Yoshimatsu (1980-06)
Identified by: S.Yoshimatsu
Culture conditions: ESM, 20° C, 40 µE/m² sec, 1M
Characteristics: Red tide, Marine, Mating type -,
Untransportable
Ac 5
- 677
Yamakawa Bay / Kagoshima (1988-03)
Axenic, Clonal, S.Yoshimatsu (1988-04)
Identified by: S.Yoshimatsu
Culture conditions: ESM, 20° C, 40 µE/m² sec, 1M
Characteristics: Red tide, Marine, Untransportable
Acy-6
- Alexandrium hiranoi* Kita et Fukuyo
612
Jogashima, Misaki / Kanagawa (1984-08)

- Unialgal, Clonal, T.Kita (1984-08)
 Identified by: T.Kita & Y.Fukuyo
 Culture conditions: ESM, 20° C, 40 µE/m²sec, 2M
 Characteristics: Toxic, Marine, Untransportable
 References: 139, 140, 186
- Alexandrium insuetum* Balech
 678
 Uchiumi Bay / Kagawa (1985-06)
 Axenic, Clonal, S.Yoshimatsu (1985-06)
 Identified by: S.Yoshimatsu
 Culture conditions: ESM, 20° C, 40 µE/m²sec, 1M
 Characteristics: Red tide, Marine, Untransportable
- Amphidinium britannicum* (Herdman) Lebour
 405
 Hasaki / Ibaraki (1987-05)
 Unialgal, Clonal, T.Sawaguchi (1987-05)
 Identified by: T.Sawaguchi
 Culture conditions: ESM, 20° C, 40 µE/m²sec, 1M
 Characteristics: Benthic, Marine, Untransportable
 HASS-1
- Amphidinium carterae* Hulburt
 331
 Iriomote Isl. / Okinawa (1986-01)
 Axenic, Clonal, T.Sawaguchi (1986-02)
 Identified by: T.Sawaguchi
 Culture conditions: ESM, 20° C, 32 µE/m²sec, 1M
 Characteristics: Marine, Unstable, Untransportable
 IIDA
- Amphidinium klebsii* Coll
 613
 Aburatsubo Bay / Kanagawa (1993-04)
 Unialgal, Clonal, M.Murata (1994-03)
 Identified by: Y.Fukuyo
 Culture conditions: f/2, ESM, 20° C, 40 µE/m²sec,
 1M
 Characteristics: Marine, Untransportable
 AK-1
- Anabaena affinis* Lemmermann
 40
 Lake Kasumigaura / Ibaraki (1974-08)
 IAM M-168, Unialgal, Clonal, M.M.Watanabe
 (1974-08)
 Identified by: M.M.Watanabe
 Culture conditions: CT, 25° C, 12 µE/m²sec, 1M
 Characteristics: Water bloom, Freshwater, Unstable
 References: 80, 169, 216, 399, 431
- Anabaena circinalis* Rabenhorst ex Bornet et Flahault
 41
 Lake Kasumigaura / Ibaraki (1974-08)
 IAM M-169, Axenic, Clonal, M.M.Watanabe
 (1974-08)
 Identified by: M.M.Watanabe
 Culture conditions: CB, 25° C, 24 µE/m²sec, 1M
 Characteristics: Water bloom, Freshwater, Unstable
 References: 80, 215, 216, 319, 399
- Anabaena compacta* (Nygaard) Hickel
 806
 Rostherne Mere, Cheshire / England
 CCAP 1403/24, Axenic, Clonal, Jaworski (1978)
 Reidentified by: R.Li
 Culture conditions: CT, 20° C, 8 µE/m²sec, 1M
 Characteristics: Freshwater, Water bloom, Formerly
 identified as *Anabaena spiroides* Klebahn
- 835
 Esthwaite Water, Cambria / England
 CCAP 1403/29, Unialgal, Clonal, Butterwick (1985)
 Reidentified by: R.Li
 Culture conditions: CT, 20° C, 8 µE/m²sec, 1M
 Characteristics: Freshwater, Water bloom, Formerly
 identified as *Anabaena spiroides* Klebahn
- Anabaena cylindrica* Lemmermann
 19
 IAM M-1, Axenic, Non-clonal
 Culture conditions: MDM(S), 20° C, 4 µE/m²sec,
 4M, (25° C, 30 µE/m²sec)
 Characteristics: Freshwater, Nitrogen fixation,
 Reidentified by M.M.Watanabe
 References: 3, 4, 12, 32, 36, 38, 39, 40, 41, 42, 43, 44,
 45, 49, 64, 80, 165, 193, 259, 260, 261, 262, 279,
 280, 281, 334, 350, 366, 373, 399, 426, 427, 428,
 429, 430, 431
- Anabaena ellipsoides* Bolochozow
 828
 Fishpond, Wuhan / China (1996-01)
 Unialgal, Clonal, R.Li (1996-01)
 Identified by: R.Li
 Culture conditions: CT, 20° C, 8 µE/m²sec, 1M
 Characteristics: Freshwater, Water bloom
 Ana HB
- Anabaena flos-aquae* Brébisson ex Bornet et Flahault
 f. *flos-aquae*
 73
 Lake Kasumigaura / Ibaraki (1978-08)

- TAC 32, Axenic, Clonal, M.Watanabe (1978-08)
 Identified by: M.Watanabe
 Culture conditions: MA, 25° C, 24 µE/m²sec, 1M,
 [Cryopreserved]
 Characteristics: Water bloom, Indicator, Freshwater,
 Unstable
 K-TAN-32
 References: 126, 216, 325, 399
- 74
 Lake Kasumigaura / Ibaraki (1978-08)
 TAC 33, Unialgal, Clonal, M.Watanabe (1978-08)
 Identified by: M.Watanabe
 Culture conditions: CT, 25° C, 24 µE/m²sec, 1M
 Characteristics: Water bloom, Indicator, Freshwater,
 Unstable
 K-TAN-33
 References: 189, 190, 191, 399
- 75
 Lake Kasumigaura / Ibaraki (1978-12)
 TAC 43, Unialgal, Clonal, M.Watanabe (1978-12)
 Identified by: M.Watanabe
 Culture conditions: CB, 25° C, 24 µE/m²sec, 1M,
 [Cryopreserved]
 Characteristics: Water bloom, Indicator, Freshwater,
 Unstable
 K-TAN-43
 Reference: 399
- Anabaena kisseleviana* Elenkin
 807
 Lake Kasumigaura / Ibaraki (1978-08)
 TAC 34, Unialgal, Clonal, M.Watanabe (1978-08)
 Identified by: R.Li
 Culture conditions: CT, 20° C, 8 µE/m²sec, 1M
 Characteristics: Freshwater, Water bloom
- Anabaena lemmermannii* Richter
 833
 Lake Steinsfjorden, Buskerud / Norway
 NIVA CYA 82, Unialgal, Clonal (1980)
 Reidentified by: R.Li
 Culture conditions: CT, 20° C, 8 µE/m²sec, 1M
 Characteristics: Freshwater, Water bloom, Unstable,
 Formerly identified as *Anabaena circinalis*
 Rabenhorst ex Bornet et Flahault
- Anabaena mendotae* Trelease
 808
 Lake Akan / Hokkaido (1990-08)
 TAC 437, Unialgal, Clonal, Y.Niiyama (1990-08)
- Identified by: R.Li
 Culture conditions: CT, 20° C, 8 µE/m²sec, 1M
 Characteristics: Freshwater, Water bloom
 A28
- Anabaena mucosa* Komárková et Eloranta
 809
 Lake Toro / Hokkaido (1990-08)
 TAC 426, Unialgal, Clonal, Y.Niiyama (1990-08)
 Identified by: R.Li
 Culture conditions: CT, 20° C, 8 µE/m²sec, 1M
 Characteristics: Freshwater, Water bloom, Unstable
 A10
- Anabaena oumiana* Watanabe
 829
 Chon Buri / Thailand (1998-05)
 Unialgal, Clonal, R.Li (1998-05)
 Identified by: R.Li
 Culture conditions: CT, 20° C, 8 µE/m²sec, 1M
 Characteristics: Freshwater, Water bloom
 Ana T1
- Anabaena planktonica* Brunthaler
 810
 Ohnuma / Hokkaido (1990-08)
 TAC 421, Axenic, Clonal, Y.Niiyama (1990-08)
 Identified by: M.Watanabe
 Culture conditions: CT, 20° C, 8 µE/m²sec, 1M
 Characteristics: Freshwater, Water bloom
 A3
 Reference: 169
- 811
 Ohnuma / Hokkaido (1990-08)
 TAC 422, Axenic, Clonal, Y.Niiyama (1990-08)
 Identified by: M.Watanabe
 Culture conditions: CT, 20° C, 8 µE/m²sec, 1M
 Characteristics: Freshwater, Water bloom
 A4
- 812
 Lake Toro / Hokkaido (1990-08)
 TAC 424, Unialgal, Clonal, Y.Niiyama (1990-08)
 Identified by: M.Watanabe
 Culture conditions: CT, 20° C, 8 µE/m²sec, 1M
 Characteristics: Freshwater, Water bloom
 A7
- 813
 Lake Tofutsu / Hokkaido (1990-08)
 TAC 434, Unialgal, Clonal, Y.Niiyama (1990-08)

- Identified by: R.Li
Culture conditions: CT, 20° C, 8 µE/m² sec, 1M
Characteristics: Freshwater, Water bloom
A25
- 814
Lake Tofutsu / Hokkaido (1990-08)
TAC 435, Axenic, Clonal, Y.Niiyama (1990-08)
Identified by: R.Li
Culture conditions: CT, 20° C, 8 µE/m² sec, 1M
Characteristics: Freshwater, Water bloom
A26
- 815
Esthwaite Water, Cambria / England
CCAP 1403/19, Axenic, Clonal, Jaworski (1968)
Reidentified by: R.Li
Culture conditions: CT, 20° C, 8 µE/m² sec, 1M
Characteristics: Freshwater, Water bloom, Formerly
identified as *Anabaena solitaria* Klebahn
- 816
Blelham Tarn, Cambria / England
CCAP 1403/27, Axenic, Clonal, Jaworski (1985)
Reidentified by: R.Li
Culture conditions: CT, 20° C, 8 µE/m² sec, 1M
Characteristics: Freshwater, Water bloom, Formerly
identified as *Anabaena solitaria* Klebahn
- 817
Inba-numa / Chiba (1995-05)
Unialgal, Clonal, R.Li (1995-05)
Identified by: R.Li
Culture conditions: CT, 20° C, 8 µE/m² sec, 1M
Characteristics: Freshwater, Water bloom, Offensive
taste and odor
Inba 2
- 834
Lake Langsævatn, Aust-Agder / Norway (1979)
NIVA CYA 66, Unialgal, Clonal
Reidentified by: R Li
Culture conditions: CT, 20° C, 8 µE/m² sec, 1M
Characteristics: Freshwater, Water bloom, Formerly
identified as *Anabaena solitaria* Klebahn f.
planktonica (Brunnthaler) Komárek
- Anabaena smithii* (Komárek) Watanabe
818
Lake Barato / Hokkaido (1989-08)
TAC 116, Unialgal, Clonal, M.Watanabe (1989-08)
Identified by: R.Li
Culture conditions: CT, 20° C, 8 µE/m² sec, 1M
Characteristics: Freshwater, Water bloom
- 819
Lake Barato / Hokkaido (1990-08)
TAC 428, Unialgal, Clonal, Y.Niiyama (1990-08)
Identified by: M.Watanabe
Culture conditions: CT, 20° C, 8 µE/m² sec, 1M
Characteristics: Freshwater, Water bloom
A14
- 820
Hirosaki / Aomori (1990-06)
TAC 431, Unialgal, Clonal, Y.Niiyama (1990-06)
Identified by: M.Watanabe
Culture conditions: CT, 20° C, 8 µE/m² sec, 1M
Characteristics: Freshwater, Water bloom
A18
- 821
Hirosaki / Aomori (1990-06)
TAC 432, Unialgal, Clonal, Y.Niiyama (1990-06)
Identified by: M.Watanabe
Culture conditions: CT, 20° C, 8 µE/m² sec, 1M
Characteristics: Freshwater, Water bloom
A19
Reference: 169
- 822
Lake Akan / Hokkaido (1991-07)
TAC 450, Unialgal, Clonal, Y.Niiyama (1991-07)
Identified by: R.Li
Culture conditions: CT, 20° C, 8 µE/m² sec, 1M
Characteristics: Freshwater, Water bloom
A51
- 823
Lake Okutama / Tokyo (1991-07)
TAC 452, Unialgal, Clonal, M.Watanabe (1991-07)
Identified by: R.Li
Culture conditions: CT, 20° C, 8 µE/m² sec, 1M
Characteristics: Freshwater, Water bloom, Offensive
taste and odor, Unstable
210
- 824
Hase River, Ogasawara Isl. / Tokyo (1998-03)
Unialgal, Clonal, R.Li (1998-03)
Identified by: R.Li
Culture conditions: CT, 20° C, 8 µE/m² sec, 1M
Characteristics: Freshwater, Water bloom, Offensive

- taste and odor
Ana Ha 1
- 830
Lam Takong / Thailand (1997-07)
Unialgal, Clonal, R.Li (1997-07)
Identified by: R.Li
Culture conditions: CT, 20° C, 8 µE/m² sec, 1M
Characteristics: Freshwater, Water bloom
Ana130
- 831
Chon Buri / Thailand (1998-05)
Unialgal, Clonal, R.Li (1998-05)
Identified by: R.Li
Culture conditions: CT, 20° C, 8 µE/m² sec, 1M
Characteristics: Freshwater, Water bloom
Ana T3
- Anabaena solitaria* Klebahn f. *solitaria*
80
Lake Kasumigaura / Ibaraki (1978-12)
TAC 42, Axenic, Clonal, M.Watanabe (1978-12)
Identified by: M.Watanabe
Culture conditions: CB, 25° C, 24 µE/m² sec, 20D
Characteristics: Water bloom, Freshwater, Unstable
K-TAN-42
References: 169, 216, 399
- Anabaena spiroides* Klebahn
76
Lake Kasumigaura / Ibaraki (1983-06)
Unialgal, Clonal, S.Suda (1983-06)
Identified by: S.Suda
Culture conditions: CA, 25° C, 24 µE/m² sec, 1M
Characteristics: Water bloom, Indicator, Freshwater,
Unstable
K-A-12
References: 169, 216, 263, 399
- Anabaena spiroides* Klebahn
f. *crassa* (Lemmermann) Elenkin
78
Lake Kasumigaura / Ibaraki (1978-07)
TAC 30, Axenic, Clonal, M.Watanabe (1978-07)
Identified by: M.Watanabe
Culture conditions: CT, 25° C, 24 µE/m² sec, 1M,
[Cryopreserved]
Characteristics: Water bloom, Indicator, Freshwater,
Unstable
K-TAN-30
References: 169, 215, 216
- Anabaena spiroides* Klebahn f. *spiroides*
77
Lake Kasumigaura / Ibaraki (1978-08)
TAC 31, Unialgal, Clonal, M.Watanabe (1978-08)
Identified by: M.Watanabe
Culture conditions: CT, 25° C, 24 µE/m² sec, 1M,
[Cryopreserved]
Characteristics: Water bloom, Indicator, Freshwater,
Unstable
K-TAN-31
References: 126, 399
- 79
Lake Kasumigaura / Ibaraki (1978-07)
TAC 28, Axenic, Clonal, M.Watanabe (1978-07)
Identified by: M.Watanabe
Culture conditions: CB, 25° C, 24 µE/m² sec, 1M,
[Cryopreserved]
Characteristics: Water bloom, Indicator, Freshwater,
Unstable
K-TAN-28
Reference: 169
- 263
Lake Kasumigaura / Ibaraki (1978-07)
TAC 27, Axenic, Clonal, M.Watanabe (1978-07)
Identified by: M.Watanabe
Culture conditions: CT, 25° C, 24 µE/m² sec, 1M
Characteristics: Water bloom, Freshwater, Unstable
K-TAN-27
Reference: 399
- Anabaena ucrainica* (Schkorb) Watanabe
825
Lake Sagami / Kanagawa (1991-08)
TAC 448, Unialgal, Clonal, Y.Niiyama (1991-08)
Identified by: M.Watanabe
Culture conditions: CT, 20° C, 8 µE/m² sec, 1M
Characteristics: Freshwater, Water bloom
A48
Reference: 169
- 826
Lake Sagami / Kanagawa (1991-08)
TAC 449, Unialgal, Clonal, Y.Niiyama (1991-08)
Identified by: M.Watanabe
Culture conditions: CT, 20° C, 8 µE/m² sec, 1M
Characteristics: Freshwater, Water bloom
A50
Reference: 169

- 832
Hochimin / Vietnam (1998-12)
Unialgal, Clonal, R.Li (1998-12)
Identified by: R.Li
Culture conditions: CT, 20° C, 8 µE/m²sec, 1M
Characteristics: Freshwater, Water bloom, Unstable
Ana V2
- Anabaena variabilis* Kützing ex Bornet et Flahault
23
IAM M-2, Axenic, Clonal
Culture conditions: MDM(S), 20° C, 4 µE/m²sec,
4M, (25° C, 30 µE/m²sec)
Characteristics: Freshwater, Non-heterocystous
variant
References: 4, 27, 28, 29, 42, 43, 44, 80, 334, 351,
366
- Anabaena viguieri* Denis et Frémy
827
Shikata-futago-ike / Hyogo (1990-09)
TAC 433, Unialgal, Clonal, Y.Niiyama (1990-09)
Identified by: R.Li
Culture conditions: CT, 20° C, 8 µE/m²sec, 1M
Characteristics: Freshwater, Water bloom
A23
- Anabaenopsis circularis*
(G.S.West) Woloszynska et Miller
21
IAM M-4, Axenic, Clonal, A.Watanabe
Identified by: Hirano
Culture conditions: MDM(S), 20° C, 4 µE/m²sec,
4M, (25° C, 30 µE/m²sec), [Cryopreserved]
Characteristics: Freshwater,
Reidentified by M.M.Watanabe
References: 4, 80, 193, 366, 372, 399
- Aphanizomenon flos-aquae* (Lemmermann) Ralfs
f. *gracile* (Lemmermann) Elenkin
81
Lake Kasumigaura / Ibaraki (1978-01)
TAC 1, Axenic, Clonal, M.Watanabe (1978-02)
Identified by: M.Watanabe
Culture conditions: CB, 25° C, 24 µE/m²sec, 1M,
[Cryopreserved]
Characteristics: Water bloom, Indicator, Freshwater,
Unstable
K-TAN-1
References: 169, 216, 325, 399
- Aphanocapsa montana* Cramer
416
Nikko / Tochigi (1987-04)
Unialgal, Non-clonal, F.Kasai (1987-04)
Identified by: M.M.Watanabe
Culture conditions: CSi, CSi+Cu, 20° C, 4 µE/m²sec,
4M, (20° C, 12 µE/m²sec), [Cryopreserved]
Characteristics: Freshwater
NK-24
Reference: 338
- **Arthrospira platensis* Gomont
See *Spirulina platensis* (Gomont) Geitler
- Asterionella glacialis* Castracane
265
Matoya Bay / Mie (1984-09)
Unialgal, Clonal, T.Sawaguchi (1984-09)
Identified by: T.Sawaguchi
Culture conditions: f/2, 10° C, 25 µE/m²sec, 1M
Characteristics: Marine
MB-B-1
- 417
Maizuru Bay / Kyoto (1985-10)
Unialgal, Clonal, C.E.Riquelme (1985-10)
Identified by: C.E.Riquelme
Culture conditions: f/2, 15° C, 20 µE/m²sec, 1M
Characteristics: Marine
- Astrephomene gubernaculifera* Pocock
418
Kaisei / Kanagawa (1981-4)
Axenic, Clonal, H.Nozaki (1981-05)
Identified by: H.Nozaki
Culture conditions: VTAC, 20° C, 12 µE/m²sec, 1M
Characteristics: Freshwater, Heterothallic,
Mating type -, Crosses with NIES-419,
rbcL gene (D63428)
1520-4 (-)
References: 222, 243
- 419
Kaisei / Kanagawa (1981-4)
Axenic, Clonal, H.Nozaki (1981-05)
Identified by: H.Nozaki
Culture conditions: VTAC, 20° C, 12 µE/m²sec, 1M
Characteristics: Freshwater, Heterothallic,
Mating type +, Crosses with NIES-418
1520-1 (+)
Reference: 222

628

Hayama / Kanagawa (1980-12)

Unialgal, Clonal, H.Nozaki (1981-07)

Identified by: H.Nozaki

Culture conditions: VTAC, 20° C, 22 µE/m² sec, 1M

Characteristics: Freshwater, Heterothallic, Isogamy,

Mating type -

1727-1(-)

Characteristics: Freshwater

AT1-7

Reference: 338

Astrephomene perforata Nozaki

564

Hayama / Kanagawa (1980-12)

UTEX 2474, Unialgal, Clonal, H.Nozaki (1981-06)

Identified by: H.Nozaki

Culture conditions: VTAC, 20° C, 22 µE/m² sec, 1M

Characteristics: Freshwater, Type strain,

Heterothallic, Isogamy, Mating type +,

Crosses with NIES-565, *rbcL* gene (D63429)

1620-3-2

References: 222, 243

Basichlamys sacculifera (Scherffel) Skuja

Syn. *Gonium sacculiferum* Scherffel

566

Fujisawa / Kanagawa (1983-08)

Unialgal, Clonal, H.Nozaki (1983-09)

Identified by: H.Nozaki

Culture conditions: AF-6, 20° C, 22 µE/m² sec, 1M

Characteristics: Freshwater, Akinete forming,

rbcL gene (D63430)

3907-1

References: 226, 243, 246

565

Hayama / Kanagawa (1980-12)

UTEX 2475, Unialgal, Clonal, H.Nozaki (1981-06)

Identified by: H.Nozaki

Culture conditions: VTAC, 20° C, 22 µE/m² sec, 1M

Characteristics: Freshwater, Type strain,

Heterothallic, Isogamy, Mating type -,

Crosses with NIES-564

1620-4-1

References: 222, 242

Botrydiopsis arrhiza Borzi

621

Shelford / England

CCAP 222/1B, Unialgal, George (1947)

Culture conditions: AF-6, 20° C, 32 µE/m² sec, 2M

Characteristics: Freshwater

Botrydium granulatum (L.) Greville

622

CCAP 805/3A, Axenic, Vischer (1939)

Culture conditions: AF-6, 20° C, 32 µE/m² sec, 2M

Characteristics: Freshwater

Aulosira laxa Kirchner ex Bornet et Flahault

50

Pusa / India

IAM M-128, Axenic, Non-clonal,

G.S.Venkataraman

Culture conditions: MDM(S), 20° C, 4 µE/m² sec,

4M, (25° C, 30 µE/m² sec), [Cryopreserved]

Characteristics: Freshwater, Nitrogen fixation,

M-128 as *Aulosira fertissima* in IAM,

Reidentified by M.M.Watanabe

References: 80, 399

Botryococcus braunii Kützing

836

Imuta-ike Pond / Kagoshima (1997-06)

Unialgal, Clonal, F.Mori (1997-06)

Identified by: F.Mori

Culture conditions: CA, 20° C, 24 µE/m² sec, 4M

Characteristics: Freshwater

Brachiomonas submarina Bohlin

375

Hachinohe Harbor / Aomori (1986-08)

Axenic, Clonal, T.Sawaguchi (1986-08)

Identified by: T.Sawaguchi

Culture conditions: ESM, 15° C, 20 µE/m² sec, 1M

Characteristics: Marine, Brackish

86-SuHH-2

Auxenochlorella protothecoides (Kruger) Kalina

629

Watarase River / Gunma (1987-08)

Unialgal, Clonal, F.Kasai (1987-08)

Identified by: F.Kasai

Culture conditions: C, 15° C, 6 µE/m² sec, 6M,

(15° C, 15 µE/m² sec)

Cachonina niei Loeblich III

420

Iriomote Isl. / Okinawa (1986-01)

Axenic, Clonal, T.Sawaguchi (1986-02)

Identified by: T.Sawaguchi

Culture conditions: ESM, 20° C, 40 µE/m² sec, 1M

- Characteristics: Marine, Untransportable
IID-1
- 614
Kashiwazaki / Niigata (1986-08)
Unialgal, Clonal, T.Sawaguchi (1986-08)
Identified by: T.Sawaguchi
Culture conditions: ESM, 20° C, 40 µE/m² sec, 1M
Characteristics: Marine, Untransportable
KSTH-29
- Calothrix brevissima* G.S.West
22
Palau Isl. (1941-09)
IAM M-7, Axenic, Non-clonal, A.Watanabe
Identified by: K.Negoro.
Culture conditions: MDM(S), 20° C, 4 µE/m² sec,
4M, (25° C, 30 µE/m² sec), [Cryopreserved]
Characteristics: Freshwater, Nitrogen fixation
Chromatic adaptation
References: 80, 278, 366
- Calothrix crustacea* Thuret ex Bornet et Flahault
266
Oshoro Bay / Hokkaido (1972-09)
IAM M-171, Unialgal, Clonal, M.M.Watanabe
(1972-09)
Identified by: M.M.Watanabe
Culture conditions: f/2, 20° C, 4 µE/m² sec, 6M,
(20° C, 12 µE/m² sec), [Cryopreserved]
Characteristics: Marine
References: 80, 387, 402
- Calothrix parasitica* Thuret ex Bornet et Flahault
267
Oshoro Bay / Hokkaido (1972-07)
IAM M-172, Axenic, Clonal, M.M.Watanabe
(1972-07)
Identified by: M.M.Watanabe
Culture conditions: f/2, 20° C, 4 µE/m² sec, 6M,
(20° C, 12 µE/m² sec), [Cryopreserved]
Characteristics: Indicator, Marine, Endophyte
in *Nemalion* (Rhodophyceae)
Reference: 80
- 334
Oshoro Bay / Hokkaido (1973-02)
IAM M-173, Unialgal, Clonal, M.M.Watanabe
(1973-02)
Identified by: M.M.Watanabe
Culture conditions: f/2, 20° C, 4 µE/m² sec, 6M,
(20° C, 12 µE/m² sec), [Cryopreserved]
Characteristics: Indicator, Marine, Endophyte
in *Codium* (Ulvophyceae)
Reference: 80
- Calothrix scopulorum* Agardh ex Bornet et Flahault
268
Oshoro Bay / Hokkaido (1972-09)
IAM M-174, Unialgal, Clonal, M.M.Watanabe
(1972-09)
Identified by: M.M.Watanabe
Culture conditions: MKM(S), 20° C, 4 µE/m² sec,
6M, (20° C, 12 µE/m² sec), [Cryopreserved]
Characteristics: Indicator, Marine
References: 80, 387, 402
- Carteria cerasiformis* Nozaki et al.
424
Lake Kasumigaura / Ibaraki (1983-08)
Axenic, Clonal, S.Suda (1983-08)
Reidentified by: H.Nozaki et al.
Culture conditions: AF-6, 20° C, 22 µE/m² sec, 2M
Characteristics: Freshwater, Formerly identified as
Carteria inversa (Korshikov) Bourrelly,
rbcL gene (D89767)
Kas-10
References: 239, 247
- 425
Tsukuba / Ibaraki (1985-11)
Axenic, Clonal, S.Suda (1985-11)
Reidentified by: H.Nozaki et al.
Culture conditions: AF-6, 20° C, 22 µE/m² sec, 2M
Characteristics: Freshwater, Formerly identified as
Carteria inversa (Korshikov) Bourrelly, Type
strain of *Carteria cerasiformis* Nozaki et al.,
rbcL gene (D89768)
w-8-15
References: 239, 247
- Carteria crucifera* Korshikov ex Pascher
421
Tsuchiura / Ibaraki (1986-02)
Axenic, Clonal, S.Suda (1986-05)
Identified by: S.Suda
Culture conditions: CYT, 20° C, 22 µE/m² sec, 2M
Characteristics: Freshwater, *rbcL* gene (D63431)
SIST3-1
References: 243, 239, 247
- 630
New Haven / USA

- UTEX 432, Unialgal, Clonal, R.A.Lewin
 Reidentified by: H.Nozaki
 Culture conditions: AF-6, 20° C, 22 μE/m² sec, 1M
 Characteristics: Freshwater, *rbcL* gene (D89758)
 References: 239, 247
- Carteria eugametos* Mitra
 631
 Saiwai-ku / Kawasaki (1990-10)
 Unialgal, Clonal, H.Nozaki (1991-04)
 Identified by: H.Nozaki
 Culture conditions: AF-6, 20° C, 22 μE/m² sec, 1M
 Characteristics: Freshwater, Homothallic, Isogamy,
rbcL gene (D89762)
 91-409-1
 References: 237, 247
- 632
 Saiwai-ku / Kawasaki (1990-10)
 Unialgal, Clonal, H.Nozaki (1991-04)
 Identified by: H.Nozaki
 Culture conditions: AF-6, 20° C, 22 μE/m² sec, 1M
 Characteristics: Freshwater, Homothallic, Isogamy,
rbcL gene (D89763)
 91-421-4
 References: 237, 239, 247
- 633
 Shirako / Chiba (1991-03)
 Unialgal, Clonal, H.Nozaki (1991-05)
 Identified by: H.Nozaki
 Culture conditions: AF-6, 20° C, 22 μE/m² sec, 1M
 Characteristics: Freshwater, Homothallic, Isogamy,
rbcL gene (D89764)
 91-504-1
 References: 237, 239, 247
- 634
 UTEX 2161, Unialgal, Clonal, B.Vandover (1972)
 Reidentified by: H.Nozaki
 Culture conditions: AF-6, 20° C, 22 μE/m² sec, 1M
 Characteristics: Freshwater, Homothallic, Isogamy,
rbcL gene (D89761)
 References: 239, 247
- 635
 Allahabad / India
 UTEX 233, Unialgal, Clonal, Pringsheim.O.
 Culture conditions: AF-6, 20° C, 22 μE/m² sec, 1M
 Characteristics: Freshwater, Type strain of *Carteria*
eugametos Mitra, *rbcL* gene (D89759)
 References: 239, 247
- 636
 California / USA
 UTEX 1032, Unialgal, Clonal, A.Waters
 Reidentified by: H.Nozaki
 Culture conditions: AF-6, 20° C, 22 μE/m² sec, 1M
 Characteristics: Freshwater, Formerly identified as
Carteria olivieri G. S. West (Starr and Zeikus
 1993), *rbcL* gene (D89760)
 References: 239, 247
- Carteria inversa* (Korshikov) Bourrelly
 422
 Tsukuba / Ibaraki (1982-11)
 Axenic, Clonal, F.Kasai (1982-11)
 Identified by: S.Suda
 Culture conditions: C, 20° C, 22 μE/m² sec, 3M
 Characteristics: Freshwater, *rbcL* gene (D89765)
 134-4
 References: 239, 247
- 423
 Higashihiroshima / Hiroshima (1983-08)
 Axenic, Clonal, M.Erata (1983-08)
 Identified by: S.Suda
 Culture conditions: C, 20° C, 22 μE/m² sec, 3M
 Characteristics: Freshwater, *rbcL* gene (D89766)
 106
 References: 239, 247
- * *Carteria inversa* (Korshikov) Bourrelly
 424
 See *Carteria cerasiformis* Nozaki et al.
- * *Carteria inversa* (Korshikov) Bourrelly
 425
 See *Carteria cerasiformis* Nozaki et al.
- Carteria klebsii* (Dangeard) Francé
 426
 Tsuchiura / Ibaraki (1986-02)
 Axenic, Clonal, S.Suda (1986-05)
 Identified by: S.Suda
 Culture conditions: AF-6, 20° C, 22 μE/m² sec, 2M
 Characteristics: Freshwater
 SIST7-4
- Carteria multifilis* (Fresenius) Dill
 427
 Kashiwa / Chiba (1986)
 Axenic, Clonal, M.M.Watanabe (1986)
 Identified by: S.Suda

- Culture conditions: VT, 20° C, 22 μE/m² sec, 2M
 Characteristics: Freshwater
 Ca1-2
- Carteria obtusa* Dill
 428
 Kashiwa / Chiba (1986-09)
 Axenic, Clonal, M.M.Watanabe (1986-09)
 Identified by: S.Suda
 Culture conditions: C, 20° C, 22 μE/m² sec, 2M
 Characteristics: Freshwater, *rbcL* gene (D89769)
 Ca-2-1
 Reference: 247
- 429
 Tsuchiura / Ibaraki (1986-02)
 Axenic, Clonal, M.Kasama (1986-03)
 Identified by: S.Suda
 Culture conditions: AF-6, 20° C, 22 μE/m² sec, 2M
 Characteristics: Freshwater
 SIS5-20
- 430
 Kashiwa / Chiba (1986-09)
 Axenic, Clonal, M.M.Watanabe (1986-09)
 Identified by: S.Suda
 Culture conditions: C, 20° C, 22 μE/m² sec, 2M
 Characteristics: Freshwater
 Ca2-3
- 431
 Tsuchiura / Ibaraki (1986-02)
 Axenic, Clonal, S.Suda (1986-05)
 Identified by: S.Suda
 Culture conditions: AF-6, 20° C, 22 μE/m² sec, 2M
 Characteristics: Freshwater
 SIST6-3
- Carteria radiosa* Korshikov ex Pascher
 432
 Tsukuba / Ibaraki (1985-11)
 Axenic, Clonal, S.Suda (1985-11)
 Identified by: S.Suda
 Culture conditions: AF-6, 20° C, 22 μE/m² sec, 2M
 Characteristics: Freshwater, *rbcL* gene (D89770)
 w-5-2
 Reference: 247
- Ceratium hirundinella* (O.F.Müller) Schrank
 376
 Lake Hinuma / Ibaraki (1986-06)
 Unialgal, Clonal, M.M.Watanabe (1986-06)
- Identified by: M.M.Watanabe
 Culture conditions: URO, 20° C, 32 μE/m² sec, 1M
 Characteristics: Brackish, Freshwater, Unstable,
 Untransportable
 860627-10
- Chaetoceros didymus* Ehrenberg
 586
 Hitachi / Ibaraki (1990-09)
 Unialgal, Non-clonal, S.Ono (1990-10)
 Identified by: S.Ono
 Culture conditions: f/2, 15° C, 10 μE/m² sec, 1M
 Characteristics: Red tide, Marine
 St-4
- Chaetoceros sociale* Lauder
 377
 Shitaru Harbor / Shizuoka (1985-05)
 Unialgal, Clonal, T.Sawaguchi (1985-05)
 Identified by: T.Sawaguchi
 Culture conditions: f/2, 5° C, 15 μE/m² sec, 20D
 Characteristics: Marine
 STHB-4
- 553
 Tokyo Bay / Tokyo (1991-10)
 Unialgal, Clonal, S.Ono (1991-10)
 Identified by: S.Ono
 Culture conditions: f/2, 5° C, 15 μE/m² sec, 1M
 Characteristics: Red tide, Marine
 T-1
- Chamaesiphon polymorphus* Geitler
 433
 Lake Mashu / Hokkaido (1987-09)
 Unialgal, Non-clonal, F.Kasai (1987-09)
 Identified by: M.M.Watanabe
 Culture conditions: CSi, 10° C, 6 μE/m² sec, 2M,
 (10° C, 15 μE/m² sec), [Cryopreserved]
 Characteristics: Freshwater
 M-29
 References: 338, 339
- Chamaesiphon subglobosus* Lemmermann
 434
 Miyata River / Ibaraki (1987-03)
 Unialgal, Non-clonal, F.Kasai (1987-05)
 Identified by: N.Takamura
 Culture conditions: CSi, CSi+Cu, 20° C, 4 μE/m² sec,
 3M, (20° C, 12 μE/m² sec), [Cryopreserved]
 Characteristics: Freshwater

- 2st-2-1
References: 337, 338, 339
- Characiochloris acuminata* Lee et Bold
637
El Tahin. Prov. Omo-Saber / Egypt
UTEX 2095, Unialgal, Clonal, F.Hindak (1963)
Identified by: K.W.Lee & H.C.Bold
Culture conditions: AF-6, 20° C, 22 μ E/m²sec, 2M
Characteristics: Freshwater, Type strain
Reference: 236
- Characiochloris sasae* Nozaki
567
Saiwai-ku, Kawasaki / Kanagawa (1990-10)
Unialgal, Clonal, H.Nozaki (1991-01)
Identified by: H.Nozaki
Culture conditions: AF-6, 20° C, 22 μ E/m²sec, 2M
Characteristics: Freshwater, Type strain,
Aplanospore forming
91-0106-1
Reference: 236
- 638
Saiwai-ku, Kawasaki / Kanagawa (1990-10)
Unialgal, Clonal, H.Nozaki (1991-01)
Identified by: H.Nozaki
Culture conditions: AF-6, 20° C, 22 μ E/m²sec, 1M
Characteristics: Freshwater, Aplanospore forming,
Endemic in Japan
91-0106-6
Reference: 236
- Characium angustum* A.Braun
639
Kinu River / Tochigi (1987-08)
Unialgal, F.Kasai, (1987-09-17)
Identified by: F.Kasai
Culture conditions: C, 15° C, 6 μ E/m²sec, 4M,
(15° C, 15 μ E/m²sec)
Characteristics: Freshwater
AK-5-2
Reference: 338
- Characium maximum* S.Watanabe
154
Sasebo / Nagasaki (1975-08)
Unialgal, Non-clonal, S.Watanabe
Identified by: S.Watanabe
Culture conditions: C(S), 20° C, 4 μ E/m²sec, 3M,
(25° C, 30 μ E/m²sec)
- Characteristics: Soil, Habitat: Garden Shrine
where *Cryptomeria japonica* was planted
6-EBO-2
Reference: 416
- Characium polymorphum* Printz
436
Between Ghorepani and Billethadi / Nepal
(1965-12)
IAM C-340, Unialgal, Clonal, T.Ichimura (1969-07)
Identified by: T.Ichimura
Culture conditions: C(S), 20° C, 4 μ E/m²sec, 3M,
(25° C, 30 μ E/m²sec)
Characteristics: Indicator, Freshwater
N-76-0
Reference: 80
- Chattonella antiqua* (Hada) Ono
1
Harima-Nada / Seto Inland Sea (1978-09)
Axenic, Clonal, M.M.Watanabe (1978-09)
Identified by: M.M.Watanabe
Culture conditions: f/2, ESM, 20° C, 32 μ E/m²sec,
1M
Characteristics: Red tide, Marine, Untransportable,
COXI gene (AF037990)
Ho-1
References: 20, 68, 90, 91, 92, 93, 94, 95, 156, 158,
194, 200, 201, 202, 203, 204, 205, 206, 207, 208,
209, 210, 217, 405, 432, 435
- 2
Osaka Bay / Osaka (1982-09)
Axenic, Clonal, S.Yamochi
Identified by: S.Yamochi
Culture conditions: f/2, ESM, 20° C, 32 μ E/m²sec,
1M
Characteristics: Red tide, Marine, Untransportable
OCH-a
Reference: 68
- 83
Off Hiketa / Seto Inland Sea (1977-08)
Axenic, Clonal, C.Ono
Culture conditions: f/2, 20° C, 32 μ E/m²sec, 1M
Characteristics: Red tide, Marine, Untransportable
KGW-2
References: 68, 357
- 84
Off Hiketa / Seto Inland Sea (1972)
Axenic, Clonal, T.Okaichi

- Culture conditions: $f/2$, 20°C , $32\ \mu\text{E}/\text{m}^2\text{sec}$, 1M
 Characteristics: Red tide, Marine, Untransportable
 KGW-6-1
 Reference: 68
- 85
 Shodo Isl. / Kagawa (1978-07)
 Axenic, Clonal, S.Yoshimatsu
 Culture conditions: $f/2$, ESM, 20°C , $32\ \mu\text{E}/\text{m}^2\text{sec}$, 1M
 Characteristics: Red tide, Marine, Untransportable
 KGW-8-5
 References: 68, 69
- 86
 Uranouchi Bay / Kochi (1980-11)
 Axenic, Clonal, S.Yoshimatsu
 Culture conditions: $f/2$, ESM, 20°C , $32\ \mu\text{E}/\text{m}^2\text{sec}$, 1M
 Characteristics: Red tide, Marine, Untransportable
 KGW-42-4
 References: 68, 69, 357
- 113
 Naoshima Isl. / Kagawa (1982-07)
 Axenic, Clonal, S.Yoshimatsu
 Culture conditions: $f/2$, 20°C , $32\ \mu\text{E}/\text{m}^2\text{sec}$, 1M
 Characteristics: Red tide, Marine, Untransportable
 KGW-59-2
 References: 6, 68
- 114
 Harima-Nada / Seto Inland Sea (1983-08)
 Axenic, Clonal, S.Yoshimatsu
 Culture conditions: $f/2$, 20°C , $32\ \mu\text{E}/\text{m}^2\text{sec}$, 1M
 Characteristics: Red tide, Marine, Untransportable
 KGW-74-8
 References: 68, 425
- 161
 Hiroshima Bay / Hiroshima
 Axenic, Clonal
 Culture conditions: $f/2$, 20°C , $32\ \mu\text{E}/\text{m}^2\text{sec}$, 1M
 Characteristics: Red tide, Marine, Untransportable
 Hiroshima-70
 References: 54, 55
- 557
 Hiroshima Bay / Hiroshima (1970-09)
 Axenic, Clonal, H.Takayama (1970-09)
 Culture conditions: $f/2$, 20°C , $32\ \mu\text{E}/\text{m}^2\text{sec}$, 1M
 Characteristics: Red tide, Marine, Untransportable
- 558
 Mikawa Bay / Aichi
 Axenic, Clonal, S.Toriumi
 Identified by: S.Toriumi
 Culture conditions: $f/2$, 20°C , $32\ \mu\text{E}/\text{m}^2\text{sec}$, 1M
 Characteristics: Red tide, Marine, Untransportable
- Chattonella marina* (Subrahmanyam) Hara et Chihara
 3
 Osaka Bay / Osaka (1982-08)
 Axenic, Clonal, S.Yamochi (1982-08)
 Culture conditions: $f/2$, 20°C , $32\ \mu\text{E}/\text{m}^2\text{sec}$, 1M
 Characteristics: Red tide, Marine, Untransportable
 OCH-m
 Reference: 357
- 14
 Harima-Nada / Seto Inland Sea (1983-02)
 Axenic, Clonal, M.M.Watanabe
 Identified by: M.M.Watanabe
 Culture conditions: $f/2$, 20°C , $32\ \mu\text{E}/\text{m}^2\text{sec}$, 1M
 Characteristics: Red tide, Marine, Untransportable
 H-53-11
 References: 68, 425
- 115
 Kinko Bay / Kagoshima (1978-06)
 Axenic, Clonal, Aramaki/Yoshimatsu
 Culture conditions: $f/2$, ESM, 20°C , $32\ \mu\text{E}/\text{m}^2\text{sec}$, 1M
 Characteristics: Red tide, Marine, Untransportable
 KGW-9-1
 Reference: 68
- 116
 Harima-Nada / Seto Inland Sea (1981-07)
 Axenic, Clonal, S.Yoshimatsu
 Culture conditions: $f/2$, 20°C , $32\ \mu\text{E}/\text{m}^2\text{sec}$, 1M
 Characteristics: Red tide, Marine, Untransportable
 KGW-46-7
 Reference: 68
- 117
 Naoshima Isl. / Kagawa (1982-07)
 Axenic, Clonal, S.Yoshimatsu
 Culture conditions: $f/2$, 20°C , $32\ \mu\text{E}/\text{m}^2\text{sec}$, 1M
 Characteristics: Red tide, Marine, Untransportable
 KGW-58-3
 Reference: 68

- 118
Harima-Nada / Seto Inland Sea (1983-07)
Axenic, Clonal, S. Yoshimatsu
Culture conditions: f/2, 20° C, 32 μE/m² sec, 1M
Characteristics: Red tide, Marine, Untransportable
KGW-75-2
References: 55, 68, 69, 288, 357
- 121
Kagoshima Bay / Kagoshima (1982)
Axenic, Clonal, T. Aramaki (1982)
Culture conditions: f/2, 20° C, 32 μE/m² sec, 1M
Characteristics: Red tide, Marine, Untransportable
KGO-57-1
References: 68, 69, 357
- 559
Maizuru Bay / Kyoto (1975-10)
Axenic, Clonal, H. Takayama (1975-10)
Identified by: S. Yoshimatsu
Culture conditions: f/2, 20° C, 32 μE/m² sec, 1M
Characteristics: Red tide, Marine, Untransportable
- Chattonella ovata* Hara et Chihara
603
Harima-Nada / Seto Inland Sea (1984-04)
Axenic, Clonal, I. Imai
Identified by: H. Nozaki
Culture conditions: f/2, 20° C, 32 μE/m² sec, 1M
Characteristics: Marine, Untransportable
References: 53, 55
- 671
Harima-Nada / Seto Inland Sea (1982-07)
Axenic, Clonal, S. Yoshimatsu (1982-07)
Identified by: S. Yoshimatsu
Culture conditions: ESM, 20° C, 40 μE/m² sec, 1M
Characteristics: Red tide, Marine, Untransportable
- Chattonella verruculosa* Hara et Chihara
670
Harima-Nada / Seto Inland Sea (1987-07)
Unialgal, Clonal, S. Yoshimatsu (1987-07)
Identified by: S. Yoshimatsu
Culture conditions: ESM, 20° C, 40 μE/m² sec, 1M
Characteristics: Red tide, Marine, Untransportable
Reference: 55
- Chilomonas paramecium* Ehrenberg
715
Sugadaira / Nagano (1985-11)
Axenic, Clonal, M. Erata (1985-11)
- Identified by: M. Erata
Culture conditions: CYT, 15° C, 15 μE/m² sec, 1M
Characteristics: Freshwater
#00210
Reference: 22
- 766
Lake Jusan-ko / Aomori (1987-07)
Unialgal, Clonal, M. Erata (1987-07)
Identified by: M. Erata
Culture conditions: CYT, 15° C, 15 μE/m² sec, 1M
Characteristics: Freshwater
#00318
- 767
Lake Jusan-ko / Aomori (1987-07)
Unialgal, Clonal, M. Erata (1987-07)
Identified by: M. Erata
Culture conditions: CYT, 15° C, 15 μE/m² sec, 1M
Characteristics: Freshwater
#00319
- Chlamydomonas augustae* Skuja
var. *ellipsoidea* S. Watanabe
158
Sumatra / Indonesia (1979-08)
Axenic, Clonal, S. Watanabe
Identified by: S. Watanabe
Culture conditions: C(S), 20° C, 4 μE/m² sec, 3M,
(25° C, 30 μE/m² sec)
Characteristics: Soil
ASE-242
References: 416, 417
- Chlamydomonas fasciata* Ettl
437
Tsukuba / Ibaraki (1984-05)
Axenic, Clonal, S. Suda (1984-05)
Identified by: S. Suda
Culture conditions: C, 20° C, 22 μE/m² sec, 2M
Characteristics: Freshwater
H-3-4-2
- Chlamydomonas monadina* Stein var. *monadina*
438
Lake Kasumigaura / Ibaraki (1983-07)
Axenic, Clonal, S. Suda (1983-07)
Identified by: S. Suda
Culture conditions: C, 20° C, 22 μE/m² sec, 2M
Characteristics: Freshwater
Kas-7

Chlamydomonas monticola S.Watanabe
157

Mt. Shiroumadake / Nagano (1980-08)
Unialgal, Clonal, S.Watanabe
Identified by: S.Watanabe
Culture conditions: C(S), 20° C, 4 µE/m² sec, 3M,
(25° C, 30 µE/m² sec)
Characteristics: Soil
KUC80-4
References: 277, 416

* *Chlamydomonas neglecta* (Pascher) Korshikov
See *Chlorogonium neglectum* Pascher

Chlamydomonas parkeae Ettl
440

Izumi Bay / Nagasaki (1986-03)
Unialgal, Clonal, S.Suda (1986-03)
Identified by: S.Suda
Culture conditions: f/2, 20° C, 22 µE/m² sec, 2M
Characteristics: Marine
I-29
References: 135, 294, 300

441

Hachinohe Harbor / Aomori (1985-01)
Axenic, Clonal, S.Suda (1985-02)
Identified by: S.Suda
Culture conditions: f/2, 20° C, 22 µE/m² sec, 2M
Characteristics: Marine
HH-5
Reference: 294

Chlamydomonas pulsatilla Wollenweber
122

Muroran / Hokkaido (1966-05)
IAM C-385, Axenic, Clonal, T.Ichimura (1966-05)
Identified by: T.Ichimura
Culture conditions: P35, 20° C, 4 µE/m² sec, 2M,
(25° C, 30 µE/m² sec)
Characteristics: Freshwater
MKF-50
References: 80, 399, 417

Chlamydomonas tetragama (Bohlin) Ettl
446

Tsuchiura / Ibaraki (1985-04)
Axenic, Clonal, S.Suda (1985-04)
Identified by: S.Suda
Culture conditions: C, 20° C, 22 µE/m² sec, 2M
Characteristics: Freshwater, Neotype strain of

Chlamydomonas tetragama (Bohlin) Ettl,
Formerly identified as *Chlorogonium*
metamorphum Skuja, *rbcL* gene (AJ001880)
413D4-4
References: 240, 255, 358

Chlorarachnion reptans Geitler
624

Puerto Penasco / Mexico
CCAP 815/1, Unialgal, Norris (1966)
Culture conditions: ESM, 20° C, 32 µE/m² sec, 2M
Characteristics: Marine

Chlorella fusca Shihira et Krauss var. *fusca*
685

IAM C-101, Unialgal, Clonal, R.A.Lewin
Culture conditions: C(S), 20° C, 4 µE/m² sec, 3M,
(25° C, 30 µE/m² sec)
Characteristics: Freshwater, Type strain

* *Chlorella pyrenoidosa* Chick

See *Graesiella emersonii*
(Shihira et Kraus) Nozaki et al.

Chlorella saccharophila (Krueger) Migula
640

Otarunai River / Hokkaido (1987-07)
Unialgal, F.Kasai (1987-07)
Identified by: F.Kasai
Culture conditions: C, 10° C, 6 µE/m² sec, 6M,
(10° C, 15 µE/m² sec)
Characteristics: Freshwater
Tst-8-2
Reference: 338

Chlorella vulgaris Beijerinck var. *vulgaris*
227

IAM C-30, Axenic, Clonal, A.Watanabe
Reidentified by: H.Nozaki
Culture conditions: C(S), 20° C, 4 µE/m² sec, 3M,
(25° C, 30 µE/m² sec)
Characteristics: Freshwater, *COXI* gene (D63763,
AB011523)
References: 66, 80, 96, 122, 131, 170, 182, 248, 271,
356, 366, 374, 417, 440

641

Miyata River / Ibaraki (1987-02)
Axenic, Clonal, F.Kasai (1987-03)
Identified by: F.Kasai
Culture conditions: C, 20° C, 8 µE / m² sec, 6M

- Characteristics: Freshwater
1st-3-26
References: 337, 338
- 642
Miyata River / Ibaraki (1987-02)
Unialgal, Clonal, F.Kasai (1987-03)
Identified by: F.Kasai
Culture conditions: C, 20° C, 8 µE/m² sec, 6M
Characteristics: Freshwater
1st-2-17
References: 337, 338
- 686
Delft / Holland
IAM C-207, Unialgal, Clonal, M.W.Beijerinck
(1892)
Culture conditions: C(S), 20° C, 4 µE/m² sec, 3M,
(25° C, 30 µE/m² sec)
Characteristics: Freshwater, Type strain
Reference: 248
- Chlorogonium capillatum* Nozaki et al.
692
Miyatoko Mire / Fukushima (1992-04)
Axenic, Clonal, H.Nozaki (1992-05)
Identified by: H.Nozaki
Culture conditions: AF-6, 10° C, 25 µE/m² sec, 1M
Characteristics: Freshwater, Type strain,
Monoecious, Isogamy, Paedogamy,
rbcL gene (AB010230)
92-912-1
Reference: 255
- 742
Czechoslovakia
UTEX 201, CCAP 12/4, Unialgal, Clonal, H.Meyer
Reidentified by: H.Nozaki
Culture conditions: AF-6, 20° C, 12 µE/m² sec, 1M
Characteristics: Freshwater, Formerly identified as
Chlorogonium tetragamum Bohlin, *rbcL* gene
(AB010234)
Reference: 255
- 743
Leveret, MA / USA
UTEX 1643, Unialgal, Clonal, P.Kugrens
Reidentified by: H.Nozaki
Culture conditions: AF-6, 20° C, 12 µE/m² sec, 1M
Characteristics: Freshwater, Formerly identified as
Chlorogonium tetragamum Bohlin, *rbcL* gene
(AB010235)
- Reference: 255
- 744
Germany
UTEX 2160, Unialgal, Clonal
Reidentified by: H.Nozaki
Culture conditions: AF-6, 20° C, 12 µE/m² sec, 1M
Characteristics: Freshwater, Formerly identified as
Chlorogonium sp., *rbcL* gene (AB010236)
Reference: 255
- 745
Berlin / Germany
CCAP 12/2A, Unialgal, Clonal, Hartmann
Reidentified by: H.Nozaki
Culture conditions: AF-6, 20° C, 12 µE/m² sec, 1M
Characteristics: Freshwater, Formerly identified as
Chlorogonium elongatum Dangeard, *rbcL* gene
(AB010231)
Reference: 255
- 746
Cape Flats / South Africa
CCAP 12/2B, Unialgal, Clonal, George (1948)
Reidentified by: H.Nozaki
Culture conditions: AF-6, 20° C, 12 µE/m² sec, 1M
Characteristics: Freshwater, Formerly identified as
Chlorogonium elongatum Dangeard, *rbcL* gene
(AB010232)
Reference: 255
- 747
CCAP 12/5, Unialgal, Clonal, E.G.Pringsheim
Reidentified by: H.Nozaki
Culture conditions: AF-6, 20° C, 12 µE/m² sec, 1M
Characteristics: Freshwater, Formerly identified as
Chlorogonium sp., *rbcL* gene (AB010233)
Reference: 255
- 748
near Prague / Czechoslovakia
SAG 12-2e, Unialgal, Clonal, E.G.Pringsheim (1936)
Reidentified by: H.Nozaki
Culture conditions: AF-6, 20° C, 12 µE/m² sec, 1M
Characteristics: Freshwater, Formerly identified as
Chlorogonium elongatum (Dangeard) Dangeard,
rbcL gene (AB010237)
Reference: 255
- 749
SAG 47.84, Unialgal, Clonal, L.Provasoli
Reidentified by: H.Nozaki

- Culture conditions: AF-6, 20° C, 12 µE/m² sec, 1M
 Characteristics: Freshwater, Formerly identified as
Chlorogonium euchlorum Ehrenberg, *rbcL* gene
 (AB010238)
 Reference: 255
- 750
 Leveret, MA / USA
 SAG 4.93, Unialgal, Clonal, P.Kugrens
 Reidentified by: H.Nozaki
 Culture conditions: AF-6, 20° C, 12 µE/m² sec, 1M
 Characteristics: Freshwater, Formerly identified as
Chlorogonium tetragamum Bohlin, *rbcL* gene
 (AB010239)
 Reference: 255
- Chlorogonium elongatum* (Dangeard) Dangeard
 751
 Caldbeck / U.K.
 IAM C-293, UTEX 204, Unialgal, Clonal,
 E.G.Pringsheim
 Reidentified by: H.Nozaki
 Culture conditions: AF-6, 20° C, 12 µE/m² sec, 1M
 Characteristics: Freshwater, Formerly identified as
Chlorogonium acus, *rbcL* gene (AJ001881)
 Reference: 255
- 752
 Austin, TX / USA
 UTEX 2571, Unialgal, Clonal, M.Wood (1990)
 Reidentified by: H.Nozaki
 Culture conditions: AF-6, 20° C, 12 µE/m² sec, 1M
 Characteristics: Freshwater, Formerly identified as
Chlorogonium sp., *rbcL* gene (AB010240)
 Reference: 255
- 753
 Austin, TX / USA
 UTEX 2572, Unialgal, Clonal, M.Wood (1990)
 Reidentified by: H.Nozaki
 Culture conditions: AF-6, 20° C, 12 µE/m² sec, 1M
 Characteristics: Freshwater, Formerly identified as
Chlorogonium sp., *rbcL* gene (AB010241)
 Reference: 255
- Chlorogonium euchlorum* (Ehrenberg) Ehrenberg
 754
 Schickley, NE / USA
 UTEX 1639, Unialgal, Clonal, P.Kugrens
 Reidentified by: H.Nozaki
 Culture conditions: AF-6, 20° C, 12 µE/m² sec, 1M
 Characteristics: Freshwater, Formerly identified as
Chlorogonium elongatum Dangeard, *rbcL* gene
 (AB010226)
 Reference: 255
- 755
 Germany
 UTEX 2010, Unialgal, Clonal, D.G.Müller
 Reidentified by: H.Nozaki
 Culture conditions: AF-6, 20° C, 12 µE/m² sec, 1M
 Characteristics: Freshwater, Formerly identified as
Chlorogonium sp., *rbcL* gene (AB010227)
 Reference: 255
- 756
 Germany
 UTEX 2011, Unialgal, Clonal, D.G.Müller
 Reidentified by: H.Nozaki
 Culture conditions: AF-6, 20° C, 12 µE/m² sec, 1M
 Characteristics: Freshwater, Formerly identified as
Chlorogonium sp., *rbcL* gene (AB010228)
 Reference: 255
- 757
 Amiens / France
 CCAP 12/2C, Unialgal, Clonal, E.G.Pringsheim
 (1949)
 Reidentified by: H.Nozaki
 Culture conditions: AF-6, 20° C, 12 µE/m² sec, 1M
 Characteristics: Freshwater, Formerly identified as
Chlorogonium elongatum (Dangeard) Dangeard,
rbcL gene (AB010224)
 Reference: 255
- 758
 Hirschberg / former Czechoslovakia
 CCAP 12/3, Unialgal, Clonal, Mainx (1936)
 Reidentified by: H.Nozaki
 Culture conditions: AF-6, 20° C, 12 µE/m² sec, 1M
 Characteristics: Freshwater, Formerly identified as
Chlorogonium euchlorum Ehrenberg, *rbcL* gene
 (AJ001882)
 Reference: 255
- 759
 CCAP 12/6, Unialgal, Clonal, E.G.Pringsheim
 Reidentified by: H.Nozaki
 Culture conditions: AF-6, 20° C, 12 µE/m² sec, 1M
 Characteristics: Freshwater, Formerly identified as
Chlorogonium sp., *rbcL* gene (AB010225)
 Reference: 255

- 760
Cape Flats / South Africa, D.K.Vlei
SAG 12-2d, Unialgal, Clonal, E.G.Pringsheim (1951)
Reidentified by: H.Nozaki
Culture conditions: AF-6, 20° C, 12 µE/m² sec, 1M
Characteristics: Freshwater, Formerly identified as
Chlorogonium elongatum (Dangeard) Dangeard,
rbcL gene (AB010229)
Reference: 255
- Chlorogonium fusiforme* Matvienko
123
Niseko / Hokkaido (1964-07)
IAM C-349, Axenic, Clonal, T.Ichimura (1964-07)
Reidentified by: H.Nozaki
Culture conditions: AF-6, 20° C, 4 µE/m² sec, 2M,
(25° C, 30 µE/m² sec)
Characteristics: Freshwater, Homothallic, Formerly
identified as *Chlorogonium metamorphum* Skuja,
rbcL gene (AB010242)
MKF-14
References: 80, 240, 255
- Chlorogonium kasakii* Nozaki
761
Cumbria / U.K.
CCAP 12/8, Unialgal, Clonal, Jaworski
Reidentified by: H.Nozaki
Culture conditions: AF-6, 20° C, 12 µE/m² sec, 1M
Characteristics: Freshwater, Formerly identified as
Chlorogonium sp., Type strain, *rbcL* gene
(AB010244)
Reference: 255
- * *Chlorogonium metamorphum* Skuja
123
See *Chlorogonium fusiforme* Matvienko
- * *Chlorogonium metamorphum* Skuja
446
See *Chlamydomonas tetragama* (Bohlin) Ettl
- Chlorogonium neglectum* Pascher
Syn. *Chlamydomonas neglecta* (Pascher) Korshikov
439
Tsukuba / Ibaraki (1984-05)
Axenic, Clonal, S.Suda (1984-05)
Reidentified by: H.Nozaki
Culture conditions: C, 20° C, 22 µE/m² sec, 2M
Characteristics: Freshwater, *rbcL* gene (AB010243)
T-4-19
- Reference: 255
- Chloromonas insignis* (Anachin) Gerloff et Ettl
447
Lake Kasumigaura / Ibaraki (1983-08)
Axenic, Clonal, S.Suda (1983-08)
Identified by: S.Suda
Culture conditions: C, 20° C, 22 µE/m² sec, 2M
Characteristics: Freshwater, *rbcL* gene (AB022226)
Kas-8
Reference: 183
- Chlorosarcinopsis caeca* S.Watanabe
160
Tottori (1972-05)
Unialgal, Non-clonal, S.Watanabe
Identified by: S.Watanabe
Culture conditions: C(S), 20° C, 4 µE/m² sec, 3M,
(25° C, 30 µE/m² sec)
Characteristics: Soil
TOT-24
Reference: 416
- Chlorosarcinopsis delicata* S.Watanabe
153
Kyoto / Kyoto (1975-04)
Unialgal, Clonal, S.Watanabe
Identified by: S.Watanabe
Culture conditions: C(S), 20° C, 4 µE/m² sec, 3M,
(25° C, 30 µE/m² sec)
Characteristics: Soil
KUC3-6
Reference: 416
- Chroomonas caudata* Geitler
712
Funada-ike / Chiba (1985-09)
Unialgal, Clonal, M.Erata (1985-09)
Identified by: M.Erata
Culture conditions: AF-6, 15° C, 15 µE/m² sec, 1M
Characteristics: Freshwater
#00171
Reference: 24
- Chroomonas coerulea* (Geitler) Skuja
713
Sugadaira / Nagano (1985-11)
Unialgal, Clonal, M.Erata (1985-11)
Identified by: M.Erata
Culture conditions: AF-6, 15° C, 15 µE/m² sec, 1M
Characteristics: Freshwater

- #00191
References: 22, 24
- 714
Sugadaira / Nagano (1985-11)
Unialgal, Clonal, M.Erata (1985-11)
Identified by: M.Erata
Culture conditions: AF-6, 15° C, 15 µE/m² sec, 1M
Characteristics: Freshwater
#00217
References: 22, 24, 25
- Chroomonas collegionis* Butcher
703
River Thames, Essex / U.K.
CCAP 978/11, Unialgal, Clonal, B.W.Butcher (1961)
Identified by: B.W.Butcher
Culture conditions: ESM, 15° C, 15 µE/m² sec, 1M
Characteristics: Marine, Type strain
- Chroomonas dispersa* Butcher
704
Bristol Channel / U.K.
CCAP 978/10, Unialgal, Clonal, B.W.Butcher
(1960-08)
Identified by: B.W.Butcher
Culture conditions: ESM, 15° C, 15 µE/m² sec, 1M
Characteristics: Marine, Type strain
- Chroomonas nordstedtii* Hansgirg
706
Sugadaira / Nagano (1976-09)
Axenic, Clonal, I.Inouye (1976-09)
Identified by: M.Erata
Culture conditions: AF-6, 15° C, 15 µE/m² sec, 1M
Characteristics: Freshwater
#00165
References: 22, 23, 24
- 707
Funada-ike / Chiba (1985-09)
Unialgal, Clonal, M.Erata (1985-09)
Identified by: M.Erata
Culture conditions: AF-6, 15° C, 15 µE/m² sec, 1M
Characteristics: Freshwater
#00173
References: 24, 25
- 708
Sapporo / Hokkaido (1987-09)
Unialgal, Clonal, M.Erata (1987-09)
Identified by: M.Erata
- Culture conditions: AF-6, 15° C, 15 µE/m² sec, 1M
Characteristics: Freshwater
#00324
Reference: 24
- 709
Sapporo / Hokkaido (1987-09)
Unialgal, Clonal, M.Erata (1987-09)
Identified by: M.Erata
Culture conditions: AF-6, 15° C, 15 µE/m² sec, 1M
Characteristics: Freshwater
#00330
Reference: 24
- 710
Sapporo / Hokkaido (1987-09)
Unialgal, Clonal, M.Erata (1987-09)
Identified by: M.Erata
Culture conditions: AF-6, 15° C, 15 µE/m² sec, 1M
Characteristics: Freshwater
#00331
Reference: 24
- 711
Mitsukaido / Ibaraki (1987-01)
Unialgal, Clonal, S.Suda (1987-12)
Identified by: M.Erata
Culture conditions: AF-6, 15° C, 15 µE/m² sec, 1M
Characteristics: Freshwater
#00354
- Chroomonas placoidea* Butcher
705
Yorkshire / U.K.
CCAP 978/8, Unialgal, Clonal, B.W.Butcher (1959)
Identified by: B.W.Butcher
Culture conditions: ESM, 15° C, 15 µE/m² sec, 1M
Characteristics: Marine, Type strain
Reference: 23
- Chrysochromulina hirta* Manton
741
Chiba Port / Chiba (1986-06)
Unialgal, Clonal, M.Kawachi (1986-07)
Identified by: M.Kawachi
Culture conditions: ESM, 15° C, 35 µE/m² sec, 1M
Characteristics: Marine, Unstable, Untransportable
CH1
- Chrysochromulina parva* Lackey
562
NIES / Tsukuba (1992-02)

- Unialgal, Clonal, N.Hatakeyama (1992-03)
 Identified by: M.Kawachi
 Culture conditions: AF-6, 15° C, 35 µE/m² sec, 1M
 Characteristics: Freshwater, Unstable,
 Untransportable
- Closterium acerosum* Ehrenberg ex Ralfs
 124
 Daramshara / Nepal (1965-10)
 Axenic, Clonal, T.Ichimura
 Identified by: T.Ichimura
 Culture conditions: C, 20° C, 8 µE/m² sec, 3M,
 (20° C, 12 µE/m² sec)
 Characteristics: Freshwater
 N-20-1
 Reference: 76
- 125
 Rukumkot / Nepal (1965-10)
 Axenic, Clonal, T.Ichimura
 Identified by: T.Ichimura
 Culture conditions: C, 20° C, 8 µE/m² sec, 3M,
 (20° C, 12 µE/m² sec)
 Characteristics: Freshwater
 N-25-22
 Reference: 76
- 127
 Sapporo / Hokkaido
 IAM C-435, Axenic, Clonal, Y.Nishihama
 Identified by: Y.Nishihama
 Culture conditions: C, 20° C, 8 µE/m² sec, 3M,
 (20° C, 12 µE/m² sec)
 Characteristics: Freshwater, Homothallic
 H-2-2
 References: 76, 80
- 448
 IAM C-314, UTEX 1075, Axenic, Clonal
 Culture conditions: C, 20° C, 8 µE/m² sec, 3M,
 (20° C, 12 µE/m² sec)
 Characteristics: Freshwater
 Reference: 80
- Closterium aciculare* T.West
 var. *subpronum* W. et G.S.West
 258
 Lake Biwa / Shiga (1983-12)
 Axenic, Clonal, M.M.Watanabe (1983-12)
 Identified by: M.M.Watanabe
 Culture conditions: CA, 20° C, 32 µE/m² sec, 2M
- Characteristics: Water bloom, Freshwater,
 Heterothallic, Mating type +, Crosses with
 NIES-259 and NIES-260
 Bca-25
 Reference: 16
- 259
 Lake Biwa / Shiga (1983-12)
 Axenic, Clonal, M.M.Watanabe (1983-12)
 Identified by: M.M.Watanabe
 Culture conditions: CA, 20° C, 32 µE/m² sec, 2M
 Characteristics: Water bloom, Freshwater,
 Heterothallic, Mating type -, Crosses with
 NIES-258
 Bca-26
- Closterium calosporum* Wittrock var. *calosporum*
 271
 Vermont / U.S.A.
 IAM C-318, Axenic, Clonal, P.W.Cook
 Culture conditions: AF-6, 20° C, 8 µE/m² sec, 3M,
 (20° C, 25 µE/m² sec)
 Characteristics: Freshwater
 References: 80, 86, 376, 377
- Closterium calosporum* Wittrock
 var. *galiciense* Gutwinski
 128
 Ibaraki
 Axenic, Clonal, M.M.Watanabe
 Identified by: M.Watanabe
 Culture conditions: CA, 20° C, 8 µE/m² sec, 3M,
 (20° C, 25 µE/m² sec)
 Characteristics: Freshwater, Heterothallic,
 Mating type -, Crosses with NIES-162
 IB-21-20
- 162
 Ibaraki
 Unialgal, Clonal, M.M.Watanabe
 Identified by: M.M.Watanabe
 Culture conditions: CA, 20° C, 8 µE/m² sec, 3M,
 (25° C, 15 µE/m² sec)
 Characteristics: Freshwater, Heterothallic,
 Mating type +, Crosses with NIES-128, NIES-163
 and NIES-168
 IB-21-21
- 163
 Ginama / Okinawa (1973-06)
 IAM C-455, Axenic, Clonal, T.Ichimura (1973-10)

Identified by: M.Watanabe
Culture conditions: CA, 20° C, 8 µE/m² sec, 3M,
(20° C, 25 µE/m² sec)
Characteristics: Freshwater, Heterothallic,
Mating type -, Crosses with NIES-162, NIES-164
and NIES-165
R-5-3
References: 86, 376, 377

164
Ginama / Okinawa (1973-06)
IAM C-454, Unialgal, Clonal, T.Ichimura (1973-10)
Identified by: M.Watanabe
Culture conditions: CA, 20° C, 8 µE/m² sec, 3M,
(20° C, 25 µE/m² sec)
Characteristics: Freshwater, Heterothallic
Mating type +, Crosses with NIES-163 and
NIES-166
R-5-2
References: 86, 376, 377

165
Iriomote Isl. / Okinawa (1973-03)
IAM C-457, Axenic, Clonal, T.Ichimura (1973-10)
Identified by: M.Watanabe
Culture conditions: CA, 20° C, 8 µE/m² sec, 3M,
(25° C, 15 µE/m² sec)
Characteristics: Freshwater, Heterothallic,
Mating type +, Crosses with NIES-163, NIES-166
and NIES-168
R-11-6
References: 86, 376, 377

166
Kagawa-cho / Kagawa (1974-09)
Axenic, Clonal, T.Ichimura
Identified by: M.Watanabe
Culture conditions: CA, 20° C, 8 µE/m² sec, 3M,
(20° C, 25 µE/m² sec)
Characteristics: Freshwater, Heterothallic,
Mating type -, Crosses with NIES-164, NIES-165
and NIES-167
J5-56-11

167
Kagawa-cho / Kagawa (1974-09)
Axenic, Clonal, T.Ichimura
Identified by: M.Watanabe
Culture conditions: CA, 20° C, 8 µE/m² sec, 3M,
(20° C, 25 µE/m² sec)
Characteristics: Freshwater, Heterothallic,

Mating type +, Crosses with NIES-166
J5-56-12

168
Iriomote Isl. / Okinawa (1973-03)
Axenic, Clonal, T.Ichimura
Identified by: M.Watanabe
Culture conditions: CA, 20° C, 8 µE/m² sec, 3M,
(20° C, 25 µE/m² sec)
Characteristics: Freshwater, Heterothallic,
Mating type -, Crosses with NIES-165
R-11-5
References: 86, 376, 377

Closterium calosporum Wittrock
var. *himalayense* M.Watanabe

169
Shewaden / Nepal (1972-06)
Axenic, Clonal, M.M.Watanabe
Identified by: M.Watanabe
Culture conditions: CA, 20° C, 8 µE/m² sec, 3M,
(20° C, 25 µE/m² sec)
Characteristics: Freshwater, Homothallic
N-134-5
References: 376, 377

170
Suke / Nepal (1972-06)
Unialgal, Clonal, M.M.Watanabe
Identified by: M.Watanabe
Culture conditions: CA, 20° C, 8 µE/m² sec, 3M,
(20° C, 25 µE/m² sec)
Characteristics: Freshwater, Homothallic
N-143-19

171
Suke / Nepal (1972-06)
Unialgal, Clonal, M.M.Watanabe
Identified by: M.Watanabe
Culture conditions: CA, 20° C, 8 µE/m² sec, 3M,
(20° C, 25 µE/m² sec)
Characteristics: Freshwater, Homothallic
N-147-3
References: 124, 376

336
Suke / Nepal (1972-06)
Axenic, Clonal, M.M.Watanabe
Identified by: M.Watanabe
Culture conditions: CA, 25° C, 15 µE/m² sec, 2M
Characteristics: Freshwater, Homothallic

- N-147-12
Reference: 376
- Closterium ehrenbergii* Meneghini ex Ralfs
228
Ebina / Kanagawa (1975-12)
Axenic, Clonal, T.Ichimura
Identified by: T.Ichimura
Culture conditions: C, 20° C, 8 µE/m² sec, 3M,
(25° C, 15 µE/m² sec)
Characteristics: Freshwater, Heterothallic,
Mating type +, Mating group B
Crosses with NIES-229
KK-33-1
References: 46, 47, 67, 78, 79, 81, 82, 84, 124, 125,
135
- 229
Ebina / Kanagawa (1975-12)
Axenic, Clonal, T.Ichimura
Identified by: T.Ichimura
Culture conditions: C, 20° C, 8 µE/m² sec, 3M,
(20° C, 12 µE/m² sec)
Characteristics: Freshwater, Heterothallic,
Mating type -, Mating group B
Crosses with NIES-228
KK-33-6
References: 46, 47, 67, 78, 79, 81, 82, 84, 124, 125
- Closterium gracile* Brébisson ex Ralfs
179
Kathmandu / Nepal (1968-05)
IAM C-444, Axenic, Clonal, T.Ichimura
Identified by: T.Ichimura
Culture conditions: CA, 20° C, 8 µE/m² sec, 3M,
(20° C, 12 µE/m² sec)
Characteristics: Freshwater, Heterothallic,
Mating type +, Crosses with NIES-180
N-90-58
References: 76, 80
- 180
Kathmandu / Nepal (1968-05)
IAM C-445, Unialgal, Clonal, T.Ichimura
Identified by: T.Ichimura
Culture conditions: CA, 20° C, 8 µE/m² sec, 3M,
(20° C, 12 µE/m² sec)
Characteristics: Freshwater, Heterothallic,
Mating type -, Crosses with NIES-179
N-90-59
References: 76, 80
- Closterium incurvum* Brébisson
181
Dhorpatan / Nepal (1965-11)
IAM C-438, Unialgal, Clonal, T.Ichimura
Identified by: T.Ichimura
Culture conditions: CA, 20° C, 8 µE/m² sec, 3M,
(20° C, 25 µE/m² sec)
Characteristics: Freshwater, Homothallic
N-34-2
References: 76, 80
- 337
Nawakot / Nepal (1965-10)
Unialgal, Non-clonal, T.Ichimura
Identified by: T.Ichimura
Culture conditions: CA, 20° C, 8 µE/m² sec, 3M,
(20° C, 25 µE/m² sec)
Characteristics: Freshwater, Homothallic
N-12-92
Reference: 76
- Closterium moniliferum* Ehrenberg ex Ralfs
var. *moniliferum*
172
Nepal
Unialgal, Non-clonal
Culture conditions: C, 20° C, 8 µE/m² sec, 3M,
(20° C, 25 µE/m² sec)
Characteristics: Freshwater, Homothallic
N-100-1
- 173
Kitaadachi-gun / Saitama (1969-01)
IAM C-432, Axenic, Clonal, T.Ichimura (1969-03)
Identified by: T.Ichimura
Culture conditions: C, 20° C, 8 µE/m² sec, 3M,
(20° C, 12 µE/m² sec)
Characteristics: Freshwater, Homothallic
S-1-22
Reference: 80
- 174
Ghorepani / Nepal (1965-12)
Unialgal, Clonal, T.Ichimura
Identified by: T.Ichimura
Culture conditions: C, 20° C, 8 µE/m² sec, 3M,
(20° C, 12 µE/m² sec)
Characteristics: Freshwater, Homothallic
N-76-30
Reference: 76

Closterium moniliferum Ehrenberg ex Ralfs
var. *submoniliferum* (Woronichin) Krieger

182

Kitaadachi-gun / Saitama (1969-01)
IAM C-433, Axenic, Clonal, T.Ichimura (1969-03)

Identified by: T.Ichimura

Culture conditions: C, 20° C, 8 μ E/m² sec, 3M,
(20° C, 12 μ E/m² sec)

Characteristics: Freshwater, Heterothallic,

Mating type +, Crosses with NIES-183

S-1-13

References: 76, 80

183

Kitaadachi-gun / Saitama (1969-01)

IAM C-434, Unialgal, Clonal, T.Ichimura (1969-03)

Identified by: T.Ichimura

Culture conditions: C, 20° C, 8 μ E/m² sec, 3M,
(20° C, 12 μ E/m² sec)

Characteristics: Freshwater, Heterothallic,

Mating type -, Crosses with NIES-182

S-1-24

References: 76, 80

Closterium navicula (Brébisson) Lütkenmüller

175

Chingkhola / Nepal (1965-11)

IAM C-443, Unialgal, Clonal, T.Ichimura

Identified by: T.Ichimura

Culture conditions: AF-6, 20° C, 8 μ E/m² sec, 3M,
(20° C, 25 μ E/m² sec)

Characteristics: Freshwater, Homothallic

N-49-7

References: 76, 80

176

Ghorepani / Nepal (1965-12)

Axenic, Clonal, T.Ichimura

Identified by: T.Ichimura

Culture conditions: C, 20° C, 8 μ E/m² sec, 3M,
(20° C, 25 μ E/m² sec)

Characteristics: Freshwater, Homothallic

N-75-10

Reference: 76

177

Billethadi / Nepal (1965-12)

Unialgal, Clonal, T.Ichimura

Identified by: T.Ichimura

Culture conditions: CA, 20° C, 8 μ E/m² sec, 3M,
(20° C, 25 μ E/m² sec)

Characteristics: Freshwater, Homothallic
N-79-26

Reference: 76

178

Shewaden / Nepal (1972-06)

Unialgal, Clonal, M.M.Watanabe (1974)

Culture conditions: CA, 20° C, 8 μ E/m² sec, 3M,
(20° C, 25 μ E/m² sec)

Characteristics: Freshwater

N-134-15

Closterium peracerosum-strigosum-littorale complex

51

Katsuta / Ibaraki (1974-08)

Unialgal, Clonal, M.M.Watanabe (1974-08)

Identified by: M.M.Watanabe

Culture conditions: CA, 15° C, 10 μ E/m² sec, 3M,
(20° C, 25 μ E/m² sec)

Characteristics: Indicator, Freshwater, Heterothallic,

Mating type +, Group II A

IB-4-2

References: 387, 392, 393, 394

52

Katsuta / Ibaraki (1974-08)

Axenic, Clonal, M.M.Watanabe (1974-08)

Identified by: M.M.Watanabe

Culture conditions: C, 15° C, 10 μ E/m² sec, 3M,
(20° C, 25 μ E/m² sec)

Characteristics: Indicator, Freshwater, Heterothallic,

Mating type -, Group II A

IB-4-9

References: 387, 392, 393, 394

53

Katsuta / Ibaraki (1974-08)

Axenic, Clonal, M.M.Watanabe (1974-08)

Identified by: M.M.Watanabe

Culture conditions: C, 15° C, 10 μ E/m² sec, 3M,
(20° C, 25 μ E/m² sec)

Characteristics: Freshwater, Heterothallic,

Mating type +, Group II A

IB-6-8

References: 387, 392, 393, 394

54

Katsuta / Ibaraki (1974-08)

Axenic, Clonal, M.M.Watanabe (1974-08)

Identified by: M.M.Watanabe

Culture conditions: C, 15° C, 10 μ E/m² sec, 3M,

- (20° C, 25 µE/m² sec)
 Characteristics: Freshwater, Heterothallic,
 Mating type -, Group II A
 IB-6-9
 References: 387, 392, 393
- 55
 Katsuta / Ibaraki (1975-05)
 Axenic, Clonal, M.M.Watanabe (1975-05)
 Identified by: M.M.Watanabe
 Culture conditions: C, 15° C, 10 µE/m² sec, 3M,
 (20° C, 25 µE/m² sec)
 Characteristics: Freshwater, Heterothallic,
 Mating type -, Group II C
 IB-8-15
 References: 387, 392, 393
- 56
 Katsuta / Ibaraki (1975-05)
 Axenic, Clonal, M.M.Watanabe (1975-05)
 Identified by: M.M.Watanabe
 Culture conditions: C, 15° C, 10 µE/m² sec, 3M,
 (20° C, 25 µE/m² sec)
 Characteristics: Freshwater, Heterothallic,
 Mating type -, Group II A
 IB-8-24
 References: 277, 387, 392, 393
- 57
 Katsuta / Ibaraki (1975-05)
 Axenic, Clonal, M.M.Watanabe (1975-05)
 Identified by: M.M.Watanabe
 Culture conditions: C, 15° C, 10 µE/m² sec, 3M,
 (20° C, 25 µE/m² sec)
 Characteristics: Freshwater, Heterothallic,
 Mating type +, Group II A
 IB-8-25
 References: 277, 387, 392, 393
- 58
 Mito / Ibaraki (1975-06)
 Unialgal, Clonal, M.M.Watanabe (1975-06)
 Identified by: M.M.Watanabe
 Culture conditions: C, 15° C, 10 µE/m² sec, 3M,
 (20° C, 25 µE/m² sec)
 Characteristics: Freshwater, Heterothallic,
 Mating type -, Group II A
 IB-10-1
 References: 387, 392, 393
- 59
 Mito / Ibaraki (1975-06)
 Axenic, Clonal, M.M.Watanabe (1975-06)
 Identified by: M.M.Watanabe
 Culture conditions: C, 15° C, 10 µE/m² sec, 3M,
 (20° C, 25 µE/m² sec)
 Characteristics: Freshwater, Heterothallic,
 Mating type +, Group II A
 IB-10-2
 References: 387, 392, 393
- 60
 Mito / Ibaraki (1975-06)
 Axenic, Clonal, M.M.Watanabe (1975-06)
 Identified by: M.M.Watanabe
 Culture conditions: C, 15° C, 10 µE/m² sec, 3M,
 (20° C, 25 µE/m² sec)
 Characteristics: Freshwater, Heterothallic,
 Mating type +, Group II B
 IB-12-1
 References: 387, 392, 393
- 61
 Mito / Ibaraki (1975-06)
 Axenic, Clonal, M.M.Watanabe (1975-06)
 Identified by: M.M.Watanabe
 Culture conditions: C, 15° C, 10 µE/m² sec, 3M,
 (20° C, 25 µE/m² sec)
 Characteristics: Freshwater, Heterothallic,
 Mating type -, Group II B
 IB-12-2
 References: 387, 392, 393
- 62
 Katsuta / Ibaraki (1975-06)
 Axenic, Clonal, M.M.Watanabe (1975-06)
 Identified by: M.M.Watanabe
 Culture conditions: C, 15° C, 10 µE/m² sec, 3M,
 (20° C, 25 µE/m² sec)
 Characteristics: Freshwater, Heterothallic,
 Mating type +, Group II A
 IB-13-1
 References: 387, 392, 393
- 63
 Katsuta / Ibaraki (1975-06)
 Unialgal, Clonal, M.M.Watanabe (1975-06)
 Identified by: M.M.Watanabe
 Culture conditions: C, 15° C, 10 µE/m² sec, 3M,
 (20° C, 25 µE/m² sec)
 Characteristics: Freshwater, Heterothallic,

- Mating type –, Group II A
 IB-13-2
 References: 387, 392, 393
- 64
 Lake Kasumigaura / Ibaraki (1974-11)
 Unialgal, Clonal, M.M.Watanabe (1974-11)
 Identified by: M.M.Watanabe
 Culture conditions: C, 15° C, 10 μ E/m² sec, 3M,
 (20° C, 25 μ E/m² sec)
 Characteristics: Freshwater, Heterothallic,
 Mating type –, Group II B
 KAS-4-29
 References: 127, 128, 129, 212, 303, 310, 387, 392,
 393, 394
- 65
 Lake Kasumigaura / Ibaraki (1974-11)
 Axenic, Clonal, M.M.Watanabe (1974-11)
 Identified by: M.M.Watanabe
 Culture conditions: C, 15° C, 10 μ E/m² sec, 3M,
 (20° C, 25 μ E/m² sec)
 Characteristics: Freshwater, Heterothallic,
 Mating type +, Group II B
 KAS-4-30
 References: 127, 128, 129, 212, 303, 310, 387, 392,
 393, 394
- 66
 Piuthan / Nepal (1965-10)
 Unialgal, Clonal, T.Ichimura
 Identified by: T.Ichimura
 Culture conditions: C, 15° C, 10 μ E/m² sec, 3M,
 (20° C, 25 μ E/m² sec)
 Characteristics: Indicator, Freshwater, Heterothallic,
 Mating type +, Group I A
 N-13-1
 References: 75, 76, 387
- 67
 Damchan / Nepal (1965-11)
 Unialgal, Clonal, T.Ichimura
 Identified by: T.Ichimura
 Culture conditions: C, 15° C, 10 μ E/m² sec, 3M,
 (20° C, 25 μ E/m² sec)
 Characteristics: Indicator, Freshwater, Heterothallic,
 Mating type +, Group I B
 N-31-19
 References: 76, 219, 301, 302, 303, 304, 305, 306,
 307, 308, 309, 310, 387
- 68
 Damchan / Nepal (1965-11)
 Axenic, Clonal, T.Ichimura
 Identified by: T.Ichimura
 Culture conditions: C, 15° C, 10 μ E/m² sec, 3M,
 (20° C, 25 μ E/m² sec)
 Characteristics: Indicator, Freshwater, Heterothallic,
 Mating type –, Group I B
 N-31-24
 References: 76, 219, 301, 302, 303, 304, 305, 306,
 309, 310, 387
- 69
 Lake Teganuma / Chiba (1974-06)
 Unialgal, Clonal, M.M.Watanabe (1974-06)
 Identified by: M.M.Watanabe
 Culture conditions: C, 15° C, 10 μ E/m² sec, 3M,
 (20° C, 25 μ E/m² sec)
 Characteristics: Freshwater, Heterothallic,
 Mating type +, Group II B
 TG-2-21
 References: 387, 392, 393
- 70
 Lake Teganuma / Chiba (1974-06)
 Axenic, Clonal, M.M.Watanabe (1974-06)
 Identified by: M.M.Watanabe
 Culture conditions: C, 15° C, 10 μ E/m² sec, 3M,
 (20° C, 25 μ E/m² sec)
 Characteristics: Freshwater, Heterothallic,
 Mating type –, Group II B
 TG-2-22
 References: 387, 392, 393
- 261
 Katsuta / Ibaraki (1974-08)
 Unialgal, Clonal, M.M.Watanabe (1974-08)
 Identified by: M.M.Watanabe
 Culture conditions: C, 15° C, 10 μ E/m² sec, 3M,
 (20° C, 25 μ E/m² sec)
 Characteristics: Freshwater, Heterothallic,
 Mating type +, Group II C
 IB-8-14
 References: 387, 392, 393
- 262
 Piuthan / Nepal (1965-10)
 Unialgal, Clonal, T.Ichimura
 Identified by: T.Ichimura
 Culture conditions: C, 15° C, 10 μ E/m² sec, 3M,
 (20° C, 25 μ E/m² sec)

- Characteristics: Freshwater, Heterothallic,
Mating type –
N-13-4
References: 75, 76, 387
- Closterium pleurodermatum* West et West
449
Iriomote Isl. / Okinawa (1973-03)
IAM C-518, Unialgal, Clonal, T.Ichimura (1973-12)
Identified by: T.Ichimura
Culture conditions: AF-6, 20° C, 8 μ E/m² sec, 3M,
(25° C, 15 μ E/m² sec)
Characteristics: Freshwater
R-11-20
- Closterium praelongum* Brébisson
var. *brevius* (Nordstedt) Krieger
450
Nawakot / Nepal (1965-10)
IAM C-447, Axenic, Clonal, T.Ichimura
Identified by: T.Ichimura
Culture conditions: CA, 20° C, 8 μ E/m² sec, 3M,
(20° C, 25 μ E/m² sec)
Characteristics: Freshwater, Homothallic
N-12-3
References: 76, 80
- 451
Billethadi / Nepal (1965-12)
Unialgal, Clonal, , T.Ichimura
Identified by: T.Ichimura
Culture conditions: MG, 20° C, 8 μ E/m² sec, 3M,
(20° C, 25 μ E/m² sec)
Characteristics: Freshwater, Homothallic
N-78-8
Reference: 76
- Closterium pusillum* Hantzsch var. *maius* Raciborski
185
Billethadi / Nepal (1965-12)
IAM C-449, Unialgal, Clonal, T.Ichimura
Identified by: T.Ichimura
Culture conditions: CA, 20° C, 8 μ E/m² sec, 3M,
(20° C, 25 μ E/m² sec)
Characteristics: Freshwater, Heterothallic
N-79-19
References: 76, 80
- Closterium rostratum* Ehrenberg ex Ralfs
var. *subrostratum* (Krieger) Krieger
Syn. *Closterium subrostratum* Krieger
- 338
Kathmandu / Nepal (1968-05)
IAM C-446, Axenic, Clonal, T.Ichimura
Identified by: T.Ichimura
Culture conditions: CA, 20° C, 8 μ E/m² sec, 3M,
(20° C, 12 μ E/m² sec)
Characteristics: Freshwater, Homothallic
N-90-55
References: 76, 80
- Closterium selenastrum* M.Watanabe
339
Mt. Yonahadake / Okinawa (1972-10)
Unialgal, Clonal, T.Ichimura
Identified by: M.Watanabe
Culture conditions: CA, 20° C, 8 μ E/m² sec, 3M,
(25° C, 15 μ E/m² sec)
Characteristics: Indicator, Freshwater, Homothallic
R-9-40
References: 86, 376, 377
- 340
Mt. Yonahadake / Okinawa (1972-10)
Axenic, Clonal, T.Ichimura
Identified by: M.Watanabe
Culture conditions: CA, 20° C, 8 μ E/m² sec, 3M,
(25° C, 15 μ E/m² sec)
Characteristics: Indicator, Freshwater, Homothallic
R-9-42
References: 86, 377
- Closterium spinosporum* Hodgetts
var. *crassum* M.Watanabe
186
Lake Akan / Hokkaido (1973-09)
Axenic, Clonal, M.Watanabe
Identified by: M.Watanabe
Culture conditions: CA, 20° C, 8 μ E/m² sec, 3M,
(20° C, 25 μ E/m² sec)
Characteristics: Indicator, Freshwater, Homothallic,
Type strain
AK-46
References: 86, 376, 377
- 187
Mt. Yonahadake / Okinawa (1973-06)
IAM C-461, Unialgal, Clonal, T.Ichimura (1973-10)
Identified by: M.Watanabe
Culture conditions: CA, 20° C, 8 μ E/m² sec, 3M,
(20° C, 25 μ E/m² sec)
Characteristics: Freshwater, Homothallic

- R-9-13
References: 86, 376, 377
- 341
Mt. Yonahadake / Okinawa (1972-10)
Axenic, Clonal, T.Ichimura
Identified by: M.Watanabe
Culture conditions: CA, 20° C, 8 $\mu\text{E}/\text{m}^2\text{sec}$, 3M,
(20° C, 25 $\mu\text{E}/\text{m}^2\text{sec}$)
Characteristics: Indicator, Freshwater, Homothallic
R-9-12
References: 86, 376, 377
- Closterium spinosporum* Hodgetts
var. *malaysiense* M.Watanabe
188
Penang / Malaysia (1974-01)
Axenic, Clonal, M.Watanabe
Identified by: M.Watanabe
Culture conditions: CA, 20° C, 8 $\mu\text{E}/\text{m}^2\text{sec}$, 3M,
(20° C, 25 $\mu\text{E}/\text{m}^2\text{sec}$)
Characteristics: Indicator, Freshwater, Heterothallic,
Mating type +
M-10-1
References: 376, 377
- 189
Penang / Malaysia (1974-01)
Axenic, Clonal, M.Watanabe
Identified by: M.Watanabe
Culture conditions: CA, 20° C, 8 $\mu\text{E}/\text{m}^2\text{sec}$, 3M,
(20° C, 25 $\mu\text{E}/\text{m}^2\text{sec}$)
Characteristics: Indicator, Freshwater, Heterothallic,
Mating type –
M-10-4
References: 376, 377
- Closterium spinosporum* Hodgetts
var. *ryukyuense* M.Watanabe
191
Iriomote Isl. / Okinawa (1973-06)
Axenic, Clonal, T.Ichimura
Identified by: M.Watanabe
Culture conditions: CA, 20° C, 8 $\mu\text{E}/\text{m}^2\text{sec}$, 3M,
(20° C, 25 $\mu\text{E}/\text{m}^2\text{sec}$)
Characteristics: Indicator, Freshwater, Homothallic
R-12-3
References: 376, 377
- 192
Iriomote Isl. / Okinawa (1973-06)
Axenic, Clonal, T.Ichimura
Identified by: M.Watanabe
Culture conditions: CA, 20° C, 8 $\mu\text{E}/\text{m}^2\text{sec}$, 3M,
(25° C, 15 $\mu\text{E}/\text{m}^2\text{sec}$)
Characteristics: Indicator, Freshwater, Homothallic
R-12-6
References: 376, 377
- 193
Iriomote Isl. / Okinawa (1973-06)
Axenic, Clonal, T.Ichimura
Identified by: M.Watanabe
Culture conditions: CA, 20° C, 8 $\mu\text{E}/\text{m}^2\text{sec}$, 3M,
(20° C, 25 $\mu\text{E}/\text{m}^2\text{sec}$)
Characteristics: Indicator, Freshwater, Homothallic,
Giant cell
R-12-2G3
Reference: 376
- Closterium spinosporum* Hodgetts var. *spinosporum*
194
Tsukude-mura / Aichi (1972-10)
Axenic, Clonal, T.Ichimura
Identified by: M.Watanabe
Culture conditions: CAM, 20° C, 8 $\mu\text{E}/\text{m}^2\text{sec}$, 3M,
(25° C, 15 $\mu\text{E}/\text{m}^2\text{sec}$)
Characteristics: Indicator, Freshwater, Homothallic
A-2-22
References: 86, 376, 377
- 195
Tsukude-mura / Aichi (1972-10)
Unialgal, Clonal, M.M.Watanabe
Identified by: M.Watanabe
Culture conditions: CAM, 20° C, 8 $\mu\text{E}/\text{m}^2\text{sec}$, 3M,
(20° C, 25 $\mu\text{E}/\text{m}^2\text{sec}$)
Characteristics: Indicator, Freshwater, Homothallic
A-7-3
Reference: 377
- 196
Tsukude-mura / Aichi (1972-10)
Unialgal, Clonal, M.M.Watanabe
Identified by: M.Watanabe
Culture conditions: CAM, 20° C, 8 $\mu\text{E}/\text{m}^2\text{sec}$, 3M,
(20° C, 25 $\mu\text{E}/\text{m}^2\text{sec}$)
Characteristics: Indicator, Freshwater, Homothallic
A-7-6
Reference: 376

- 197
Tsukude-mura / Aichi (1972-10)
Unialgal, Clonal, M.M.Watanabe
Identified by: M.Watanabe
Culture conditions: CA, 20° C, 8 μE/m² sec, 3M,
(20° C, 25 μE/m² sec)
Characteristics: Indicator, Freshwater, Homothallic
A-13-4
References: 376, 377
- * *Closterium subrostratum* Krieger
See *Closterium rostratum* Ehrenberg ex Ralfs
var. *subrostratum* (Krieger) Krieger
- Closterium tumidum* Johnson
198
Billethadi / Nepal (1965-12)
IAM C-450, Unialgal, Clonal, T.Ichimura
Identified by: T.Ichimura
Culture conditions: C, 20° C, 8 μE/m² sec, 3M,
(20° C, 25 μE/m² sec)
Characteristics: Freshwater, Homothallic
N-79-11
References: 76, 80
- Closterium venus* Kützing ex Ralfs
199
Kathmandu / Nepal (1968)
Unialgal, Clonal, T.Ichimura
Identified by: T.Ichimura
Culture conditions: CA, 20° C, 8 μE/m² sec, 3M,
(20° C, 25 μE/m² sec)
Characteristics: Freshwater
N-90-48
- Closterium wallichii* Turner
200
Kitaadachi-gun / Saitama (1969-01)
IAM C-451, Unialgal, Clonal, T.Ichimura
Identified by: T.Ichimura
Culture conditions: C, 20° C, 8 μE/m² sec, 3M,
(20° C, 12 μE/m² sec)
Characteristics: Freshwater, Homothallic
S-1-0
Reference: 80
- 201
Lake Kasumigaura / Ibaraki (1983-09)
Axenic, Clonal, F.Kasai (1983-09)
Identified by: F.Kasai
Culture conditions: C, 20° C, 8 μE/m² sec, 3M,
(20° C, 12 μE/m² sec)
Characteristics: Indicator, Freshwater, Homothallic
F60-21
- 202
Ghasa / Nepal (1965-11)
Axenic, Clonal, T.Ichimura
Identified by: T.Ichimura
Culture conditions: C, 20° C, 8 μE/m² sec, 3M,
(20° C, 12 μE/m² sec)
Characteristics: Freshwater, Homothallic
N-63-0
Reference: 76
- Coelastrum astroideum* De Notaris
129
Lake Shoji / Yamanashi (1981-10)
TAC 56, Axenic, Clonal, M.Watanabe
Identified by: M.Watanabe
Culture conditions: C, 20° C, 8 μE/m² sec, 2M,
(25° C, 30 μE/m² sec)
Characteristics: Freshwater
TAN-56-7
- 130
Lake Shoji / Yamanashi (1981-08)
TAC 51-9A, Axenic, Clonal, M.Watanabe
Identified by: M.Watanabe
Culture conditions: C, 20° C, 8 μE/m² sec, 2M,
(25° C, 30 μE/m² sec)
Characteristics: Freshwater
TAN-51-9A
- 244
Lake Kasumigaura / Ibaraki (1983-08)
Unialgal, Clonal, F.Kasai (1983-08)
Identified by: M.Watanabe
Culture conditions: C(S), 20° C, 4 μE/m² sec, 3M,
(25° C, 30 μE/m² sec)
Characteristics: Freshwater
- 342
Lake Kawaguchi / Yamanashi (1981-10)
TAC 54, Unialgal, Clonal, M.Watanabe
Identified by: M.Watanabe
Culture conditions: C, 20° C, 8 μE/m² sec, 2M
Characteristics: Freshwater
TAN-54-1
- Coelastrum morus* W. et G.S.West
231

Hachijo Isl. / Tokyo (1984-04)
Axenic, Clonal, F.Kasai (1984-05)
Identified by: M.Watanabe
Culture conditions: C(S), 20° C, 4 µE/m² sec, 3M,
(25° C, 30 µE/m² sec)
Characteristics: Freshwater
F78-4-2
Reference: 126

Coelastrum proboscideum Bohlin

131
Near Tukucha / Nepal (1965-11)
IAM C-344, Axenic, Clonal, T.Ichimura (1969-07)
Identified by: T.Ichimura
Culture conditions: C(S), 20° C, 4 µE/m² sec, 3M,
(25° C, 30 µE/m² sec)
Characteristics: Freshwater
N-63-20
References: 80, 399

Coelastrum reticulatum (Dangeard) Senn

132
Lake Yamanaka / Yamanashi (1981-10)
TAC 53-5A, Axenic, Clonal, M.Watanabe
Identified by: M.Watanabe
Culture conditions: C, 20° C, 8 µE/m² sec, 2M,
(25° C, 30 µE/m² sec)
Characteristics: Indicator, Freshwater
TAN-53-5A

Coelastrum reticulatum (Dangeard) Senn

var. *reticulatum*
245
Lake Kasumigaura / Ibaraki (1983-10)
Axenic, Clonal, F.Kasai (1983-10)
Identified by: M.Watanabe
Culture conditions: C(S), 20° C, 4 µE/m² sec, 3M,
(25° C, 30 µE/m² sec)
Characteristics: Indicator, Freshwater
F63-3

Coolia monotis Meunier

343
Hachijo Isl. / Tokyo (1984-04)
Axenic, Clonal, S.Suda (1984-04)
Identified by: S.Suda
Culture conditions: ESM, 20° C, 12 µE/m² sec, 3M
Characteristics: Marine, Tide pool, Unstable,
Untransportable
8-1

615
Motobu / Okinawa (1993-06)
Unialgal, Clonal, H.Kobayashi (1993-06)
Identified by: Y.Fukuyo
Culture conditions: f/2, 20° C, 32 µE/m² sec, 1M
Characteristics: Toxic, Marine, Untransportable
CM-01

Cosmarium askenasyi Schmidle

768
Near Cairns, Queensland / Australia (1988-09)
Unialgal, Clonal, F.Kasai (1988-09)
Identified by: F.Kasai
Culture conditions: MG, 20° C, 8 µE/m² sec, 3M,
(25° C, 30 µE/m² sec)
Characteristics: Freshwater, Crosses with NIES-769
88-8-37

769
Near Cairns, Queensland / Australia (1988-09)
Unialgal, Clonal, F.Kasai (1988-09)
Identified by: F.Kasai
Culture conditions: MG, 20° C, 8 µE/m² sec, 3M,
(25° C, 30 µE/m² sec)
Characteristics: Freshwater, Crosses with NIES-768
88-8-38

770
Near Cairns, Queensland / Australia (1988-09)
Unialgal, Clonal, F.Kasai (1988-09)
Identified by: F.Kasai
Culture conditions: MG, 20° C, 8 µE/m² sec, 3M,
(25° C, 30 µE/m² sec)
Characteristics: Freshwater, Crosses with NIES-771
88-8-39

771
Near Cairns, Queensland / Australia (1988-09)
Unialgal, Clonal, F.Kasai (1988-09)
Identified by: F.Kasai
Culture conditions: MG, 20° C, 8 µE/m² sec, 3M,
(25° C, 30 µE/m² sec)
Characteristics: Freshwater, Crosses with NIES-770
88-8-40

Cosmarium contractum Kirchner

133
Lake Yamanaka / Yamanashi (1981-10)
TAC 53, Unialgal, Clonal, M.Watanabe
Identified by: M.Watanabe
Culture conditions: C, 20° C, 8 µE/m² sec, 2M,

- (20° C, 12 µE/m² sec)
 Characteristics: Indicator, Freshwater
 TAN-53-2
- Cosmarium dilatatum* Lütkenmüller
 in Tärnefeld et Grönblad
 839
 Ryoanji Temple / Kyoto (1998-06)
 Unialgal, Clonal, A. Gontcharov (1998-07)
 Identified by: A. Gontcharov
 Culture conditions: C, 20° C, 8 µE/m² sec, 3M,
 (25° C, 30 µE/m² sec)
 Characteristics: Freshwater
 Reference: 48
- Cosmarium hians* Borge
 452
 Lake Yamanaka / Yamanashi (1981-06)
 Axenic, Clonal, M.H. Watanabe (1981-06)
 Identified by: M.H. Watanabe
 Culture conditions: C, 20° C, 8 µE/m² sec, 2M
 Characteristics: Indicator, Freshwater
 YAMA-Cos-4
- Cosmocladium constrictum* (Archer) Archer
 248
 Lake Biwa / Shiga (1983-12)
 Axenic, Clonal, F. Kasai (1983-12)
 Identified by: M. Watanabe
 Culture conditions: C, 20° C, 8 µE/m² sec, 3M,
 (20° C, 12 µE/m² sec)
 Characteristics: Freshwater
 F75-2
- Cricosphaera roscoffensis*
 (Dangeard) Gayral et Fresnel
 8
 Osaka Bay / Osaka (1978-09)
 Axenic, Clonal, S. Yamochi
 Identified by: S. Yamochi
 Culture conditions: f/2, 20° C, 32 µE/m² sec, 1M
 Characteristics: Red tide, Marine,
 COXI gene (AB000117)
 OCri
 References: 65, 270
- Cryptomonas acuta* Butcher
 697
 Conway, N. Wales / U.K.
 CCAP 979/10, Unialgal, Clonal, B.W. Butcher
 Identified by: B.W. Butcher
- Culture conditions: ESM, 15° C, 15 µE/m² sec, 1M
 Characteristics: Marine, Type strain
 Reference: 23
- Cryptomonas irregularis* Butcher
 698
 Plymouth, Devon / U.K.
 CCAP 979/7, Unialgal, Clonal, B.W. Butcher (1960)
 Identified by: B.W. Butcher
 Culture conditions: ESM, 15° C, 15 µE/m² sec, 1M
 Characteristics: Marine, Type strain
 Reference: 23
- Cryptomonas ovata* Ehrenberg
 274
 Tsuchiura / Ibaraki (1982-10)
 Axenic, Clonal, M. Ishimitsu (1982-10)
 Identified by: M. Ishimitsu
 Culture conditions: VT, 10° C, 25 µE/m² sec, 2M
 Characteristics: Freshwater, COXI gene (AB009419)
 #00046
 References: 96, 106
- 275
 Tsuchiura / Ibaraki (1982-09)
 Axenic, Clonal, M. Ishimitsu (1982-09)
 Identified by: M. Ishimitsu
 Culture conditions: VT, 10° C, 25 µE/m² sec, 2M
 Characteristics: Freshwater
 #00042
 References: 106, 126
- Cryptomonas platyuris* Skuja
 276
 Higashihiroshima / Hiroshima (1983-08)
 Axenic, Clonal, M. Ishimitsu (1983-08)
 Identified by: M. Ishimitsu
 Culture conditions: VT, 10° C, 25 µE/m² sec, 1M
 Characteristics: Freshwater
 #00096
 Reference: 106
- 344
 Higashihiroshima / Hiroshima (1983-08)
 Axenic, Clonal, M. Ishimitsu (1983-08)
 Identified by: M. Ishimitsu
 Culture conditions: VT, 10° C, 25 µE/m² sec, 2M
 Characteristics: Freshwater
 #00103
 Reference: 106

Cryptomonas rostratiformis Skuja

277

Hongo / Hiroshima (1983-10)
Axenic, Clonal, M.Ishimitsu (1983-10)
Identified by: M.Ishimitsu
Culture conditions: VT, 15° C, 20 µE/m² sec, 1M
Characteristics: Freshwater
#00148
Reference: 106

278

Hongo / Hiroshima (1983-10)
Axenic, Clonal, M.Ishimitsu (1983-10)
Identified by: M.Ishimitsu
Culture conditions: VT, 15° C, 20 µE/m² sec, 1M
Characteristics: Freshwater
#00154
Reference: 106

345

Sugadaira / Nagano (1982-07)
Axenic, Clonal, M.Ishimitsu (1982-08)
Identified by: M.Ishimitsu
Culture conditions: VT, 10° C, 25 µE/m² sec, 2M
Characteristics: Freshwater
#00006
Reference: 106

Cryptomonas tetrapyrenoidosa Skuja

279

Higashihiroshima / Hiroshima (1983-08)
Axenic, Clonal, M.Ishimitsu (1983-08)
Identified by: M.Ishimitsu
Culture conditions: VT, 10° C, 25 µE/m² sec, 2M
Characteristics: Freshwater
#00099
Reference: 106

280

Sugadaira / Nagano (1982-07)
Axenic, Clonal, M.Ishimitsu (1982-08)
Identified by: M.Ishimitsu
Culture conditions: VT, 10° C, 25 µE/m² sec, 2M
Characteristics: Freshwater
#00014
References: 106, 126

281

Minamiizu / Shizuoka (1983-05)
Axenic, Clonal, M.Ishimitsu (1983-05)
Identified by: M.Ishimitsu
Culture conditions: VT, 5° C, 15 µE/m² sec, 2M

Characteristics: Freshwater

#00073

Reference: 106

282

Tsuchiura / Ibaraki (1982-09)
Axenic, Clonal, M.Ishimitsu (1982-09)
Identified by: M.Ishimitsu
Culture conditions: VT, 15° C, 20 µE/m² sec, 1M
Characteristics: Freshwater
#00056
References: 9, 10, 11, 106

346

Sugadaira / Nagano (1982-07)
Axenic, Clonal, M.Ishimitsu (1982-08)
Identified by: M.Ishimitsu
Culture conditions: VT, 5° C, 15 µE/m² sec, 2M
Characteristics: Freshwater
#00009
Reference: 106

347

Minamiizu / Shizuoka (1983-05)
Axenic, Clonal, M.Ishimitsu (1983-05)
Identified by: M.Ishimitsu
Culture conditions: VT, 5° C, 15 µE/m² sec, 2M
Characteristics: Freshwater
#00072
Reference: 106

348

Higashihiroshima / Hiroshima (1983-08)
Axenic, Clonal, M.Ishimitsu (1983-08)
Identified by: M.Ishimitsu
Culture conditions: VT, 10° C, 25 µE/m² sec, 2M
Characteristics: Freshwater
#00109
Reference: 106

Cyanidioschyzon merolae De Luca et al.

549

Unialgal, Non-clonal
Identified by: A.Merola et al.
Culture conditions: Allen, 20° C, 4 µE/m² sec, 6M,
(20° C, 12 µE/m² sec)
Characteristics: Acidophilic
3
Reference: 175

- Cyanidium caldarium* (Tilden) Geitler
250
See *Galdieria sulphuraria* (Galdieri) Merola
- 551
Unialgal, Non-clonal
Identified by: A.Merola et al.
Culture conditions: Allen, 20° C, 4 µE/m² sec, 6M,
(20° C, 12 µE/m² sec)
Characteristics: Acidophilic
086
Reference: 175
- Cyanophora paradoxa* Korshikov
547
England
UTEX 555, Axenic, Clonal, E.G.Pringsheim (1943)
Identified by: E.G.Pringsheim
Culture conditions: C, 20° C, 8 µE/m² sec, 2M,
(25° C, 30 µE/m² sec)
Characteristics: Alkaline water
- 763
Mitsukaido / Ibaraki (1987-01)
Axenic, Clonal, S.Suda (1991-08)
Identified by: S.Suda
Culture conditions: CSi, 20° C, 12 µE/m² sec, 2M
Characteristics: Freshwater, Indicator
S117
- Cyanophora tetracyanea* Korshikov
764
Mitsukaido / Ibaraki (1987-01)
Axenic, Clonal, S.Suda (1991-08)
Identified by: S.Suda
Culture conditions: CSi, 20° C, 12 µE/m² sec, 2M
Characteristics: Freshwater, Indicator
S118
- Cyclotella meneghiniana* Kützing
803
Mitsukaido / Ibaraki (1993-05)
Unialgal, Clonal, F.Kasai (1993-05)
Identified by: Y.Niiyama
Culture conditions: CSi, 15° C, 15 µE/m² sec, 1M
Characteristics: Freshwater
506-26
Reference: 123
- 804
Mitsukaido / Ibaraki (1993-05)
- Unialgal, Clonal, F.Kasai (1993-05)
Identified by: Y.Niiyama
Culture conditions: CSi, 15° C, 15 µE/m² sec, 1M
Characteristics: Freshwater
518-39
Reference: 123
- 805
Mitsukaido / Ibaraki (1993-06)
Unialgal, Clonal, F.Kasai (1993-06)
Identified by: Y.Niiyama
Culture conditions: CSi, 15° C, 15 µE/m² sec, 1M
Characteristics: Freshwater
613-2
Reference: 123
- Cylindrocystis brehissonii* (Ralfs) De Bary
var. *brehissonii*
349
Lake Onuma / Hokkaido (1967-06)
IAM C-354, Axenic, Clonal, M.Haga (1968-01)
Identified by: M.Haga
Culture conditions: C(S), 20° C, 8 µE/m² sec, 4M,
(25° C, 30 µE/m² sec)
Characteristics: Freshwater, Homothallic
6801-68
- Dictyochloropsis irregularis* Nakano et Isagi
378
Akkeshi / Hokkaido (1982-07)
Axenic, Clonal, Y.Isagi (1982-08)
Identified by: T.Nakano
Culture conditions: C(S), 20° C, 4 µE/m² sec, 3M,
(25° C, 30 µE/m² sec)
Characteristics: Aerial on the surface of the
bark of *Picea jezoensis*
CCHU-2227
Reference: 211
- Dictyosphaerium pulchellum* Wood
453
Lake Kasumigaura / Ibaraki (1988-12)
Unialgal, Clonal, T.Yanai (1988-12)
Identified by: Y.Niiyama
Culture conditions: MG, 15° C, 15 µE/m² sec, 2M
Characteristics: Freshwater
- Dimorphococcus lunatus* A.Brown
134
Ozegahara / Gunma (1983-08)
Unialgal, Clonal, F.Kasai (1983-09)

Identified by: M. Watanabe
Culture conditions: CA, 20° C, 4 $\mu\text{E}/\text{m}^2\text{sec}$, 2M,
(25° C, 30 $\mu\text{E}/\text{m}^2\text{sec}$)
Characteristics: Freshwater
34-5

135
Tsuchiura / Ibaraki (1983-10)
Axenic, Clonal, F. Kasai (1983-10)
Identified by: M. Watanabe
Culture conditions: CA, 20° C, 4 $\mu\text{E}/\text{m}^2\text{sec}$, 2M,
(25° C, 30 $\mu\text{E}/\text{m}^2\text{sec}$)
Characteristics: Freshwater
F-61-4
Reference: 399

Dinobryon divergens Imhof
284
Lake Biwa / Shiga (1983-12)
Unialgal, Non-clonal, F. Kasai (1983-12)
Identified by: F. Kasai
Culture conditions: AF-6/2, 15° C, 20 $\mu\text{E}/\text{m}^2\text{sec}$, 4M
Characteristics: Freshwater
F-75-26

Ditylum brightwellii (T. West) Grunow et Heurck
350
Shimoda / Shizuoka (1985-05)
Unialgal, Clonal, T. Sawaguchi (1985-05)
Identified by: T. Sawaguchi
Culture conditions: f/2, 5° C, 15 $\mu\text{E}/\text{m}^2\text{sec}$, 1M
Characteristics: Marine
KBB-10

Docidium undulatum Bailey var. *undulatum*
285
Oze / Fukushima (1983-08)
Unialgal, Clonal, F. Kasai (1983-09)
Identified by: F. Kasai
Culture conditions: SW(Bi), 20° C, 8 $\mu\text{E}/\text{m}^2\text{sec}$, 3M
Characteristics: Freshwater
41-11

Draparnaldia plumosa (Vaucher) Agardh
454
Shirai River / Sapporo (1987-10)
Unialgal, Non-clonal, F. Kasai (1987-10)
Identified by: F. Kasai
Culture conditions: C, 10° C, 6 $\mu\text{E}/\text{m}^2\text{sec}$, 3M,
(10° C, 15 $\mu\text{E}/\text{m}^2\text{sec}$)
Characteristics: Freshwater

2Tst-2-1
Reference: 338

Echinosphaeridium nordstedtii Lemmermann
137
Lake Kasumigaura / Ibaraki (1983-08)
Axenic, Clonal, F. Kasai (1983-08)
Identified by: M. Watanabe
Culture conditions: C(S), 20° C, 4 $\mu\text{E}/\text{m}^2\text{sec}$, 3M,
(25° C, 30 $\mu\text{E}/\text{m}^2\text{sec}$)
Characteristics: Indicator, Freshwater
F-56-3
References: 126, 399

Emiliana huxleyi (Lohmann) Hay et Mohler
837
Great Barrier Reef / Australia (1990-11)
Unialgal, Clonal, I. Inouye (1990-11)
Identified by: I. Inouye
Culture conditions: ESM, 20° C, 12 $\mu\text{E}/\text{m}^2\text{sec}$, 1M
Characteristics: Marine, Red tide
EH-01
References: 297, 311, 312, 313, 314

Eremosphaera gigas (Archer) Fott et Kalina
379
Shinobugaoka / Osaka (1968-11)
IAM C-338, Unialgal, Clonal, T. Ichimura (1969-01)
Identified by: T. Nakano
Culture conditions: C(S), 20° C, 4 $\mu\text{E}/\text{m}^2\text{sec}$, 3M,
(25° C, 30 $\mu\text{E}/\text{m}^2\text{sec}$)
Characteristics: Freshwater
O-2
References: 80, 399

Eremosphaera viridis De Bary
380
Oze / Fukushima (1983-08)
Unialgal, Clonal, F. Kasai (1983-09)
Identified by: T. Nakano
Culture conditions: CAM, 20° C, 4 $\mu\text{E}/\text{m}^2\text{sec}$, 3M,
(25° C, 30 $\mu\text{E}/\text{m}^2\text{sec}$)
Characteristics: Freshwater
43-23

643
Miyatoko Mire / Fukushima (1992-04)
Unialgal, Clonal, H. Nozaki (1992-04)
Identified by: H. Nozaki
Culture conditions: AF-6, 20° C, 22 $\mu\text{E}/\text{m}^2\text{sec}$, 2M
Characteristics: Freshwater

- 92-604-E-5
 644
 Miyatoko Mire / Fukushima (1992-04)
 Unialgal, Clonal, H.Nozaki (1992-04)
 Identified by: H.Nozaki
 Culture conditions: AF-6, 20° C, 22 $\mu\text{E}/\text{m}^2\text{sec}$, 2M
 Characteristics: Freshwater
 92-604-E-3
 (25° C, 30 $\mu\text{E}/\text{m}^2\text{sec}$)
 Characteristics: Freshwater, Heterothallic, Crosses
 with NIES-772
 84-15-76
- Errerella bornhemiensis* Conrad
 455
 Between Ghorepani and Billethadi / Nepal
 (1965-12)
 IAM C-341, Axenic, Clonal, T.Ichimura (1972-05)
 Identified by: T.Ichimura
 Culture conditions: C(S), 20° C, 4 $\mu\text{E}/\text{m}^2\text{sec}$, 3M,
 (25° C, 30 $\mu\text{E}/\text{m}^2\text{sec}$)
 Characteristics: Indicator, Freshwater
 N-76-1
 Reference: 80
- Euastrum biverrucosum*
 Gontcharov et M.M.Watanabe
 Syn. *Euastrum englerii* Schmidle
 var. *madagascariense* Bourrelly et Mangium
 840
 Hirosawa-ike Pond / Kyoto (1998-06)
 Unialgal, Clonal, A.Gontcharov (1998-07)
 Identified by: A.Gontcharov
 Culture conditions: CAM, 20° C, 8 $\mu\text{E}/\text{m}^2\text{sec}$, 3M,
 (25° C, 30 $\mu\text{E}/\text{m}^2\text{sec}$)
 Characteristics: Freshwater
 Reference: 48
- Euastrum turgidum* Wallich
 772
 Ishigaki Isl / Okinawa (1984-03)
 Unialgal, Clonal, T.Ichimura (1984-03)
 Identified by: T.Ichimura
 Culture conditions: MG, 20° C, 8 $\mu\text{E}/\text{m}^2\text{sec}$, 3M,
 (25° C, 30 $\mu\text{E}/\text{m}^2\text{sec}$)
 Characteristics: Freshwater, Heterothallic, Crosses
 with NIES-773
 84-15-75
- 773
 Ishigaki Isl / Okinawa (1984-03)
 Unialgal, Clonal, T.Ichimura (1984-03)
 Identified by: T.Ichimura
 Culture conditions: MG, 20° C, 8 $\mu\text{E}/\text{m}^2\text{sec}$, 3M,
 (25° C, 30 $\mu\text{E}/\text{m}^2\text{sec}$)
 Characteristics: Freshwater, Heterothallic, Crosses
 with NIES-772
 84-15-76
- Eudorina cylindrica* Korshikov
 722
 IA / USA
 UTEX 1197, Axenic, Clonal, A.W.Coleman
 (1957-04)
 Culture conditions: AF-6, 20° C, 12 $\mu\text{E}/\text{m}^2\text{sec}$, 1M
 Characteristics: Freshwater, Heterothallic,
rbcL gene (D86833)
 Reference: 244
- Eudorina elegans* Ehrenberg
 351
 Lake Biwa / Shiga (1983-12)
 Axenic, Clonal, S.Suda (1983-12)
 Identified by: S.Suda
 Culture conditions: CA, 20° C, 12 $\mu\text{E}/\text{m}^2\text{sec}$, 1M
 Characteristics: Freshwater, Homothallic
 B-Eud-6
 Reference: 325
- Eudorina elegans* Ehrenberg var. *carteri*
 721
 KY / USA
 UTEX 1212, Axenic, Clonal, P.Cock (1960-04)
 Culture conditions: AF-6, 20° C, 12 $\mu\text{E}/\text{m}^2\text{sec}$, 1M
 Characteristics: Freshwater, Homothallic,
 Monoecious, *rbcL* gene (D88806)
 Reference: 245
- Eudorina elegans* Ehrenberg var. *elegans*
 456
 Chiyoda-ku / Tokyo (1977-09)
 Axenic, Clonal, H.Nozaki (1977-09)
 Identified by: H.Nozaki
 Culture conditions: VT, 20° C, 12 $\mu\text{E}/\text{m}^2\text{sec}$, 1M
 Characteristics: Freshwater, Heterothallic, Male,
 Crosses with NIES-457, *rbcL* gene (D63432)
 A-5 (m)
 References: 223, 242, 243
- 457
 Chiyoda-ku / Tokyo (1977-09)
 Axenic, Clonal, H.Nozaki (1977-09)
 Identified by: H.Nozaki
 Culture conditions: VT, 20° C, 12 $\mu\text{E}/\text{m}^2\text{sec}$, 1M
 Characteristics: Freshwater, Heterothallic, Female,
 Crosses with NIES-456

- I-14 (f)
Reference: 223
- 717
Indiana / USA (1959-06)
UTEX 1193, Axenic, Clonal, M.E.Goldstein
Culture conditions: AF-6, 20° C, 12 µE/m² sec, 1M
Characteristics: Freshwater, Heterothallic,
rbcL gene (D88803)
References 245
- 718
Indiana / USA
UTEX 1195, Axenic, Clonal, A.W.Coleman
Culture conditions: AF-6, 20° C, 12 µE/m² sec, 1M
Characteristics: Freshwater, Heterothallic,
rbcL gene (D88810)
Reference: 245
- 719
Indiana / USA
UTEX 1199, Axenic, Clonal, A.W.Coleman (1956)
Culture conditions: AF-6, 20° C, 12 µE/m² sec, 1M
Characteristics: Freshwater, Heterothallic,
rbcL gene (D88804)
Reference: 245
- 720
Indiana / USA (1959-09)
UTEX 1205, Axenic, Clonal, M.E.Goldstein
Culture conditions: AF-6, 20° C, 12 µE/m² sec, 1M
Characteristics: Freshwater, Heterothallic,
rbcL gene (D88805)
Reference: 245
- Eudorina elegans* Ehrenberg
var. *synoica* Goldstein
458
Midori-ku / Yokohama / Kanagawa (1980-01)
Axenic, Clonal, H.Nozaiki (1980-04)
Identified by: H.Nozaiki
Culture conditions: VT, 20° C, 12 µE/m² sec, 1M
Characteristics: Freshwater, Homothallic,
Monoecious, *rbcL* gene (D88807)
04427-1
References: 229, 245
- 568
Kathmandu / Nepal (1986-09)
Axenic, Clonal, H.Nozaiki (1987-09)
Identified by: H.Nozaiki
Culture conditions: CA, 20° C, 12 µE/m² sec, 1M
Characteristics: Freshwater, Homothallic,
Monoecious, *rbcL* gene (D88808)
7914-E-6
References: 230, 245
- Eudorina illinoisensis* (Kofoid) Pascher
459
Saiwai-ku / Kawasaki / Kanagawa (1984-01)
Axenic, Clonal, H.Nozaiki (1985-06)
Identified by: H.Nozaiki
Culture conditions: VT, 20° C, 12 µE/m² sec, 1M
Characteristics: Freshwater, Heterothallic, Female,
Crosses with NIES-460
5607-E-14 (F)
References: 227, 254
- 460
Saiwai-ku / Kawasaki / Kanagawa (1984-01)
Axenic, Clonal, H.Nozaiki (1985-06)
Identified by: H.Nozaiki
Culture conditions: VT, 20° C, 12 µE/m² sec, 1M
Characteristics: Freshwater, Heterothallic, Male,
Crosses with NIES-459, *rbcL* gene (D63433)
5630-E-3 (m)
References: 227, 242, 243
- 723
Missouri / USA (1956-11)
UTEX 808, Axenic, Clonal, J.Stein
Reidentified by: H.Nozaiki
Culture conditions: AF-6, 20° C, 12 µE/m² sec, 1M
Characteristics: Freshwater, Heterothallic,
rbcL gene (D88809)
Reference: 245
- Eudorina unicocca* G.M.Smith
var. *peripheralis* Goldstein
726
British Columbia / Canada (1961-05)
UTEX 1218, Axenic, Clonal, M.E.Goldstein
Culture conditions: AF-6, 20° C, 12 µE/m² sec, 1M
Characteristics: Freshwater, Heterothallic,
rbcL gene (D86830)
Reference: 244
- Eudorina unicocca* G.M.Smith var. *unicocca*
724
Indiana / USA
UTEX 737, Axenic, Clonal, R.C.Starr
Culture conditions: AF-6, 20° C, 12 µE/m² sec, 1M
Characteristics: Freshwater, Heterothallic,
rbcL gene (D86829)

- Reference: 244
- 725
Ohio / USA (1961-04)
UTEX 1215, Axenic, Clonal, M.E.Goldstein
Culture conditions: AF-6, 20° C, 12 $\mu\text{E}/\text{m}^2\text{sec}$, 1M
Characteristics: Freshwater, Heterothallic,
rbcL gene (D63434)
Reference: 243
- Euglena clara* Skuja
253
Higashiyata River / Ibaraki (1983-07)
Unialgal, Clonal, S.Suda (1983-07)
Identified by: S.Suda
Culture conditions: AF-6, 20° C, 12 $\mu\text{E}/\text{m}^2\text{sec}$, 1M,
(25° C, 30 $\mu\text{E}/\text{m}^2\text{sec}$)
Characteristics: Indicator, Freshwater
USI-21
- Euglena gracilis* Klebs
47
IAM E-3, Axenic, Clonal
Culture conditions: HUT(SS), 20° C, 4 $\mu\text{E}/\text{m}^2\text{sec}$,
1M, (25° C, 30 $\mu\text{E}/\text{m}^2\text{sec}$)
Characteristics: Indicator, Freshwater, Material for
Vitamin B₁₂ bioassay
References: 80, 112, 366
- 48
IAM E-6 (Z strain), Axenic, Clonal
Culture conditions: HUT(SS), 20° C, 4 $\mu\text{E}/\text{m}^2\text{sec}$,
1M, (25° C, 30 $\mu\text{E}/\text{m}^2\text{sec}$)
Characteristics: Freshwater, Material for Vitamin
B₁₂ bioassay
References: 21, 80, 88, 112, 174, 180, 193, 273, 274,
275, 276, 348, 349, 442
- Euglena gracilis* Klebs var. *bacillaris* Pringsheim
49
IAM E-2, Axenic, Clonal
Culture conditions: HUT, 20° C, 4 $\mu\text{E}/\text{m}^2\text{sec}$, 2M,
(25° C, 30 $\mu\text{E}/\text{m}^2\text{sec}$)
Characteristics: Freshwater
References: 80, 112, 366
- Euglena mutabilis* Schmitz
286
Takatori River / Ibaraki (1984-10)
Axenic, Clonal, S.Suda (1984-10)
Identified by: S.Suda
- Culture conditions: AF-6, 20° C, 12 $\mu\text{E}/\text{m}^2\text{sec}$, 1M,
(25° C, 30 $\mu\text{E}/\text{m}^2\text{sec}$)
Characteristics: Indicator, Freshwater
- Eunotia pectinalis* (Kützing) Rabenhorst
var. *minor* (Kützing) Rabenhorst
461
Mt.Tsukuba / Ibaraki (1987-04)
Unialgal, Non-clonal, F.Kasai (1987-05)
Identified by: N.Takamura
Culture conditions: CSI, 15° C, 15 $\mu\text{E}/\text{m}^2\text{sec}$, 4M
Characteristics: Freshwater
(1)-16
Reference: 338
- Eutreptiella gymnastica* Thronsen
381
Yashima Bay / Kagawa (1982-10)
Axenic, Clonal, S.Yoshimatsu
Identified by: S.Yoshimatsu
Culture conditions: f/2, ESM, 20° C, 32 $\mu\text{E}/\text{m}^2\text{sec}$,
1M
Characteristics: Red tide, Marine,
COXI gene (AB000136)
KGW-63-1
Reference: 97
- Fibrocapsa japonica* Toriumi et Takano
136
Tsuda Bay / Kagawa (1978-07)
Axenic, Clonal, K.Yuki
Identified by: K.Yuki
Culture conditions: f/2, 20° C, 32 $\mu\text{E}/\text{m}^2\text{sec}$, 1M
Characteristics: Red tide, Marine, Untransportable
KGW-20-2
Reference: 361
- 462
Hasaki / Ibaraki (1987-05)
Axenic, Clonal, T.Sawaguchi (1987-05)
Identified by: T.Sawaguchi
Culture conditions: ESM, 20° C, 40 $\mu\text{E}/\text{m}^2\text{sec}$, 1M
Characteristics: Red tide, Marine, Untransportable
HASS-8
- 560
Mikawa bay / Aichi
Axenic, Non-clonal, S.Toriumi
Identified by: T.Honjou
Culture conditions: ESM, 20° C, 32 $\mu\text{E}/\text{m}^2\text{sec}$, 1M
Characteristics: Red tide, Marine, Untransportable

- 605
Seto Inland Sea / Yamaguchi (1970-08)
Axenic, Clonal, H.Iwasaki (1970-08)
Identified by: H.Takano
Culture conditions: f/2, 20° C, 40 $\mu\text{E}/\text{m}^2\text{sec}$, 1M
Characteristics: Red tide, Marine, Untransportable
- Fischerella major* Gomont
592
Yukawa-hot spring / Iwate (1990-09)
Unialgal, Clonal, T.Hagiwara (1990-10)
Identified by: T.Hagiwara
Culture conditions: CB, 20° C, 4 $\mu\text{E}/\text{m}^2\text{sec}$, 2M,
(25° C, 30 $\mu\text{E}/\text{m}^2\text{sec}$), [Cryopreserved]
Characteristics: Benthic
Yu-50
- Fragilaria capucina* Desmazières
391
Lake Kasumigaura / Ibaraki (1985-04)
Unialgal, Clonal, T.Sawaguchi (1985-04)
Identified by: M.Idei
Culture conditions: CSi, M Chu No.10, 15° C,
20 $\mu\text{E}/\text{m}^2\text{sec}$, 1M
Characteristics: Freshwater
KEB-24
- Galdieria sulphuraria* (Galdieri) Merola
250
IAM M-8, Unialgal, Non-clonal
Culture conditions: Allen, 20° C, 4 $\mu\text{E}/\text{m}^2\text{sec}$, 4M,
(20° C, 12 $\mu\text{E}/\text{m}^2\text{sec}$)
Characteristics: Hot spring, Formerly identified as
Cyanidium caldarium (Tilden) Geitler
References: 80, 134, 193
- 550
Unialgal, Non-clonal, Pinto
Identified by: A.Merola et al.
Culture conditions: Allen, 20° C, 4 $\mu\text{E}/\text{m}^2\text{sec}$, 6M,
(25° C, 15 $\mu\text{E}/\text{m}^2\text{sec}$)
Characteristics: Acidophilic, Type strain
002
Reference: 175
- Gephyrocapsa oceanica* Kamptner
353
Tsushima / Nagasaki (1986-03)
Axenic, Clonal, T.Sawaguchi (1986-05)
Identified by: I.Inouye
Culture conditions: ESM, 20° C, 12 $\mu\text{E}/\text{m}^2\text{sec}$, 20D
Characteristics: Marine, COXI gene (AB000118)
TMCO-2
References: 65, 160
- 838
Mutsu Bay / Aomori (1990-11)
Unialgal, Clonal, M.Kawachi (1990-11)
Identified by: M.Kawachi
Culture conditions: ESM, 20° C, 12 $\mu\text{E}/\text{m}^2\text{sec}$, 1M
Characteristics: Marine, Red tide
GO-01
Reference: 297
- Glenodiniopsis uliginosa* (Schilling) Woloszynska
463
Shizukuishi / Iwate (1984-09)
Axenic, Clonal, T.Sawaguchi (1984-09)
Identified by: T.Sawaguchi
Culture conditions: AF-6/2, 20° C, 40 $\mu\text{E}/\text{m}^2\text{sec}$,
2M
Characteristics: Freshwater, Unstable,
Untransportable
TM3D-6
- Gloeomonas lateperforata* (Skuja) Ettl
464
Tsukuba / Ibaraki (1982-11)
Axenic, Clonal, F.Kasai (1982-11)
Identified by: S.Suda
Culture conditions: C, 20° C, 22 $\mu\text{E}/\text{m}^2\text{sec}$, 2M
Characteristics: Freshwater
- Gomphonema angustatum* (Kützing) Rabenhorst
var. *obtusatum* (Kützing) Grunow
620
Mt.Tsukuba / Ibaraki (1987-04-17)
Unialgal, Clonal, F.Kasai (1987-05)
Identified by: N.Takamura
Culture conditions: CSi, 15° C, 15 $\mu\text{E}/\text{m}^2\text{sec}$, 2M
Characteristics: Freshwater
1-36
Reference: 338
- Gomphonema gracile* Ehrenberg var. *gracile*
465
Ashio / Gunma (1987-08)
Unialgal, Clonal, F.Kasai (1987-08)
Identified by: N.Takamura
Culture conditions: CSi, 15° C, 15 $\mu\text{E}/\text{m}^2\text{sec}$, 2M
Characteristics: Freshwater
Ast-1-1
Reference: 338

Gomphonema parvulum Kützting var. *parvulum*
466

Shirai River / Sapporo (1987-07)
Unialgal, Non-clonal, F.Kasai (1987-07)
Identified by: N.Takamura
Culture conditions: CSi, 10° C, 15 µE/m²sec, 2M
Characteristics: Freshwater
Tst-1-18
Reference: 338

467

Shirai River / Sapporo (1987-07)
Unialgal, Clonal, F.Kasai (1987-07)
Identified by: N.Takamura
Culture conditions: CSi, 10° C, 15 µE/m²sec, 2M
Characteristics: Freshwater
Tst-4-3
Reference: 338

Gonatozygon brebissonii De Bary

138

Lake Kasumigaura / Ibaraki (1974-11)
Axenic, Clonal
Culture conditions: C, 20° C, 8 µE/m²sec, 4M,
(20° C, 12 µE/m²sec)
Characteristics: Freshwater
KAS-4-43

139

Lake Shoji / Yamanashi (1981-10)
TAC 56-1, Axenic, Clonal, M.Watanabe
Identified by: M.Watanabe
Culture conditions: C, 20° C, 8 µE/m²sec, 4M,
(20° C, 12 µE/m²sec)
Characteristics: Freshwater
TAN-56-1

Gonatozygon monotaenium De Bary

247

Tsukiyono / Gunma (1984-06)
Axenic, Clonal, F.Kasai (1984-06)
Identified by: F.Kasai
Culture conditions: C, 20° C, 8 µE/m²sec, 3M,
(20° C, 12 µE/m²sec)
Characteristics: Freshwater, Homothallic
84-25-109

287

Lake Yamanaka / Yamanashi (1981-10)
TAC 53-3, Unialgal, Clonal, M.Watanabe
Identified by: M.Watanabe

Culture conditions: MG, 20° C, 8 µE/m²sec, 2M,
(20° C, 12 µE/m²sec)

Characteristics: Freshwater
TAN-53-3

Gonium multicocum Pocock

737

UTEX 2580, Axenic, Clonal, H.Nozaki
Identified by: H.Nozaki
Culture conditions: AF-6, 20° C, 12 µE/m²sec, 1M
Characteristics: Freshwater, F1 clone of UTEX 2579,
rbcL gene (D63435)
90-530-F1-5
References: 243, 252

Gonium pectorale Müller var. *pectorale*

468

Kohoku-ku / Yokohama / Kanagawa (1979-04)
Axenic, Clonal, H.Nozaki (1979-04)
Identified by: H.Nozaki
Culture conditions: VT, 20° C, 12 µE/m²sec, 1M
Characteristics: Freshwater, Heterothallic,
Mating type -, Crosses with NIES-469
9406-10
References: 135, 224, 234, 249

469

Kohoku-ku / Yokohama / Kanagawa (1979-04)
Axenic, Clonal, H.Nozaki (1979-04)
Identified by: H.Nozaki
Culture conditions: VT, 20° C, 12 µE/m²sec, 1M
Characteristics: Freshwater, Heterothallic,
Mating type +, Crosses with NIES-468
9406-12
Reference: 224

569

Kourakuen / Okayama (1988-10)
Unialgal, Clonal, H.Nozaki
Identified by: H.Nozaki
Culture conditions: VTAC, 20° C, 22 µE/m²sec, 1M
Characteristics: Freshwater, Heterothallic, Isogamy,
Mating type +, Crosses with NIES-570,
rbcL gene (D63437)
88-1113-G-1
Reference: 243

570

Kourakuen / Okayama (1988-10)
Unialgal, Clonal, H.Nozaki
Identified by: H.Nozaki

- Culture conditions: VTAC, 20° C, 22 μE/m² sec, 1M
 Characteristics: Freshwater, Heterothallic, Isogamy,
 Mating type –, Crosses with NIES-569
 88-1113-G-2
- 645
 Near Goshokake Hot Spring / Akita (1985-07)
 Unialgal, Clonal, H.Nozaki (1985-09)
 Identified by: H.Nozaki
 Culture conditions: AF-6, 20° C, 22 μE/m² sec, 1M
 Characteristics: Freshwater, Heterothallic, Isogamy,
 Mating type +
 5912-6(+)
- 646
 Near Goshokake Hot Spring / Akita (1985-07)
 Unialgal, Clonal, H.Nozaki (1985-09)
 Identified by: H.Nozaki
 Culture conditions: AF-6, 20° C, 22 μE/m² sec, 1M
 Characteristics: Freshwater, Heterothallic, Isogamy,
 Mating type –
 5912-6(-)
- Gonium quadratum* Pringsheim ex Nozaki
- 647
 Unialgal, Clonal, H.Nozaki (1990-08)
 Identified by: H.Nozaki
 Culture conditions: AF-6, 20° C, 22 μE/m² sec, 1M
 Characteristics: Freshwater, Heterothallic, Isogamy,
 Mating type –, F1 clone of NIES-652 × NIES-653,
 Sister clone to NIES-648, 649, and 650 from one
 zygote
 90-809-F1-2-1
- 648
 Unialgal, Clonal, H.Nozaki (1990-08)
 Identified by: H.Nozaki
 Culture conditions: AF-6, 20° C, 22 μE/m² sec, 1M
 Characteristics: Freshwater, Heterothallic, Isogamy,
 Mating type +, F1 clone of NIES-652 × NIES-653,
 Sister clone to NIES-647, 649, and 650 from one
 zygote
 90-809-F1-2-2
- 649
 Unialgal, Clonal, H.Nozaki (1990-08)
 Identified by: H.Nozaki
 Culture conditions: AF-6, 20° C, 22 μE/m² sec, 1M
 Characteristics: Freshwater, Heterothallic, Isogamy,
 Mating type +, F1 clone of NIES-652 × NIES-653,
 Sister clone to NIES-647, 648, and 650 from one
 zygote
 90-809-F1-2-3
- 650
 Unialgal, Clonal, H.Nozaki (1990-08)
 Identified by: H.Nozaki
 Culture conditions: AF-6, 20° C, 22 μE/m² sec, 1M
 Characteristics: Freshwater, Heterothallic, Isogamy,
 Mating type –, F1 clone of NIES-652 × NIES-653,
 Sister clone to NIES-647, 648, and 649 from one
 zygote
 90-809-F1-2-4
- 651
 Klausen / Italy
 UTEX 956, Unialgal, Clonal, E.G.Pringsheim
 (1957)
 Identified by: E.G.Pringsheim
 Culture conditions: AF-6, 20° C, 22 μE/m² sec, 1M
 Characteristics: Freshwater, Type strain
 References: 234, 235
- 652
 Itahari / Nepal (1989-10)
 Unialgal, Clonal, H.Nozaki (1990-04)
 Identified by: H.Nozaki
 Culture conditions: AF-6, 20° C, 22 μE/m² sec, 1M
 Characteristics: Freshwater, Heterothallic, Isogamy,
 Mating type –, Crosses with NIES-653
 90-423-3
 Reference: 235
- 653
 Itahari / Nepal (1989-10)
 Unialgal, Clonal, H.Nozaki (1990-04)
 Identified by: H.Nozaki
 Culture conditions: AF-6, 20° C, 22 μE/m² sec, 1M
 Characteristics: Freshwater, Heterothallic, Isogamy,
 Mating type +, Crosses with NIES-652,
rbcL gene (D63438)
 90-423-2
 References: 235, 243, 249
- * *Gonium sacculiferum* Scherffel
 See *Basichlamys sacculifera* (Scherffel) Skuja
- * *Gonium sociale* (Dujardin) Warming var. *sociale*
 See *Tetrabaena socialis* (Dujardin) Nozaki et Ito
 var. *socialis*

Gonium viridistellatum M.Watanabe

288

Okinawa / Okinawa (1973-06)

Axenic, Clonal, M.Watanabe

Identified by: M.Watanabe

Culture conditions: CA, 20° C, 12 µE/m² sec, 1M

Characteristics: Indicator, Freshwater, Heterothallic,

Mating type -, Crosses with NIES-289 and 290

G4

References: 232, 375

289

Okinawa / Okinawa (1973-06)

Axenic, Clonal, M.Watanabe

Identified by: M.Watanabe

Culture conditions: CA, 20° C, 12 µE/m² sec, 1M

Characteristics: Indicator, Freshwater, Type strain,

Heterothallic, Mating type +, Crosses with

NIES-288

G3

References: 232, 375

290

Okinawa / Okinawa (1973-06)

Axenic, Clonal, M.Watanabe

Identified by: M.Watanabe

Culture conditions: CA, 20° C, 12 µE/m² sec, 1M

Characteristics: Indicator, Freshwater, Type strain,

Heterothallic, Mating type +, Crosses with

NIES-288

G1

References: 232, 375

654

Midori-ku / Yokohama (1980-01)

UTEX 2519, Unialgal, Clonal, H.Nozaki (1985-11)

Identified by: H.Nozaki

Culture conditions: VT, 20° C, 22 µE/m² sec, 1M

Characteristics: Freshwater, Heterothallic, Isogamy,

Mating type +, *rbcl* gene (D86831)

KY-4 (+)

References: 232, 242, 244

655

Midori-ku / Yokohama (1980-01)

UTEX 2520, Unialgal, Clonal, H.Nozaki (1985-11)

Identified by: H.Nozaki

Culture conditions: VT, 20° C, 22 µE/m² sec, 1M

Characteristics: Freshwater, Heterothallic, Isogamy,

Mating type -

KY-7 (-)

Reference: 232

Graesiella emersonii (Shihira et Kraus) Nozaki et al.

Syn. *Chlorella emersonii* Shihira et Krauss

Chlorella fusca Shihira et Krauss var. *vacuolata*

Shihira et Krauss

226

IAM C-28, Axenic, Clonal, E.G.Pringsheim

Culture conditions: C(S), 20° C, 4 µE/m² sec, 3M,

(25° C, 30 µE/m² sec)

Characteristics: Freshwater, Formerly identified as

Chlorella pyrenoidosa Chick

References: 80, 176, 214, 248, 340, 366, 417, 438,

439, 440, 441

687

USA

IAM C-104, CCAP 211/8B, Unialgal, Clonal,

R.Emerson (1923)

Culture conditions: C(S), 20° C, 4 µE/m² sec, 3M,

(25° C, 30 µE/m² sec)

Characteristics: Freshwater, Type strain of

Chlorella fusca Shihira et Krauss var. *vacuolata*

Shihira et Krauss

Reference: 248

688

CCAP 211/8G, Unialgal, Clonal, R.Emerson

Culture conditions: C(S), 20° C, 4 µE/m² sec, 3M,

(25° C, 30 µE/m² sec)

Characteristics: Freshwater

Reference: 248

689

CCAP 211/8H, Unialgal, Clonal, R.Emerson

Culture conditions: C(S), 20° C, 4 µE/m² sec, 3M,

(25° C, 30 µE/m² sec)

Characteristics: Freshwater

Reference: 248

690

CCAP 211/11N, Unialgal, Clonal, R.Emerson

(1939)

Culture conditions: C(S), 20° C, 4 µE/m² sec, 3M,

(25° C, 30 µE/m² sec)

Characteristics: Freshwater, Type strain of

Chlorella emersonii Shihira et Krauss

Reference: 248

Gymnodinium fuscum Stein

470

Tsuchiura / Ibaraki (1986-02)

Unialgal, Clonal, T.Sawaguchi (1986-05)

Identified by: T.Sawaguchi
Culture conditions: AF-6/2, 20° C, 40 µE/m² sec, 1M
Characteristics: Freshwater, Unstable,
Untransportable
SPSDG

Gymnodinium mikimotoi Miyake et Kominami
ex Oda

Syn. *Gymnodinium nagasakiense*
Takayama et Adachi

680

Uchiumi Bay / Kagawa (1992-10)
Unialgal, Clonal, S.Yoshimatsu (1992-10)
Identified by: S.Yoshimatsu
Culture conditions: ESM, 20° C, 40 µE/m² sec, 1M
Characteristics: Red tide, Marine, Untransportable

* *Gymnodinium nagasakiense* Takayama et Adachi
See *Gymnodinium mikimotoi*
Miyake et Mominami ex Oda

Gyrodinium instriatum Freudenthal et Lee
143

Shodo Isl. / Kagawa (1978-06)
Unialgal, Clonal, K.Yuki
Identified by: K.Yuki
Culture conditions: f/2, ESM, 20° C, 32 µE/m² sec,
1M
Characteristics: Red tide, Marine Unstable,
Untransportable
KGW-17-1

Haematococcus lacustris

(Girod-Chantrans) Rostafinski

Syn. *Haematococcus pluvialis* Flotow
144

Sapporo / Hokkaido (1964-07)
IAM C-392, Axenic, Clonal, T.Ichimura (1964-07)
Identified by: T.Ichimura
Culture conditions: C(S), 20° C, 4 µE/m² sec, 3M,
(25° C, 30 µE/m² sec)
Characteristics: Freshwater, Homothallic, Isogamy
MKF-8
References: 80, 114, 115, 142, 143, 144, 145, 146,
147, 148, 149, 151, 150, 152, 344, 343, 345, 399

* *Haematococcus pluvialis* Flotow
See *Haematococcus lacustris*
(Girod-Chantrans) Rostafinski

Hafniomonas montana (Geitler) Ettl et Moestrup
257

Tsukuba / Ibaraki (1983-10)
Axenic, Clonal, S.Suda (1983-10)
Identified by: I.Inouye
Culture conditions: C, 20° C, 12 µE/m² sec, 1M,
(20° C, 25 µE/m² sec)
Characteristics: Freshwater
OUT-5
References: 327, 399

656

Tsukuba / Ibaraki (1986-04-30)
Axenic, Clonal, S.Suda (1986-05)
Identified by: S.Suda
Culture conditions: C, 20° C, 22 µE/m² sec, 1M
Characteristics: Freshwater
430M3-3

Hantzschia amphioxys (Ehrenberg) Grunow
var. *compacta* Hustedt
587

Tsukuba / Ibaraki (1990-04)
Unialgal, Clonal, T.Hagiwara (1990-04)
Identified by: T.Hagiwara
Culture conditions: CSi, 15° C, 35 µE/m² sec, 1M
Characteristics: Freshwater
Wn-24

Haramonas dimorpha Horiguchi
716

Daintree River / Australia (1991-09)
Unialgal, Clonal, T.Horiguchi (1991-10)
Identified by: T.Horiguchi
Culture conditions: f/2, ESM, 20° C, 40 µE/m² sec,
1M
Characteristics: Brackish, Type strain,
Untransportable
Reference: 72

Hemidinium nasutum Stein
471

Tsuchiura / Ibaraki (1987-08)
Unialgal, Clonal, T.Sawaguchi (1987-08)
Identified by: T.Sawaguchi
Culture conditions: AF-6/2, 20° C, 40 µE/m² sec,
1M
Characteristics: Freshwater, Untransportable
87SPD-1

Heterocapsa pygmaea Loeblich III et al.
472

- Kashiwazaki / Niigata (1986-08)
 Unialgal, Clonal, T.Sawaguchi (1986-08)
 Identified by: T.Sawaguchi
 Culture conditions: ESM, 20° C, 40 $\mu\text{E}/\text{m}^2\text{sec}$, 1M
 Characteristics: Marine, Untransportable
 KSTH-23
- 473
 Izuhara / Nagasaki (1986-03)
 Unialgal, Clonal, T.Sawaguchi (1986-03)
 Identified by: T.Sawaguchi
 Culture conditions: ESM, 20° C, 40 $\mu\text{E}/\text{m}^2\text{sec}$, 1M
 Characteristics: Marine, Untransportable
 TMUD-2
- Heterocapsa triquetra* Stein
 7
 Osaka Bay / Osaka (1981-04)
 Axenic, Clonal, S.Yamochi
 Identified by: S.Yamochi
 Culture conditions: f/2, ESM, 20° C, 32 $\mu\text{E}/\text{m}^2\text{sec}$, 1M
 Characteristics: Red tide, Marine, Untransportable
 OHet
 Reference: 160
- 235
 Harima-Nada / Seto Inland Sea (1982-03)
 Axenic, Clonal, S.Yoshimatsu
 Identified by: S.Yoshimatsu
 Culture conditions: f/2, 20° C, 32 $\mu\text{E}/\text{m}^2\text{sec}$, 1M
 Characteristics: Red tide, Marine, Untransportable
 KGW-57
 Reference: 361
- Heterosigma akashiwo* (Hada) Hada
 4
 Fukuyama Bay / Hiroshima (1966-06)
 Axenic, Clonal, H.Iwasaki et al.
 Identified by: H.Iwasaki et al.
 Culture conditions: f/2, M-ASP7, 20° C, 32 $\mu\text{E}/\text{m}^2\text{sec}$, 1M
 Characteristics: Red tide, Marine, Untransportable
 FHE
 References: 17, 58, 110, 300
- 5
 Gokasho Bay / Mie (1966)
 Axenic, Clonal, H.Iwasaki et al.
 Identified by: Y.Hara
 Culture conditions: f/2, M-ASP7, 20° C, 32 $\mu\text{E}/\text{m}^2\text{sec}$, 1M
- Characteristics: Red tide, Marine, Untransportable
 GHE
 References: 111, 288
- 6
 Osaka Bay / Osaka (1979-08)
 Axenic, Clonal, M.M.Watanabe
 Identified by: M.M.Watanabe
 Culture conditions: f/2, M-ASP7, 20° C, 32 $\mu\text{E}/\text{m}^2\text{sec}$, 1M
 Characteristics: Red tide, Marine, Untransportable
 OHE-1
 References: 57, 60, 153, 154, 155, 157, 160, 161, 178, 179, 181, 194, 296, 323, 324, 331, 362, 363, 364, 378, 379, 380, 381, 382, 383, 384, 385, 401, 403, 404, 405, 406, 407, 414, 433, 435
- 9
 Harima-Nada / Seto Inland Sea (1983-02)
 Axenic, Clonal, M.M.Watanabe (1983-05)
 Identified by: M.M.Watanabe
 Culture conditions: f/2, M-ASP7, 20° C, 32 $\mu\text{E}/\text{m}^2\text{sec}$, 1M
 Characteristics: Red tide, Marine, Untransportable
 H-28
- 10
 Harima-Nada / Seto Inland Sea (1983-02)
 Axenic, Clonal, M.M.Watanabe (1983-05)
 Identified by: M.M.Watanabe
 Culture conditions: f/2, M-ASP7, 20° C, 32 $\mu\text{E}/\text{m}^2\text{sec}$, 1M
 Characteristics: Red tide, Marine, Untransportable
 H-40
- 145
 Nomaike / Kagoshima (1978-05)
 Axenic, Clonal, S.Yoshimatsu
 Identified by: S.Yoshimatsu
 Culture conditions: f/2, M-ASP7, 20° C, 32 $\mu\text{E}/\text{m}^2\text{sec}$, 1M
 Characteristics: Red tide, Marine, Untransportable
 KGW-11-5
 Reference: 361
- 146
 Shido Bay / Kagawa (1978-06)
 Axenic, Clonal, K.Yuki
 Identified by: K.Yuki
 Culture conditions: f/2, M-ASP7, 20° C, 32 $\mu\text{E}/\text{m}^2\text{sec}$, 1M
 Characteristics: Red tide, Marine, Untransportable

- KGW-21-2
293
Onagawa Bay / Miyagi (1984-08)
Axenic, Clonal, S.Suda (1984-09)
Identified by: S.Suda
Culture conditions: f/2, 20° C, 32 $\mu\text{E}/\text{m}^2\text{ sec}$, 1M
Characteristics: Red tide, Marine, Untransportable
8280G21-1
Reference: 17
- 561
Mikawa Bay / Aichi
Axenic, Clonal, S.Toriumi
Culture conditions: f/2, 20° C, 32 $\mu\text{E}/\text{m}^2\text{ sec}$, 1M
Characteristics: Red tide, Marine, Untransportable
- Hyalotheca dissiliens* Brébisson ex Ralfs
147
Nagatoro / Saitama (1969-11)
IAM C-510, Unialgal, Clonal, T.Ichimura (1972-06)
Identified by: T.Ichimura
Culture conditions: C, 20° C, 8 $\mu\text{E}/\text{m}^2\text{ sec}$, 3M,
(20° C, 12 $\mu\text{E}/\text{m}^2\text{ sec}$)
Characteristics: Freshwater, Heterothallic,
Crosses with NIES-148
S-9-18
- 148
Nagatoro / Saitama (1969-11)
IAM C-511, Axenic, Clonal, T.Ichimura (1972-06)
Identified by: T.Ichimura
Culture conditions: C, 20° C, 8 $\mu\text{E}/\text{m}^2\text{ sec}$, 3M,
(20° C, 12 $\mu\text{E}/\text{m}^2\text{ sec}$)
Characteristics: Freshwater, Heterothallic,
Crosses with NIES-147
S-9-22
- 149
Lake Kasumigaura / Ibaraki (1975-12)
IAM C-512, Axenic, Clonal, T.Ichimura (1975-12)
Identified by: T.Ichimura
Culture conditions: VT, 20° C, 12 $\mu\text{E}/\text{m}^2\text{ sec}$, 1M
Characteristics: Freshwater, Heterothallic,
Crosses with NIES-150
KAS-7-3
- 150
Lake Kasumigaura / Ibaraki (1975-12)
IAM C-513, Axenic, Clonal, T.Ichimura (1975-12)
Identified by: T.Ichimura
- Culture conditions: VT, 20° C, 12 $\mu\text{E}/\text{m}^2\text{ sec}$, 1M
Characteristics: Freshwater, Heterothallic,
Crosses with NIES-149
KAS-7-8
- Hyalotheca dissiliens* Brébisson ex Ralfs
var. *dissiliens* f. *tridentula* (Nordstedt) Bold
294
Tsukuba / Ibaraki (1982)
Unialgal, Clonal, F.Kasai (1983-02)
Identified by: F.Kasai
Culture conditions: VT, 20° C, 8 $\mu\text{E}/\text{m}^2\text{ sec}$, 3M,
(20° C, 12 $\mu\text{E}/\text{m}^2\text{ sec}$)
Characteristics: Freshwater, Homothallic
H-1
- Hydrococcus rivularis* Kützing
593
Yukawa-hot spring / Iwate (1990-09)
Unialgal, Clonal, T.Hagiwara (1990-10)
Identified by: T.Hagiwara
Culture conditions: CB, 20° C, 4 $\mu\text{E}/\text{m}^2\text{ sec}$, 4M,
(25° C, 30 $\mu\text{E}/\text{m}^2\text{ sec}$), [Cryopreserved]
Characteristics: Benthic
Yu-52
- Hydrodictyon reticulatum* (Lagerheim) Lagerheim
295
Kitakawachi-gun / Osaka (1968-11)
IAM C-335, Unialgal, Clonal, T.Ichimura (1969-01)
Identified by: T.Ichimura
Culture conditions: C(S), 20° C, 4 $\mu\text{E}/\text{m}^2\text{ sec}$, 3M,
(25° C, 30 $\mu\text{E}/\text{m}^2\text{ sec}$)
Characteristics: Freshwater, Homothallic
O-2
Reference: 80
- Katodinium rotundatum* (Lohmann) Loeblich III
356
Hachinohe Harbor / Aomori (1985-01)
Axenic, Clonal, T.Sawaguchi (1985-01)
Identified by: T.Sawaguchi
Culture conditions: f/2, ESM, 5° C, 6 $\mu\text{E}/\text{m}^2\text{ sec}$,
1M (10° C, 15 $\mu\text{E}/\text{m}^2\text{ sec}$)
Characteristics: Marine, Unstable, Untransportable
HHD-1
- Lagerheimia ciliata* (Lagerheim) Chodat
382
Lake Kasumigaura / Ibaraki (1983-08)
Axenic, Clonal, F.Kasai (1983-08)

- Identified by: Y.Niiyama
 Culture conditions: C, 20° C, 4 $\mu\text{E}/\text{m}^2\text{sec}$, 3M,
 (25° C, 30 $\mu\text{E}/\text{m}^2\text{sec}$)
 Characteristics: Freshwater
 F37-1
- Lithodesmium variable* Takano
 588
 Hitachi / Ibaraki (1990-09)
 Unialgal, Non-clonal, S.Ono (1990-10)
 Identified by: S.Ono
 Culture conditions: f/2, 15° C, 10 $\mu\text{E}/\text{m}^2\text{sec}$, 1M
 Characteristics: Red tide, Marine
 St-12
- Lobomonas monstrosa* Korshikov
 474
 Iwaki / Fukushima (1984-08)
 Axenic, Clonal, S.Suda (1984-08)
 Identified by: S.Suda
 Culture conditions: C, 20° C, 22 $\mu\text{E}/\text{m}^2\text{sec}$, 2M
 Characteristics: Freshwater
 FL
- Melosira granulata* (Ehrenberg) Ralfs
 var. *angustissima* Müller f. *spiralis* Müller
 333
 Lake Kasumigaura / Ibaraki (1983-05)
 Axenic, Clonal, T.Hiwatari (1983-05)
 Identified by: M.Mizuno
 Culture conditions: CSi, 15° C, 10 $\mu\text{E}/\text{m}^2\text{sec}$, 1M,
 (20° C, 25 $\mu\text{E}/\text{m}^2\text{sec}$)
 Characteristics: Indicator, Freshwater, Unstable
 K-Melo
 Reference: 325
- Merismopedia tenuissima* Lemmermann
 230
 Tsukuba / Ibaraki (1984-05)
 Unialgal, Clonal, F.Kasai (1984-05)
 Identified by: M.M.Watanabe
 Culture conditions: C, 20° C, 12 $\mu\text{E}/\text{m}^2\text{sec}$, 1M,
 [Cryopreserved]
 Characteristics: Freshwater
 F98-2
 Reference: 126
- Mesostigma viride* Lauterborn
 296
 Mitsukaido / Ibaraki (1985-07)
 Axenic, Clonal, S.Suda (1985-07)
- Identified by: I.Inouye
 Culture conditions: C, 20° C, 40 $\mu\text{E}/\text{m}^2\text{sec}$, 1M
 Characteristics: Freshwater
 KY-14
- 475
 Mitsukaido / Ibaraki (1986-01)
 Axenic, Clonal, S.Suda (1987-12)
 Identified by: S.Suda
 Culture conditions: C, 20° C, 22 $\mu\text{E}/\text{m}^2\text{sec}$, 20D
 Characteristics: Freshwater, Heterothallic,
 Mating type +
 KY-Mes-2
- 476
 Mitsukaido / Ibaraki (1986-01)
 Axenic, Clonal, S.Suda (1986-12)
 Identified by: S.Suda
 Culture conditions: C, 20° C, 22 $\mu\text{E}/\text{m}^2\text{sec}$, 20D
 Characteristics: Freshwater, Heterothallic,
 Mating type –
 KY-Mes-1
- 477
 Mitsukaido / Ibaraki (1986-01)
 Axenic, Clonal, S.Suda (1986-12)
 Identified by: S.Suda
 Culture conditions: AF-6, 20° C, 22 $\mu\text{E}/\text{m}^2\text{sec}$, 20D
 Characteristics: Freshwater, Heterothallic,
 Mating type –
 KY-Mes-3
- Mesotaenium kramstae* Lemmermann
 657
 IAM C-330, Unialgal, Clonal
 Culture conditions: C, 20° C, 4 $\mu\text{E}/\text{m}^2\text{sec}$, 3M,
 (25° C, 30 $\mu\text{E}/\text{m}^2\text{sec}$)
 Characteristics: Freshwater, Heterothallic, Crosses
 with NIES-658
- 658
 IAM C-331, Unialgal, Clonal
 Culture conditions: C, 20° C, 4 $\mu\text{E}/\text{m}^2\text{sec}$, 3M,
 (25° C, 30 $\mu\text{E}/\text{m}^2\text{sec}$)
 Characteristics: Freshwater, Heterothallic, Crosses
 with NIES-657
- Micractinium pusillum* Fresenius
 151
 Lake Kasumigaura / Ibaraki (1983-07)
 Axenic, Clonal, F.Kasai (1983-07)

- Identified by: F.Kasai
 Culture conditions: C(S), 20° C, 4 µE/m² sec, 3M,
 (25° C, 30 µE/m² sec)
 Characteristics: Indicator, Freshwater
 F-19-4
 References: 126, 399
- Micrasterias anomala* Turner
 774
 Cairns, Queensland / Australia (1988-09)
 Unialgal, Clonal, T.Ichimura (1988)
 Identified by: T.Ichimura
 Culture conditions: MG, 20° C, 8 µE/m² sec, 3M,
 (25° C, 30 µE/m² sec)
 Characteristics: Freshwater, Homothallic
 88-95-12
- 775
 Near Melaka / Malaysia (1985-08)
 Unialgal, Clonal, T.Ichimura (1985)
 Identified by: T.Ichimura
 Culture conditions: MG, 20° C, 8 µE/m² sec, 3M,
 (25° C, 30 µE/m² sec)
 Characteristics: Freshwater, Heterothallic, Crosses
 with NIES-776
 85-30-30
- 776
 Near Melaka / Malaysia (1985-08)
 Unialgal, Clonal, T.Ichimura (1985)
 Identified by: T.Ichimura
 Culture conditions: MG, 20° C, 8 µE/m² sec, 3M,
 (25° C, 30 µE/m² sec)
 Characteristics: Freshwater, Heterothallic, Crosses
 with NIES-775
 85-30-38
- Micrasterias crux-melitensis* Ralfs
 152
 Kathmandu / Nepal (1968-05)
 IAM C-427, Unialgal, Clonal, T.Ichimura (1970-12)
 Identified by: T.Ichimura
 Culture conditions: VT, 20° C, 8 µE/m² sec, 3M,
 (25° C, 30 µE/m² sec)
 Characteristics: Freshwater, Homothallic
 N-90-27
 Reference: 80
- Micrasterias foliacea* Bailey ex Ralfs
 777
 2 km southeast of Melaka / Malaysia (1985-08)
- Unialgal, Clonal, T.Ichimura (1985)
 Identified by: T.Ichimura
 Culture conditions: MG, 20° C, 8 µE/m² sec, 3M,
 (25° C, 30 µE/m² sec)
 Characteristics: Freshwater, Heterothallic, Crosses
 with NIES-778
 M2-1
- 778
 2 km southeast of Melaka / Malaysia (1985-08)
 Unialgal, Clonal, T.Ichimura (1985)
 Identified by: T.Ichimura
 Culture conditions: MG, 20° C, 8 µE/m² sec, 3M,
 (25° C, 30 µE/m² sec)
 Characteristics: Freshwater, Heterothallic, Crosses
 with NIES-777
 M2-2
- Micrasterias foliacea* Bailey ex Ralfs var. *foliacea*
 297
 Higashihiroshima / Hiroshima (1983-10)
 Unialgal, Clonal, F.Kasai (1983-10)
 Identified by: F.Kasai
 Culture conditions: MG, 20° C, 8 µE/m² sec, 3M,
 (25° C, 15 µE/m² sec)
 Characteristics: Freshwater
 83-24-24
- Micrasterias mahabuleshwariensis* Hobson
 779
 2 km southeast of Melaka / Malaysia (1985-08)
 Unialgal, Clonal, T.Ichimura (1985)
 Identified by: T.Ichimura
 Culture conditions: MG, 20° C, 8 µE/m² sec, 3M,
 (25° C, 30 µE/m² sec)
 Characteristics: Freshwater, Heterothallic, Crosses
 with NIES-780
 M2-6
- 780 (PS-952)
 2 km southeast of Melaka / Malaysia (1985-08)
 Unialgal, Clonal, T.Ichimura (1985)
 Identified by: T.Ichimura
 Culture conditions: MG, 20° C, 8 µE/m² sec, 3M,
 (25° C, 30 µE/m² sec)
 Characteristics: Freshwater, Heterothallic, Crosses
 with NIES-779
 M2-7
- Micrasterias thomasiana* Archer
 var. *notata* (Nordstedt) Grönblad

- 781
2 km southeast of Melaka / Malaysia (1985-08)
Unialgal, Clonal, T.Ichimura (1985)
Identified by: T.Ichimura
Culture conditions: MG, 20° C, 8 µE/m² sec, 3M,
(25° C, 30 µE/m² sec)
Characteristics: Freshwater, Heterothallic, Crosses
with NIES-782
85-28-14
- 782
2 km southeast of Melaka / Malaysia (1985-08)
Unialgal, Clonal, T.Ichimura (1985)
Identified by: T.Ichimura
Culture conditions: MG, 20° C, 8 µE/m² sec, 3M,
(25° C, 30 µE/m² sec)
Characteristics: Freshwater, Heterothallic, Crosses
with NIES-781
85-28-57
- Micrasterias truncata* (Corda) Brébisson ex Ralfs
var. *pusilla* G.S.West
- 783
Centennial Park, Sydney / Australia (1988-09)
Unialgal, Clonal, F.Kasai (1988-09)
Identified by: F.Kasai
Culture conditions: MG, 20° C, 8 µE/m² sec, 3M,
(25° C, 30 µE/m² sec)
Characteristics: Freshwater, Heterothallic, Crosses
with NIES-784
88-7-2
- 784
Near Cairns, Queensland / Australia (1988-09)
Unialgal, Clonal, F.Kasai (1988-09)
Identified by: F.Kasai
Culture conditions: MG, 20° C, 8 µE/m² sec, 3M,
(25° C, 30 µE/m² sec)
Characteristics: Freshwater, Heterothallic, Crosses
with NIES-783
88-8-5
- Microcystis aeruginosa* (Kützing) Lemmermann
f. *aeruginosa*
- 44
Lake Kasumigaura / Ibaraki (1974-08)
IAM M-176, Axenic, Clonal, M.M.Watanabe
(1974-08).
Identified by: M.M.Watanabe
Culture conditions: CB, 25° C, 24 µE/m² sec, 20D,
[Cryopreserved]
- Characteristics: Water bloom, Indicator, Freshwater
References: 5, 52, 80, 87, 118, 119, 126, 169, 216,
325, 341, 399, 410, 431
- 87
Lake Kasumigaura / Ibaraki (1982-09)
Axenic, Clonal, M.H.Watanabe (1982-09)
Identified by: M.H.Watanabe
Culture conditions: MA, 25° C, 24 µE/m² sec, 20D,
[Cryopreserved]
Characteristics: Water bloom, Indicator, Freshwater
K-MA-11
References: 169, 216, 271, 347, 399
- 88
Lake Kawaguchi / Yamanashi (1981-06)
Unialgal, Clonal, M.H.Watanabe (1981-06)
Identified by: M.H.Watanabe
Culture conditions: MA, 25° C, 24 µE/m² sec, 20D,
[Cryopreserved]
Characteristics: Water bloom, Indicator, Freshwater
KW-MA1-3
References: 100, 102, 132, 325, 399
- 89
Lake Kawaguchi / Yamanashi (1981-06)
Unialgal, Clonal, M.H.Watanabe (1981-06)
Identified by: M.H.Watanabe
Culture conditions: MA, 25° C, 24 µE/m² sec, 20D,
[Cryopreserved]
Characteristics: Water bloom, Indicator, Freshwater
KW-MA2-5
References: 169, 215, 216, 399, 411
- 90
Lake Kawaguchi / Yamanashi (1981-06)
Axenic, Clonal, M.H.Watanabe (1981-06)
Identified by: M.H.Watanabe
Culture conditions: MA, 25° C, 24 µE/m² sec, 20D,
[Cryopreserved]
Characteristics: Water bloom, Indicator, Freshwater
KW-MB-2
References: 101, 105, 399, 410
- 91
Lake Kasumigaura / Ibaraki (1982-09)
Unialgal, Clonal, M.H.Watanabe (1982-09)
Identified by: M.H.Watanabe
Culture conditions: MA, 25° C, 24 µE/m² sec, 20D,
[Cryopreserved]
Characteristics: Water bloom, Indicator, Freshwater
K-MB-13

- Reference: 399
- 99
Lake Suwa / Nagano (1982-08)
Unialgal, Clonal, M.H.Watanabe (1982-08)
Identified by: M.H.Watanabe
Culture conditions: MA, 25° C, 24 µE/m² sec, 20D,
[Cryopreserved]
Characteristics: Water bloom, Indicator, Freshwater
S-MA-S5
References: 399, 431
- 100
Lake Suwa / Nagano (1982-08)
Unialgal, Clonal, M.H.Watanabe (1982-08)
Identified by: M.H.Watanabe
Culture conditions: MA, 25° C, 24 µE/m² sec, 20D,
[Cryopreserved]
Characteristics: Water bloom, Indicator, Freshwater
S-MB-S7
References: 266, 268, 287, 386, 399, 425
- 101
Lake Suwa / Nagano (1982-10)
TAC 48, Unialgal, Clonal, M.Watanabe (1982-10)
Identified by: M.Watanabe
Culture conditions: CB, 25° C, 24 µE/m² sec, 20D,
[Cryopreserved]
Characteristics: Water bloom, Indicator, Freshwater
S-TAN-48
References: 198, 386, 399
- 298
Lake Kasumigaura / Ibaraki (1982-09)
TAC 47, Axenic, Clonal, M.Watanabe (1982-09)
Culture conditions: CB, 25° C, 24 µE/m² sec, 20D,
[Cryopreserved]
Characteristics: Water bloom, Toxic, Freshwater
K-TAN-47
References: 52, 126, 166, 167, 169, 187, 216, 267,
287, 386, 411
- 299
Lake Kasumigaura / Ibaraki (1979-08)
Unialgal, Clonal, N.Takamura (1979-08)
Identified by: N.Takamura
Culture conditions: MA, 25° C, 24 µE/m² sec, 20D,
[Cryopreserved]
Characteristics: Water bloom, Freshwater
KN1133
References: 52, 103
- Microcystis aeruginosa* (Kützing) Lemmermann
f. *flos-aquae* (Wittrock) Elenkin
98
Lake Kasumigaura / Ibaraki (1982-09)
Axenic, Clonal, M.H.Watanabe (1982-09)
Identified by: M.H.Watanabe
Culture conditions: MA, 25° C, 24 µE/m² sec, 20D,
[Cryopreserved]
Characteristics: Water bloom, Indicator, Freshwater
K-MF-K-3
References: 2, 89, 169, 184, 216, 386, 399
- 478
Lake Kasumigaura / Ibaraki (1977-09)
Unialgal, Non-clonal, O.Yagi (1978-04)
Identified by: O.Yagi
Culture conditions: MA, 20° C, 4 µE/m² sec, 3M,
(25° C, 15 µE/m² sec), [Cryopreserved]
Characteristics: Freshwater
K-5
References: 104, 418, 419, 420
- Microcystis elabens* Kützing var. *minor* Nygaard
42
Lake Kasumigaura / Ibaraki (1974-08)
IAM M-177, Unialgal, Clonal, M.M.Watanabe
(1974-08)
Identified by: M.M.Watanabe
Culture conditions: CT, 25° C, 24 µE/m² sec, 1M,
[Cryopreserved]
Characteristics: Water bloom, Freshwater
References: 80, 215, 216, 299, 425, 431
- Microcystis holsatica* Lemmermann
43
Lake Kasumigaura / Ibaraki (1974-08)
IAM M-179, Unialgal, Clonal, M.M.Watanabe
(1974-08)
Identified by: M.M.Watanabe
Culture conditions: CT, 25° C, 24 µE/m² sec, 1M,
[Cryopreserved]
Characteristics: Water bloom, Freshwater
References: 80, 216, 299, 425
- Microcystis viridis* (A.Brown) Lemmermann
102
Lake Kasumigaura / Ibaraki (1982-09)
Axenic, Clonal, M.H.Watanabe (1982-09)
Identified by: M.H.Watanabe
Culture conditions: MA, 25° C, 24 µE/m² sec, 1M,
[Cryopreserved]
Characteristics: Water bloom, Indicator, Toxic,

- Freshwater
K-MV-20
References: 98, 107, 109, 126, 133, 164, 169, 172,
215, 216, 272, 287, 332, 333, 389, 411, 425, 431
- 103
Lake Kasumigaura / Ibaraki (1978-12)
TAC 44, Unialgal, Clonal, M.Watanabe (1978-12)
Identified by: M.Watanabe
Culture conditions: MA, 25° C, 24 µE/m² sec, 1M,
[Cryopreserved]
Characteristics: Water bloom, Indicator, Toxic,
Freshwater
K-TAN-44
References: 185, 386
- Microcystis wesenbergii* Komárek
- 104
Chiyoda-ku / Tokyo (1982-11)
Axenic, Clonal, M.H.Watanabe (1982-11)
Identified by: M.H.Watanabe
Culture conditions: CB, MA, 25° C, 24 µE/m² sec,
1M, [Cryopreserved]
Characteristics: Water bloom, Indicator, Freshwater
MW-H1
References: 169, 325, 425
- 105
Lake Kasumigaura / Ibaraki (1982-09)
Unialgal, Clonal, M.H.Watanabe (1982-09)
Identified by: M.H.Watanabe
Culture conditions: MA, 25° C, 24 µE/m² sec, 1M,
[Cryopreserved]
Characteristics: Water bloom, Indicator, Freshwater
K-MW-K4
- 106
Lake Kasumigaura / Ibaraki (1982-09)
Unialgal, Clonal, M.H.Watanabe (1982-09)
Identified by: M.H.Watanabe
Culture conditions: MA, 25° C, 24 µE/m² sec, 1M,
[Cryopreserved]
Characteristics: Water bloom, Indicator, Freshwater,
(A) large size
K-MW-19
Reference: 287
- 107
Lake Kawaguchi / Yamanashi (1981-06)
Unialgal, Clonal, M.H.Watanabe (1981-06)
Identified by: M.H.Watanabe
Culture conditions: CB, MA, 25° C, 24 µE/m² sec,
1M, [Cryopreserved]
Characteristics: Water bloom, Indicator, Freshwater
KW-MW-7
References: 215, 216, 411
- 108
Lake Suwa / Nagano (1982-08)
Unialgal, Clonal, M.H.Watanabe (1982-08)
Identified by: M.H.Watanabe
Culture conditions: MA, 25° C, 24 µE/m² sec, 1M,
[Cryopreserved]
Characteristics: Water bloom, Indicator, Freshwater
S-MW-52
- 109
Lake Yogo / Shiga (1982-07)
Unialgal, Clonal, M.H.Watanabe (1982-07)
Identified by: M.H.Watanabe
Culture conditions: MA, 25° C, 24 µE/m² sec, 1M,
[Cryopreserved]
Characteristics: Water bloom, Indicator, Freshwater
Y-MW-24
- 110
Lake Kasumigaura / Ibaraki (1978-08)
TAC 36, Unialgal, Clonal, M.Watanabe (1978-08)
Identified by: M.Watanabe
Culture conditions: CT, 25° C, 24 µE/m² sec, 1M,
[Cryopreserved]
Characteristics: Water bloom, Indicator, Freshwater
K-TAN-36
- 111
Lake Kasumigaura / Ibaraki (1978-08)
TAC 37, Axenic, Clonal, M.Watanabe (1978-08)
Identified by: M.Watanabe
Culture conditions: MA, 25° C, 24 µE/m² sec, 1M,
[Cryopreserved]
Characteristics: Water bloom, Indicator, Freshwater
K-TAN-37
References: 169, 215, 216, 325, 411
- 112
Lake Suwa / Nagano (1982-10)
TAC 52, Axenic, Clonal, M.Watanabe (1982-10)
Identified by: M.Watanabe
Culture conditions: MA, 25° C, 24 µE/m² sec, 1M,
[Cryopreserved]
Characteristics: Water bloom, Indicator, Freshwater
S-TAN-52
References: 169, 216, 386, 431

604

Lake Kasumigaura / Ibaraki (1977-09)
Axenic, Clonal, O.Yagi (1978-04)
Identified by: O.Yagi
Culture conditions: MA, 20° C, 4 μ E/m² sec, 3M,
(25° C, 15 μ E/m² sec), [Cryopreserved]
Characteristics: Water bloom, Freshwater, Formerly
identified as *Microcystis aeruginosa*
K-3A
References: 51, 163, 169, 195, 315, 418

Microthamnion kützingianum Nägeli
479

Toyohira River / Sapporo (1987-07)
Unialgal, Clonal, F.Kasai (1987-07)
Identified by: F.Kasai
Culture conditions: C, 10° C, 6 μ E/m² sec, 6M,
(10° C, 15 μ E/m² sec)
Characteristics: Freshwater
Tst11-6
References: 338, 339

Monomastix minuta Skuja
255

Tsuchiura / Ibaraki (1983-07)
Axenic, Clonal, S.Suda (1983-07)
Identified by: S.Suda
Culture conditions: C, 20° C, 40 μ E/m² sec, 1M
Characteristics: Freshwater
SIS-Mono
Reference: 126

256

Oze / Gunma (1983-08)
Axenic, Clonal, S.Suda (1983-11)
Identified by: S.Suda
Culture conditions: AF-6, 20° C, 40 μ E/m² sec, 1M
Characteristics: Freshwater
Oz-35-m

* *Monoraphidium capricornutum* (Printz) Nygaard
See *Selenastrum capricornutum* Printz

Monoraphidium circinale (Nygaard) Nygaard
480

Tsuchiura / Ibaraki (1983-07)
Axenic, Clonal, S.Suda (1983-07)
Identified by: F.Kasai
Culture conditions: C(S), 20° C, μ E/m² sec, 3M,
(25° C, 30 μ E/m² sec)
Characteristics: Freshwater

SIS-1-M

Monoraphidium contortum

(Thuret) Komárková-Legnerová
384
Lake Unagiike / Kagoshima (1985-02)
Unialgal, Clonal, T.Sawaguchi (1985-02)
Identified by: Y.Niiyama
Culture conditions: C, 20° C, 4 μ E/m² sec, 3M,
(25° C, 30 μ E/m² sec)
Characteristics: Freshwater
Ep-i

Monoraphidium griffithii

(Berkeley) Komárková-Legnerová
385
Urizura / Ibaraki (1984-10)
Axenic, Clonal, T.Sawaguchi (1984-12)
Identified by: Y.Niiyama
Culture conditions: C, 20° C, 4 μ E/m² sec, 3M,
(25° C, 30 μ E/m² sec)
Characteristics: Freshwater
AWA

Myxosarcina burmensis Skuja

481
Mt. Tsukuba / Ibaraki (1987-04)
Unialgal, Non-clonal, F.Kasai (1987-05)
Identified by: M.M.Watanabe
Culture conditions: MDM(S), 20° C, 4 μ E/m² sec,
5M, (20° C, 12 μ E/m² sec), [Cryopreserved]
Characteristics: Freshwater
(1)-45
Reference: 338

Nephroselmis astigmatica Inouye et Pienaar

252
Tateyama Harbor / Chiba (1983-08)
Axenic, Clonal, I.Inouye (1983-08)
Identified by: I.Inouye
Culture conditions: f/2, ESM, 20° C, 32 μ E/m² sec,
1M
Characteristics: Red tide, Marine
810-13

Nephroselmis olivacea Stein

483
Tsuchiura / Ibaraki (1986-02)
Axenic, Clonal, S.Suda (1986-05)
Identified by: S.Suda
Culture conditions: AF-6, 20° C, 22 μ E/m² sec, 20D

- Characteristics: Freshwater, Heterothallic,
Mating type +
S-N-2-1
References: 171, 328
- 484
Tsuchiura / Ibaraki (1986-02)
Axenic, Clonal, S.Suda (1986-05)
Identified by: S.Suda
Culture conditions: AF-6, 20° C, 22 μ E/m²sec, 20D
Characteristics: Freshwater, Heterothallic,
Mating type –
S-N-5-8
References: 354, 355
- 485
Tsuchiura / Ibaraki (1986-02)
Axenic, Clonal, S.Suda (1986-05)
Identified by: S.Suda
Culture conditions: AF-6, 20° C, 22 μ E/m²sec, 20D
Characteristics: Freshwater, Heterothallic,
Mating type –
S-N-3-4
References: 171, 328
- Nephroselmis viridis* Inouye, nom. nud.
486
Harima-Nada / Seto Inland Sea (1983-02)
Axenic, Clonal, S.Suda (1983-09)
Identified by: I.Inouye
Culture conditions: ESM, 20° C, 12 μ E/m²sec, 1M
Characteristics: Red tide, Marine, Type strain
H-70-2
- Nitzschia palea* (Kützing) W.Smith
487
Miyata River / Ibaraki (1987-04)
Unialgal, Non-clonal, F.Kasai (1987-05)
Identified by: N.Takamura
Culture conditions: CSi, 15° C, 15 μ E/m²sec, 2M
Characteristics: Freshwater
3st-0-57
Reference: 338
- 488
Miyata River / Ibaraki (1987-02)
Unialgal, Non-clonal, F.Kasai (1987-03)
Identified by: N.Takamura
Culture conditions: CSi, 15° C, 15 μ E/m²sec, 2M
Characteristics: Freshwater
1st-3-39
Reference: 338
- 489
Ashio / Gunma (1987-08)
Unialgal, Clonal, F.Kasai (1987-08)
Identified by: N.Takamura
Culture conditions: CSi, 15° C, 15 μ E/m²sec, 1M
Characteristics: Freshwater
Ast-2-2
References: 338, 339
- Nostoc commune* Vaucher ex Bornet et Flahault
24
Kurobe Valley / Toyama
IAM M-13, Unialgal, Non-clonal, A.Watanabe
Identified by: H.Fukushima
Culture conditions: MDM(S), 20° C, 4 μ E/m²sec,
4M, (25° C, 30 μ E/m²sec), [Cryopreserved]
Characteristics: Freshwater, Reidentified by
M.M.Watanabe
References: 80, 216, 336, 366, 399
- 38
Marble Point
IAM M-115, Unialgal, Non-clonal, O.Holm-Hansen
Identified by: M.M.Watanabe
Culture conditions: MDM(S), 20° C, 4 μ E/m²sec,
4M, (25° C, 30 μ E/m²sec), [Cryopreserved]
Characteristics: Freshwater, From dry lichens and
algae in sand
M-48-a
Reference: 80
- Nostoc linckia* Bornet ex Bornet et Flahault
25
Kagoshima / Kagoshima
IAM M-16, Axenic, Non-clonal, M.Ishikawa
Identified by: M.M.Watanabe
Culture conditions: MDM(S), 20° C, 4 μ E/m²sec,
4M, (25° C, 30 μ E/m²sec)
Characteristics: Freshwater
Reference: 366
- Nostoc linckia* Bornet ex Bornet et Flahault
var. *arvense* C.B.Rao
28
Kagoshima / Kagoshima
IAM M-30, Axenic, Non-clonal, M.Ishikawa
Identified by: Fukushima/Maruyama
Culture conditions: MDM(S), 20° C, 4 μ E/m²sec,
4M, (25° C, 30 μ E/m²sec), [Cryopreserved]
Characteristics: Freshwater, Reidentified by
M.M.Watanabe

- References: 80, 366
- Nostoc minutum* Desmazières ex Bornet et Flahault
26
Ishigaki Isl. / Okinawa
IAM M-17, Unialgal, Non-clonal, M.Ishikawa
Identified by: M.M.Watanabe
Culture conditions: MDM(S), 20° C, 4 µE/m² sec,
4M, (25° C, 30 µE/m² sec), [Cryopreserved]
Characteristics: Freshwater, Chromatic adaptation
References: 192, 269, 366, 399
- 29
Ishigaki Isl. / Okinawa
IAM M-31, Unialgal, Non-clonal, M.Ishikawa
Identified by: M.M.Watanabe
Culture conditions: MDM(S), 20° C, 4 µE/m² sec,
4M, (25° C, 30 µE/m² sec), [Cryopreserved]
Characteristics: Freshwater
References: 366, 399
- Odontella aurita* Agardh
589
Penzance / England (1991-03)
Unialgal, Non-clonal, S.Ono (1991-04)
Identified by: S.Ono
Culture conditions: f/2, 15° C, 10 µE/m² sec, 1M
Characteristics: Red tide, Marine
St-22
- Odontella longicuris* (Greville) Hoban
590
Hitachi / Ibaraki (1990-09)
Unialgal, Non-clonal, S.Ono (1990-10)
Identified by: S.Ono
Culture conditions: f/2, 15° C, 10 µE/m² sec, 1M
Characteristics: Red tide, Marine
St-11
- Oedogonium obesum* Witrock ex Hirn
203
IAM C-348, Axenic, Clonal, E.Saito
Identified by: E.Saito
Culture conditions: C, 20° C, 4 µE/m² sec, 3M,
(25° C, 30 µE/m² sec)
Characteristics: Indicator, Freshwater
807
Reference: 80
- Olisthodiscus luteus* Carter
15
Tamano / Okayama / Seto Inland Sea
Axenic, Clonal, I.Inouye
Identified by: I.Inouye
Culture conditions: f/2, 20° C, 12 µE/m² sec, 1M
Characteristics: Red tide, Marine, Untransportable
Olisth
References: 56, 194, 361, 435
- Oltmannsiellopsis geminata* Inouye et Chihara
672
Harima-Nada / Seto Inland Sea (1986-06)
Axenic, Clonal, S.Yoshimatsu (1986-06)
Identified by: S.Yoshimatsu
Culture conditions: ESM, 20° C, 40 µE/m² sec, 1M
Characteristics: Marine, Mutant
- Oltmannsiellopsis unicellularis* Inouye et Chihara
359
Ieshima Isls. / Hyogo (1984-08)
Axenic, Clonal, S.Suda (1984-08)
Identified by: I.Inouye
Culture conditions: ESM, 20° C, 12 µE/m² sec, 2M
Characteristics: Red tide, Marine, Type strain
810YB-6
Reference: 14
- Oltmannsiellopsis viridis*
(Hargraves et Steele) Chihara et Inouye
360
Onagawa Bay / Miyagi (1984-08)
Axenic, Clonal, S.Suda (1984-09)
Identified by: S.Suda
Culture conditions: ESM, 20° C, 32 µE/m² sec, 2M
Characteristics: Marine, 18S rDNA gene (D86495)
8280G41-2
References: 14, 213
- Oocystis borgei* Snow
659
Watarase River / Gunma (1987-08)
Unialgal, F.Kasai (1987-09)
Identified by: F.Kasai
Culture conditions: C, 15° C, 6 µE/m² sec, 6M,
(15° C, 15 µE/m² sec)
Characteristics: Freshwater
AT2-26
Reference: 338
- Oocystis lacustris* Chodat
660
Watarase River / Gunma (1987-08)
Unialgal, Clonal, F.Kasai (1987-08)

- Identified by: F.Kasai
 Culture conditions: C, 15° C, 6 $\mu\text{E}/\text{m}^2\text{sec}$, 6M,
 (15° C, 15 $\mu\text{E}/\text{m}^2\text{sec}$)
 Characteristics: Freshwater
 Ast-3-1
 Reference: 338
- 661
 Miyata River / Ibaraki (1987-05)
 Unialgal, Clonal, F.Kasai (1987-06)
 Identified by: F.Kasai
 Culture conditions: C, 20° C, 8 $\mu\text{E}/\text{m}^2\text{sec}$, 6M
 Characteristics: Freshwater
 4st-3-9
 Reference: 338
- 662
 Miyata River / Ibaraki (1987-02)
 Axenic, Clonal, F.Kasai (1987-03)
 Identified by: F.Kasai
 Culture conditions: C, 20° C, 8 $\mu\text{E}/\text{m}^2\text{sec}$, 6M
 Characteristics: Freshwater
 1st-2-9
 References: 337, 338
- Oscillatoria agardhii* Gomont
 204
 Lake Kasumigaura / Ibaraki (1983-08)
 Axenic, Clonal, S.Suda (1983-08)
 Identified by: S.Suda
 Culture conditions: CB, 25° C, 24 $\mu\text{E}/\text{m}^2\text{sec}$, 1M,
 [Cryopreserved]
 Characteristics: Water bloom, Indicator, Freshwater
 K-O-A
 References: 169, 188, 216, 316, 318, 320, 321, 399,
 417
- 205
 Lake Kasumigaura / Ibaraki (1982-09)
 TAC 53, Unialgal, Clonal, M.Watanabe (1982-09)
 Identified by: M.Watanabe
 Culture conditions: MA, 25° C, 24 $\mu\text{E}/\text{m}^2\text{sec}$, 1M,
 [Cryopreserved]
 Characteristics: Water bloom, Indicator, Freshwater
 K-TAN-53
 References: 169, 317
- 594
 North Ireland / U.K.
 Axenic, Clonal
 Culture conditions: CT, 20° C, 4 $\mu\text{E}/\text{m}^2\text{sec}$, 2M,
 (20° C, 12 $\mu\text{E}/\text{m}^2\text{sec}$), [Cryopreserved]
 Characteristics: Freshwater
 k-8
- 595
 North Ireland / U.K.
 Axenic, Clonal
 Culture conditions: CT, 20° C, 4 $\mu\text{E}/\text{m}^2\text{sec}$, 2M,
 (20° C, 12 $\mu\text{E}/\text{m}^2\text{sec}$), [Cryopreserved]
 Characteristics: Freshwater
 3A②
- 596
 Veluwemeer / Holland
 Axenic, Clonal
 Culture conditions: CT, 20° C, 4 $\mu\text{E}/\text{m}^2\text{sec}$, 2M,
 (20° C, 12 $\mu\text{E}/\text{m}^2\text{sec}$), [Cryopreserved]
 Characteristics: Freshwater
 VLOA7
 Reference: 26
- 610
 See *Oscillatoria rubescens* (DC.) ex Gomont
- Oscillatoria amphibia* Agardh ex Gomont
 361
 Asaji Bay / Nagasaki (1985-07)
 Unialgal, Clonal, M.M.Watanabe (1985-07)
 Identified by: M.M.Watanabe
 Culture conditions: f/2, 20° C, 12 $\mu\text{E}/\text{m}^2\text{sec}$, 1M,
 [Cryopreserved]
 Characteristics: Marine, Benthic
 Oa
- Oscillatoria animalis* Agardh ex Gomont
 206
 IAM M-75, Unialgal, Clonal, F.Murano
 Identified by: H.Fukushima
 Culture conditions: MDM(S), 20° C, 4 $\mu\text{E}/\text{m}^2\text{sec}$,
 4M, (25° C, 30 $\mu\text{E}/\text{m}^2\text{sec}$), [Cryopreserved]
 Characteristics: Freshwater, Reidentified by
 M.M.Watanabe
 Reference: 80
- Oscillatoria laetevirens* Gomont
 31
 Kawaji / Tochigi
 IAM M-42, Unialgal, Clonal, M.Ishikawa
 Identified by: H.Fukushima
 Culture conditions: MDM(S), 20° C, 4 $\mu\text{E}/\text{m}^2\text{sec}$,
 4M, (25° C, 30 $\mu\text{E}/\text{m}^2\text{sec}$), [Cryopreserved]

- Characteristics: Freshwater, Hot spring, Reidentified by M.M.Watanabe
References: 80, 399
- Oscillatoria limnetica* Lemmermann
36
Nakano / Tokyo
IAM M-92, Unialgal, Clonal, F.Murano
Identified by: H.Fukushima
Culture conditions: MDM(S), 20° C, 4 μ E/m² sec, 4M, (25° C, 30 μ E/m² sec), [Cryopreserved]
Characteristics: Freshwater, Reidentified by M.M.Watanabe
References: 4, 80, 263
- Oscillatoria raciborskii* Woloszynska
207
Lake Kasumigaura / Ibaraki (1983-06)
Axenic, Clonal, S.Suda (1983-06)
Identified by: S.Suda
Culture conditions: CB, CT, 25° C, 24 μ E/m² sec, 20D
Characteristics: Water bloom, Offensive taste and odor, Freshwater, Unstable
K-O-R
References: 169, 216, 399
- Oscillatoria rosea* Utermöhl
208
Asaji Bay / Nagasaki (1983-08)
Axenic, Clonal, Y.Ichimura (1983-08)
Identified by: M.M.Watanabe
Culture conditions: f/2, 20° C, 32 μ E/m² sec, 1M, [Cryopreserved]
Characteristics: Indicator, Marine
NGS-1
Reference: 300
- Oscillatoria rubescens* (DC.) ex Gomont
610
Lake Gjersjoen / Norway
CCAP 1459/22, Axenic, Romstad (1971)
Reidentified by: S.Suda
Culture conditions: CB, MA, 20° C, 40 μ E/m² sec, 1M, [Cryopreserved]
Characteristics: Freshwater, Formerly identified as *Oscillatoria agardhii* Gomont
NIVA CYA 18
References: 290, 291, 292
- Oscillatoria tenuis* Agardh ex Gomont
33
- Setagaya / Tokyo
IAM M-50, Unialgal, Clonal, M.Ishikawa
Identified by: K.Maruyama
Culture conditions: MDM(S), 20° C, 4 μ E/m² sec, 4M, (25° C, 30 μ E/m² sec), [Cryopreserved]
Characteristics: Indicator, Freshwater, Reidentified by M.M.Watanabe
Reference: 80
- Oxyrrhis marina* Dujardin
494
Hachinohe / Aomori (1988-08)
Mixed, Clonal, T.Sawaguchi (1989-01)
Identified by: T.Sawaguchi
Culture conditions: f/2, 20° C, 40 μ E/m² sec, 1M
Characteristics: Predator, Marine, Feeds on NIES-254, Untransportable
370OX
- Pandorina colemaniae* Nozaki
572
Kourakuen / Okayama (1988-10)
Unialgal, Clonal, H.Nozaki (1988-10)
Identified by: H.Nozaki
Culture conditions: AF-6, 20° C, 22 μ E/m² sec, 1M
Characteristics: Freshwater, Type strain, Isogamy, Mating type +, Crosses with NIES-573, *rbcL* gene (D63441)
88-1025-1
References: 243, 253
- 573
Kourakuen / Okayama (1988-10)
Unialgal, Clonal, H.Nozaki (1989-01)
Identified by: H.Nozaki
Culture conditions: AF-6, 20° C, 22 μ E/m² sec, 1M
Characteristics: Freshwater, Type strain, Isogamy, Mating type -, Crosses with NIES-572
89-0131-P-3
Reference: 253
- Pandorina morum* (O. F. Müller) Bory
242
Lake Ozenuma / Fukushima (1983-08)
Axenic, Clonal, S.Suda (1983-09)
Identified by: S.Suda
Culture conditions: CA, 20° C, 12 μ E/m² sec, 1M
Characteristics: Indicator, Freshwater, Heterothallic, Mating type +, Crosses with NIES-243 and 362
Oz-Pa-2

243

Lake Ozenuma / Fukushima (1983-08)
Axenic, Clonal, S.Suda (1983-09)
Identified by: S.Suda
Culture conditions: CA, 20° C, 12 μ E/m² sec, 1M
Characteristics: Freshwater, Heterothallic,
Mating type -, Crosses with NIES-242
Oz-Pa-3

362

Lake Ozenuma / Fukushima (1983-08)
Axenic, Clonal, S.Suda (1983-09)
Identified by: S.Suda
Culture conditions: CA, 20° C, 12 μ E/m² sec, 1M
Characteristics: Freshwater, Heterothallic,
Mating type -, Crosses with NIES-242
Oz-Pa-1

Pandorina morum (O. F. Müller) Bory var. *morum*
574

Nepal (1986-09)
Unialgal, Clonal, H.Nozaki (1987-09)
Identified by: H.Nozaki
Culture conditions: VT, 20° C, 22 μ E/m² sec, 1M
Characteristics: Freshwater, Heterothallic, Isogamy,
Mating type +, Crosses with NIES-575,
rbcL gene (D63442)
7916-P-7
References: 230, 243

575

Nepal (1986-09)
Unialgal, Clonal, H.Nozaki (1987-09)
Identified by: H.Nozaki
Culture conditions: AF-6, 20° C, 22 μ E/m² sec, 1M
Characteristics: Freshwater, Heterothallic, Isogamy,
Mating type -, Crosses with NIES-574
7916-P-8
Reference: 230

* *Pandorina unicocca* Rayburn et Starr
See *Yamagishiella unicocca*
(Rayburn et Starr) Nozaki

Paulschulzia pseudovolvox Skuja
727

Tvärminne / Finland
UTEX 167, Axenic, Clonal, M.R.Droop (1951)
Culture conditions: AF-6, 20° C, 12 μ E/m² sec, 1M
Characteristics: Freshwater, *rbcL* gene (D86837)
Reference: 244

Pavlova gyrans Butcher

623
Matoya Bay / Mie (1984-09)
Unialgal, Clonal, T.Sawaguchi (1984-09)
Identified by: S.Suda
Culture conditions: ESM, 20° C, 22 μ E/m² sec, 2M
Characteristics: Marine
MB-D-24

Pediastrum angulosum Meneghini
var. *angulosum*

300
Higashihiroshima / Hiroshima (1983-10)
Axenic, Clonal, F.Kasai (1983-10)
Identified by: M.Watanabe
Culture conditions: C, 20° C, 4 μ E/m² sec, 3M,
(25° C, 30 μ E/m² sec)
Characteristics: Freshwater
83-24-1-7

Pediastrum boryanum (Turpin) Meneghini
209

Lake Kasumigaura / Ibaraki (1982-12)
Axenic, Clonal, M.H.Watanabe (1982-12)
Identified by: M.H.Watanabe
Culture conditions: C, 20° C, 8 μ E/m² sec, 2M
Characteristics: Indicator, Freshwater,
COXI gene (D63659)
K-P-40
Reference: 66

301

Lake Shoji / Yamanashi (1981-10)
TAC 56-3A, Axenic, Clonal, M.Watanabe
Culture conditions: C, 20° C, 8 μ E/m² sec, 2M
Characteristics: Freshwater
TAN-56-3A
Reference: 199

Pediastrum duplex Meyen
212

Lake Kawaguchi / Yamanashi (1981-06)
Unialgal, Clonal, M.H.Watanabe (1981-06)
Identified by: M.H.Watanabe
Culture conditions: C, 20° C, 8 μ E/m² sec, 2M
Characteristics: Indicator, Freshwater
KW-P-1
Reference: 359

Pediastrum duplex Meyen var. *duplex*
210

Tsukuba / Ibaraki (1983-05)

- Axenic, Clonal, A. Yuri (1983-05)
 Identified by: A. Yuri
 Culture conditions: C, 20° C, 4 $\mu\text{E}/\text{m}^2\text{sec}$, 3M,
 (25° C, 30 $\mu\text{E}/\text{m}^2\text{sec}$)
 Characteristics: Indicator, Freshwater, Reidentified
 by M. Watanabe
 Pe-16
- 213
 Tsukuba / Ibaraki (1983-05)
 Axenic, Clonal, T. Hiwatari (1983-06)
 Identified by: T. Hiwatari
 Culture conditions: C, 20° C, 8 $\mu\text{E}/\text{m}^2\text{sec}$, 2M
 Characteristics: Indicator, Freshwater, Reidentified
 by M. Watanabe
 AQ-P-1
 References: 70, 399
- Pediastrum duplex* Meyen
 var. *gracillimum* W. et G.S. West
 211
 Lake Kasumigaura / Ibaraki (1983-08)
 Axenic, Clonal, F. Kasai (1983-08)
 Identified by: M. Watanabe
 Culture conditions: C(S), 20° C, 4 $\mu\text{E}/\text{m}^2\text{sec}$, 3M,
 (25° C, 30 $\mu\text{E}/\text{m}^2\text{sec}$)
 Characteristics: Indicator, Freshwater
 F50-1
- 214
 Tsukuba / Ibaraki (1983-08)
 Axenic, Clonal, T. Hiwatari (1983-08)
 Identified by: T. Hiwatari
 Culture conditions: C, 20° C, 8 $\mu\text{E}/\text{m}^2\text{sec}$, 2M
 Characteristics: Indicator, Freshwater, Reidentified
 by M. Watanabe
 KR-P-2
- Pediastrum simplex* Meyen
 215
 Lake Biwa / Shiga (1982-07)
 Axenic, Clonal, M.H. Watanabe (1982-07)
 Identified by: M.H. Watanabe
 Culture conditions: C, 20° C, 8 $\mu\text{E}/\text{m}^2\text{sec}$, 2M
 Characteristics: Indicator, Freshwater
 B-P-18
- 302
 Lake Kasumigaura / Ibaraki (1983-08)
 Axenic, Clonal, F. Kasai (1983-08)
 Culture conditions: C, 20° C, 4 $\mu\text{E}/\text{m}^2\text{sec}$, 3M,
 (25° C, 30 $\mu\text{E}/\text{m}^2\text{sec}$)
 Characteristics: Indicator, Freshwater
 F-26-4
- Pediastrum tetras* (Ehrenberg) Ralfs
 216
 Lake Kasumigaura / Ibaraki (1982-12)
 Axenic, Clonal, M.H. Watanabe (1982-12)
 Identified by: M.H. Watanabe
 Culture conditions: C, 20° C, 8 $\mu\text{E}/\text{m}^2\text{sec}$, 2M
 Characteristics: Indicator, Freshwater
 K-P-30
- Pedinomonas minor* Korshikov
 363
 Tsukuba / Ibaraki (1984-05)
 Axenic, Clonal, S. Suda (1984-05)
 Identified by: S. Suda
 Culture conditions: C(S), 20° C, 4 $\mu\text{E}/\text{m}^2\text{sec}$, 3M,
 (25° C, 30 $\mu\text{E}/\text{m}^2\text{sec}$)
 Characteristics: Freshwater
 H31P4
- Penium margaritaceum* Brébisson
 217
 Rumalbhara / Nepal (1965-11)
 IAM C-397, Axenic, Clonal, T. Ichimura (1972-05)
 Identified by: T. Ichimura
 Culture conditions: C, 20° C, 8 $\mu\text{E}/\text{m}^2\text{sec}$, 2M,
 (25° C, 30 $\mu\text{E}/\text{m}^2\text{sec}$)
 Characteristics: Indicator, Freshwater, Heterothallic
 N-76-20
 Reference: 80
- 303
 Tsukiyono / Gunma (1984-06)
 Axenic, Clonal, F. Kasai (1984-06)
 Identified by: F. Kasai
 Culture conditions: C, 20° C, 8 $\mu\text{E}/\text{m}^2\text{sec}$, 3M,
 (25° C, 30 $\mu\text{E}/\text{m}^2\text{sec}$)
 Characteristics: Freshwater
 84-25-1
- Peridinium bipes* Stein f. *globosum* Lindermann
 495
 Lake Onogawa / Fukushima (1985-07)
 Unialgal, Clonal, T. Sawaguchi (1985-08)
 Identified by: T. Sawaguchi
 Culture conditions: AF-6, 15° C, 35 $\mu\text{E}/\text{m}^2\text{sec}$, 2M
 Characteristics: Freshwater, Untransportable
 LOND-9

Peridinium bipes Stein

f. *occultatum* (Lindermann) Lefèvre

364

Lake Unagiike / Kagoshima (1985-02)

Axenic, Clonal, T.Sawaguchi (1985-02)

Identified by: T.Sawaguchi

Culture conditions: URO, Carefoot, 15° C,

35 µE/m² sec, 2M

Characteristics: Red tide, Freshwater,

Untransportable

EPD-7

496

Isobe / Mie (1986-10)

Unialgal, Clonal, T.Sawaguchi (1986-11)

Identified by: T.Sawaguchi

Culture conditions: URO, 15° C, 35 µE/m² sec, 2M

Characteristics: Red tide, Freshwater,

Untransportable

KDD-1

497

Lake Kizaki / Nagano (1988-04)

Unialgal, Clonal, T.Sawaguchi (1988-04)

Identified by: T.Sawaguchi

Culture conditions: Carefoot, 15° C, 35 µE/m² sec,

2M

Characteristics: Red tide, Freshwater,

Untransportable

LK420

Peridinium bipes Stein var. *tabulatum*

(Ehrenberg) Lefèvre

600

Shishizuka / Tsuchiura / Ibaraki (1990-04)

Unialgal, Clonal, T.Hagiwara (1990-04)

Identified by: T.Hagiwara

Culture conditions: URO, 15° C, 35 µE/m² sec, 3M

Characteristics: Red tide, Freshwater, Planktonic,

Untransportable

CCZ-1

Peridinium inconspicuum Lemmermann

subsp. *remotum* (Lefèvre) Lefèvre

499

Iwai / Ibaraki (1985-10)

Unialgal, Clonal, T.Sawaguchi (1985-11)

Identified by: T.Sawaguchi

Culture conditions: MW/5, 15° C, 35 µE/m² sec,

2M

Characteristics: Freshwater, Untransportable

TOM-1

Peridinium polonicum Woloszynska

500

Shiogama / Miyagi (1988-07)

Axenic, Clonal, T.Sawaguchi (1988-07)

Identified by: T.Sawaguchi

Culture conditions: AF-6/2, 20° C, 40 µE/m² sec,

2M

Characteristics: Freshwater, Untransportable

KAP-2

Peridinium volzii Lemmermann

365

Ajiro / Iwate (1984-09)

Axenic, Clonal, T.Sawaguchi (1984-09)

Identified by: T.Sawaguchi

Culture conditions: Carefoot, 15° C, 35 µE/m² sec,

2M

Characteristics: Freshwater, Untransportable

HND-1

501

Tsuchiura / Ibaraki (1986-04)

Unialgal, Clonal, T.Sawaguchi (1986-05)

Culture conditions: Carefoot, 15° C, 35 µE/m² sec,

2M

Characteristics: Freshwater, Homothallic,

Untransportable

SPSP-2

Peridinium wierzejskii Woloszynska

502

Tsuchiura / Ibaraki (1985-04)

Unialgal, Clonal, T.Sawaguchi (1985-04)

Identified by: T.Sawaguchi

Culture conditions: MW/5, 15° C, 35 µE/m² sec, 2M

Characteristics: Freshwater, Homothallic,

Untransportable

SPD-7

Peridinium willei Huitfeldt-Kaas

304

Tsukiyono / Gunma (1984-06)

Axenic, Clonal, T.Sawaguchi (1984-06)

Identified by: T.Sawaguchi

Culture conditions: Carefoot, 15° C, 35 µE/m² sec,

2M

Characteristics: Freshwater, Homothallic,

Untransportable

8423-P

366

Tsuchiura / Ibaraki (1985-04)

- Axenic, Clonal, T.Sawaguchi (1985-04)
 Identified by: T.Sawaguchi
 Culture conditions: Carefoot, 15° C, 35 $\mu\text{E}/\text{m}^2\text{sec}$,
 2M
 Characteristics: Freshwater, Homothallic,
 Untransportable
 SPD-1
- Phacus agilis* Skuja
 387
 Kashiwa / Chiba (1986-09)
 Axenic, Clonal, M.M.Watanabe (1986-09)
 Identified by: M.M.Watanabe
 Culture conditions: MAF-6, AF-6, 20° C,
 32 $\mu\text{E}/\text{m}^2\text{sec}$, 1M
 Characteristics: Freshwater, Umetatechi-
 shinshutsusui lagoon
 PhD-3
- Phaeocystis pouchetii* (Hariot) Lagerheim
 388
 Hachijo Isl. / Tokyo (1984-04)
 Unialgal, Non-clonal, T.Sawaguchi (1984-04)
 Identified by: T.Sawaguchi
 Culture conditions: ESM, 15° C, 20 $\mu\text{E}/\text{m}^2\text{sec}$, 20D,
 (20° C, 40 $\mu\text{E}/\text{m}^2\text{sec}$)
 Characteristics: Red tide, Marine, Unstable,
 Untransportable, *COXI* gene (AB000120)
 8-P
 Reference: 65
- Phormidium foveolarum* Gomont
 32
 Lake Shirakaba / Nagano
 IAM M-43, Unialgal, Non-clonal, M.Ishikawa
 Identified by: H.Fukushima
 Culture conditions: MDM(S), 20° C, 4 $\mu\text{E}/\text{m}^2\text{sec}$,
 4M, (25° C, 30 $\mu\text{E}/\text{m}^2\text{sec}$), [Cryopreserved]
 Characteristics: Freshwater, Reidentified by
 M.M.Watanabe
 References: 80, 346, 356
- 34
 Sendai / Miyagi
 IAM M-59, Unialgal, Non-clonal, M.Ishikawa
 Identified by: K.Maruyama
 Culture conditions: MDM(S), 20° C, 4 $\mu\text{E}/\text{m}^2\text{sec}$,
 4M, (25° C, 30 $\mu\text{E}/\text{m}^2\text{sec}$), [Cryopreserved]
 Characteristics: Freshwater, Reidentified by
 M.M.Watanabe
 Reference: 80
- 503
 Mt. Tsukuba / Ibaraki (1987-04)
 Unialgal, Non-clonal, F.Kasai (1987-05)
 Identified by: M.M.Watanabe
 Culture conditions: CSi, CSi+Cu, 20° C,
 4 $\mu\text{E}/\text{m}^2\text{sec}$, 3M, (20° C, 12 $\mu\text{E}/\text{m}^2\text{sec}$),
 [Cryopreserved]
 Characteristics: Freshwater
 (1)-48
 Reference: 338
- 504
 Miyata River / Ibaraki (1987-03)
 Unialgal, Non-clonal, F.Kasai (1987-05)
 Identified by: M.M.Watanabe
 Culture conditions: CSi, CSi+Cu, 20° C,
 4 $\mu\text{E}/\text{m}^2\text{sec}$, 3M, (20° C, 12 $\mu\text{E}/\text{m}^2\text{sec}$),
 [Cryopreserved]
 Characteristics: Freshwater
 2st-2-4
 References: 337, 338, 339
- 505
 Watarase River / Gunma (1987-08)
 Unialgal, Non-clonal, F.Kasai (1987-10)
 Identified by: M.M.Watanabe
 Culture conditions: CSi, CSi+Cu, 20° C,
 4 $\mu\text{E}/\text{m}^2\text{sec}$, 2M, (20° C, 12 $\mu\text{E}/\text{m}^2\text{sec}$),
 [Cryopreserved]
 Characteristics: Freshwater
 AT4-17
 References: 338, 339
- Phormidium jenkelianum* G.Schmid
 506
 Watarase River / Gunma (1987-08)
 Unialgal, Non-clonal, F.Kasai (1987-09)
 Identified by: M.M.Watanabe
 Culture conditions: CSi, CSi+Cu, 20° C, 4 $\mu\text{E}/\text{m}^2\text{sec}$,
 2M, (20° C, 12 $\mu\text{E}/\text{m}^2\text{sec}$), [Cryopreserved]
 Characteristics: Freshwater
 AT5-37
 Reference: 338
- 507
 Watarase River / Gunma (1987-08)
 Unialgal, Non-clonal, F.Kasai (1987-08)
 Identified by: M.M.Watanabe
 Culture conditions: CSi, CSi+Cu, 20° C, 4 $\mu\text{E}/\text{m}^2\text{sec}$,
 2M, (20° C, 12 $\mu\text{E}/\text{m}^2\text{sec}$), [Cryopreserved]
 Characteristics: Freshwater

- Ast-1-4
References: 338, 339
- Phormidium molle* Gomont
509
Watarase River / Gunma (1987-08)
Unialgal, Non-clonal, F.Kasai (1987-08)
Identified by: M.M.Watanabe
Culture conditions: CSi, CSi+Cu, 20° C, 4 µE/m² sec,
2M, (20° C, 12 µE/m² sec), [Cryopreserved]
Characteristics: Freshwater
AT2-17
References: 338, 339
- Phormidium mucicola* Huber-Pestalozzi et Naum
510
Mt. Tsukuba / Ibaraki (1987-04)
Unialgal, Non-clonal, F.Kasai (1987-05)
Identified by: M.M.Watanabe
Culture conditions: CSi, CSi+Cu, 20° C, 4 µE/m² sec,
4M, (20° C, 12 µE/m² sec), [Cryopreserved]
Characteristics: Freshwater
(1)-23
Reference: 338
- Phormidium ramosum* Boye-Petersen
305
Takatori River / Ibaraki (1984-12)
Unialgal, Clonal, S.Suda (1984-12)
Identified by: S.Suda
Culture conditions: CSi, CSi+Cu, 20° C, 4 µE/m² sec,
4M, (20° C, 12 µE/m² sec), [Cryopreserved]
Characteristics: Freshwater
841211St5-1
References: 337, 338
- Phormidium tenue* Gomont
30
Akita / Akita
IAM M-40, Unialgal, Non-clonal, M.Ishikawa
Identified by: H.Fukushima
Culture conditions: MDM(S), 20° C, 4 µE/m² sec,
4M, (25° C, 30 µE/m² sec), [Cryopreserved]
Characteristics: Freshwater, Reidentified by
M.M.Watanabe
References: 80, 322
- 512
Nagoya / Aichi (1981-11)
Axenic, Non-clonal, N.Yamada (1985-05)
Identified by: N.Yamada
Culture conditions: CT, 20° C, 4 µE/m² sec, 20D,
(20° C, 12 µE/m² sec), [Cryopreserved]
Characteristics: Offensive taste and odor,
Freshwater, Nakaku Honmaru (a moat of the
Nagoya Castle)
PM-81A
References: 166, 167, 263, 423, 424
- 611
Lake Biwa / Shiga (1987-06)
Unialgal, Clonal, S.Ichise (1987-06)
Identified by: M.M.Watanabe
Culture conditions: CT, 25° C, 30 µE/m² sec, 1M,
[Cryopreserved]
Characteristics: Freshwater
Bpt
- Planctonema lauterbornii* Schmidle
514
Lake Kasumigaura / Ibaraki (1988-08)
Axenic, Clonal, Y.Niiyama (1988-08)
Identified by: Y.Niiyama
Culture conditions: C, 20° C, 8 µE/m² sec, 2M
Characteristics: Freshwater
K880818
- Platydorina caudata* Kofoid
728
Kansas / USA (1965-09)
UTEX 1658, Unialgal, Clonal, D.O.Harris
Culture conditions: AF-6, 20° C, 12 µE/m² sec, 1M
Characteristics: Freshwater, *rbcL* gene (D86828)
Reference: 244
- 729
Kansas / USA (1965-09)
UTEX 1661, Unialgal, Clonal, D.O.Harris
Culture conditions: AF-6, 20° C, 12 µE/m² sec, 1M
Characteristics: Freshwater, *rbcL* gene (D86827)
Reference: 244
- Plectonema radiosum* Gomont
515
Nikko / Tochigi (1987-04)
Axenic, Clonal, F.Kasai (1987-04)
Identified by: M.M.Watanabe
Culture conditions: CSi, 20° C, 4 µE/m² sec, 3M,
(20° C, 12 µE/m² sec), [Cryopreserved]
Characteristics: Freshwater
NK-12
References: 173, 338, 339

Pleodorina californica Shaw

576

Hachiman / Gifu (1990-08)

Axenic, Clonal, Y.Ogasawara (1990-08)

Identified by: Y.Ogasawara

Culture conditions: VT, 25° C, 30 µE/m²sec, 1M

Characteristics: Freshwater

735

Indiana / USA

UTEX 809, Axenic, Clonal, R.C.Starr

Culture conditions: AF-6, 20° C, 12 µE/m²sec, 1M

Characteristics: Freshwater, *rbcl* gene (D63439)

Reference: 243

Pleodorina indica (Iyengar) Nozaki

736

Mexico

UTEX 1990, Unialgal, Clonal, S.Morro

Reidentified by: H.Nozaki

Culture conditions: AF-6, 20° C, 12 µE/m²sec, 1M

Characteristics: Freshwater, *rbcl* gene (D86834)

Reference: 244

Pleodorina japonica Nozaki

577

Fuji / Shizuoka (1986-07)

UTEX 2523, Unialgal, Clonal, H.Nozaki (1986-07)

Identified by: H.Nozaki

Culture conditions: AF-6, 20° C, 22 µE/m²sec, 1M

Characteristics: Freshwater, Type strain, Homothallic,

Dioecious, Anisogamy, *rbcl* gene (D63440)

6715-7

References: 243, 250

Pleurotaenium cylindricum (Turner) Schmidle

var. *stuhlmannii* (Hieronymus) Krieger

306

Niimi / Okayama (1983-09)

Unialgal, Clonal, F.Kasai (1983-09)

Identified by: F.Kasai

Culture conditions: MG, 25° C, 15 µE/m²sec, 1M

Characteristics: Freshwater, Homothallic

F57-18-4

Pleurotaenium ehrenbergii (Ralfs) De Bary

var. *curtum* Krieger

307

Naka-gun / Wakayama (1969-10)

IAM C-378, Axenic, Clonal, T.Ichimura (1969-11)

Identified by: T.Ichimura

Culture conditions: CA, 20° C, 8 µE/m²sec, 3M,

(25° C, 30 µE/m²sec)

Characteristics: Freshwater, Heterothallic,

Mating type +, Crosses with NIES-308

W-1-1

308

Naka-gun / Wakayama (1969-10)

IAM C-379, Axenic, Clonal, T.Ichimura (1969-11)

Identified by: T.Ichimura

Culture conditions: CA, 20° C, 8 µE/m²sec, 3M,

(25° C, 30 µE/m²sec)

Characteristics: Freshwater, Heterothallic,

Mating type -, Crosses with NIES-307

W-1-3

311

Iriomote Isl. / Okinawa (1973-06)

IAM C-430, Unialgal, Clonal, T.Ichimura (1973-11)

Culture conditions: MG, 20° C, 8 µE/m²sec, 3M,

(25° C, 30 µE/m²sec)

Characteristics: Freshwater, Heterothallic,

Mating type +

R-13-19

Pleurotaenium ehrenbergii (Ralfs) De Bary

var. *ehrenbergii*

309

Iriomote Isl. / Okinawa (1973-06)

IAM C-467, Unialgal, Clonal, T.Ichimura (1973-10)

Culture conditions: MG, 20° C, 8 µE/m²sec, 3M,

(25° C, 30 µE/m²sec)

Characteristics: Freshwater, Heterothallic,

Mating type +, Crosses with NIES-310

R-13-27

Reference: 80

310

Iriomote Isl. / Okinawa (1973-06)

IAM C-468, Unialgal, Clonal, T.Ichimura (1973-10)

Culture conditions: MG, 20° C, 8 µE/m²sec, 3M,

(25° C, 30 µE/m²sec)

Characteristics: Freshwater, Heterothallic,

Mating type -, Crosses with NIES-309

R-13-30

Reference: 80

Pleurotaenium nodosum (Bailey ex Ralfs) Lundell

var. *borgei* Grönblad

663

Miyatoko Mire / Fukushima (1993-09)

Unialgal, Clonal, H.Nozaki (1993-09)

- Identified by: H.Nozaki
 Culture conditions: AF-6, 20° C, 22 $\mu\text{E}/\text{m}^2\text{sec}$, 2M
 Characteristics: Freshwater
 93-913-Gon-1
- 664
 Miyatoko Mire / Fukushima (1993-09)
 Unialgal, Clonal, H.Nozaki (1993-09)
 Identified by: H.Nozaki
 Culture conditions: AF-6, 20° C, 22 $\mu\text{E}/\text{m}^2\text{sec}$, 2M
 Characteristics: Freshwater
 93-913-Gon-3
- Pleurotaenium nodosum* (Bailey ex Ralfs) Lundell
 var. *gutwinskii* Krieger
 787
 4 km northwest of Melaka / Malaysia (1985-08)
 Unialgal, Clonal, T.Ichimura (1985)
 Identified by: T.Ichimura
 Culture conditions: MG, 20° C, 8 $\mu\text{E}/\text{m}^2\text{sec}$, 3M,
 (25° C, 30 $\mu\text{E}/\text{m}^2\text{sec}$)
 Characteristics: Freshwater, Heterothallic, Crosses
 with NIES-788
 85-30-9
- 788
 4 km northwest of Melaka / Malaysia (1985-08)
 Unialgal, Clonal, T.Ichimura (1985)
 Identified by: T.Ichimura
 Culture conditions: MG, 20° C, 8 $\mu\text{E}/\text{m}^2\text{sec}$, 3M,
 (25° C, 30 $\mu\text{E}/\text{m}^2\text{sec}$)
 Characteristics: Freshwater, Heterothallic, Crosses
 with NIES-787
 85-30-56
- Pleurotaenium nodosum* (Bailey ex Ralfs) Lundell
 var. *nodosum*
 312
 Higashihiroshima / Hiroshima (1983-10)
 Unialgal, Clonal, F.Kasai (1983-10)
 Identified by: F.Kasai
 Culture conditions: CAM, 20° C, 8 $\mu\text{E}/\text{m}^2\text{sec}$, 3M,
 (25° C, 30 $\mu\text{E}/\text{m}^2\text{sec}$)
 Characteristics: Freshwater
 83-24-3
- 785
 Imuta-ike Pond / Kagoshima (1986-10)
 Unialgal, Clonal, T.Ichimura (1986)
 Identified by: T.Ichimura
 Culture conditions: MG, 20° C, 8 $\mu\text{E}/\text{m}^2\text{sec}$, 3M,
 (25° C, 30 $\mu\text{E}/\text{m}^2\text{sec}$)
 Characteristics: Freshwater, Heterothallic, Crosses
 with NIES-786
 86-7-15
- 786
 Imuta-ike Pond / Kagoshima (1986-10)
 Unialgal, Clonal, T.Ichimura (1986)
 Identified by: T.Ichimura
 Culture conditions: MG, 20° C, 8 $\mu\text{E}/\text{m}^2\text{sec}$, 3M,
 (25° C, 30 $\mu\text{E}/\text{m}^2\text{sec}$)
 Characteristics: Freshwater, Heterothallic, Crosses
 with NIES-785
 86-7-16
- Pleurotaenium ovatum* Nordstedt
 313
 Niimi / Okayama (1983-09)
 Unialgal, Clonal, F.Kasai (1983-09)
 Identified by: F.Kasai
 Culture conditions: C, 20° C, 8 $\mu\text{E}/\text{m}^2\text{sec}$, 3M,
 (25° C, 30 $\mu\text{E}/\text{m}^2\text{sec}$)
 Characteristics: Freshwater
 F57-17-8
- Polyedriopsis spinulosa* (Schmidle) Schmidle
 232
 Tsukuba / Ibaraki (1984-05)
 Unialgal, Clonal, F.Kasai (1984-05)
 Identified by: F.Kasai
 Culture conditions: C, 20° C, 4 $\mu\text{E}/\text{m}^2\text{sec}$, 3M,
 (25° C, 30 $\mu\text{E}/\text{m}^2\text{sec}$)
 Characteristics: Freshwater
 F128
- Prorocentrum dentatum* Stein
 682
 Hiuchi-Nada / Seto Inland Sea (1979-12)
 Unialgal, Clonal, S.Yoshimatsu (1980-01)
 Identified by: S.Yoshimatsu
 Culture conditions: ESM, 20° C, 40 $\mu\text{E}/\text{m}^2\text{sec}$, 1M
 Characteristics: Red tide, Marine, Untransportable
- Prorocentrum gracile* Schütt
 315
 Harima-Nada / Seto Inland Sea
 Axenic, Clonal, S.Yoshimatsu (1984-08)
 Identified by: S.Yoshimatsu
 Culture conditions: ESM, 20° C, 32 $\mu\text{E}/\text{m}^2\text{sec}$, 1M
 Characteristics: Red tide, Marine, Untransportable
 80

Prorocentrum lima (Ehrenberg) Dodge

617

Motobu / Okinawa (1993-06)
Unialgal, Clonal, H.Kobayashi (1993-06)
Identified by: Y.Fukuyo
Culture conditions: f/2, ESM, 20° C, 32 µE/m² sec, 1M
Characteristics: Toxic, Marine, Untransportable
PL-03

Prorocentrum mexicanum Osorio Tafall

317

Harima-Nada / Seto Inland Sea
Axenic, Clonal, S.Yoshimatsu (1984-08)
Identified by: S.Yoshimatsu
Culture conditions: f/2, 20° C, 32 µE/m² sec, 1M
Characteristics: Red tide, Marine, Untransportable
KGW-83

618

Motobu / Okinawa (1993-06)
Unialgal, Clonal, H.Kobayashi (1993-06)
Identified by: Y.Fukuyo
Culture conditions: ESM, 20° C, 32 µE/m² sec, 1M
Characteristics: Marine, Untransportable
PX-01

Prorocentrum micans Ehrenberg

12

Osaka Bay / Osaka (1981-07)
Axenic, Clonal, S.Yamochi
Culture conditions: ESM, 20° C, 32 µE/m² sec, 1M
Characteristics: Red tide, Marine, Untransportable,
COXI gene (AB000133, AB000134)
OPm
References: 96, 97, 194, 361, 435

218

Yashima Bay / Kagawa (1978-08)
Axenic, Clonal, K.Yuki
Culture conditions: f/2, 20° C, 32 µE/m² sec, 1M
Characteristics: Red tide, Marine, Untransportable
KGW-13-7

316

Matoya Bay / Mie (1984-09)
Axenic, Clonal, T.Sawaguchi (1984-09)
Identified by: T.Sawaguchi
Culture conditions: f/2, ESM, 20° C, 32 µE/m² sec, 2M
Characteristics: Red tide, Marine, Untransportable
MB-D-4

601

Mikawa bay / Aichi
Unialgal, Clonal, S.Toriumi
Culture conditions: f/2, 20° C, 32 µE/m² sec, 1M
Characteristics: Marine, Untransportable

608

Ise Bay / Mie (1978-06)
Unialgal, Clonal, H.Iwasaki (1978-06)
Identified by: K.Steidnger
Culture conditions: ESM, 20° C, 40 µE/m² sec, 2M
Characteristics: Red tide, Marine, Untransportable

Prorocentrum minimum (Pavillard) Schiller

237

Osaka Bay / Osaka (1982-08)
Axenic, Clonal, M.M.Watanabe (1982-08)
Culture conditions: ESM, 20° C, 32 µE/m² sec, 1M
Characteristics: Red tide, Marine, Untransportable
OPmin

238

Harima-Nada / Seto Inland Sea (1983-04)
Unialgal, Clonal, S.Yoshimatsu
Culture conditions: f/2, ESM, 20° C, 32 µE/m² sec, 1M
Characteristics: Red tide, Marine, Unstable,
Untransportable
KGW-14-2-5

Prorocentrum sigmoides Bohm

683

Uchiumi Bay / Kagawa (1985-10)
Axenic, Clonal, S.Yoshimatsu (1985-10)
Identified by: S.Yoshimatsu
Culture conditions: ESM, 20° C, 40 µE/m² sec, 1M
Characteristics: Red tide, Marine, Untransportable

Prorocentrum triestinum Schiller

219

Nomi Bay / Kochi (1980-04)
Unialgal, Clonal, S.Yoshimatsu
Culture conditions: ESM, 20° C, 32 µE/m² sec, 1M
Characteristics: Red tide, Marine, Untransportable
KGW-28-1
Reference: 361

Protoceratium reticulatum

(Claparède et Lachmann) Bütschli

318

Matoya Bay / Mie (1984-09)
Axenic, Clonal, T.Sawaguchi (1984-09)

- Identified by: T.Sawaguchi
 Culture conditions: ESM, 20° C, 40 μ E/m²sec, 1M
 Characteristics: Red tide, Marine, Unstable,
 Untransportable
 MB-D-25
- 319
 Naoshima Isl. / Kagawa (1982-07)
 Axenic, Clonal, S.Yoshimatsu
 Identified by: S.Yoshimatsu
 Culture conditions: f/2, ESM, 20° C, 40 μ E/m²sec,
 1M
 Characteristics: Red tide, Marine, Unstable,
 Untransportable
 KGW-62
 Reference: 361
- * *Protogonyaulax catenella* (Whedon et Kofoid)
 Taylor
 See *Alexandrium catenella*
 (Whedon et Kofoid) Balech
- Pseudocarteria mucosa* (Korshikov) Ettl
 522
 Izumi / Miyagi (1985-08)
 Axenic, Clonal, S.Suda (1985-08)
 Identified by: S.Suda
 Culture conditions: AF-6, 20° C, 22 μ E/m²sec, 1M
 Characteristics: Freshwater, Homothallic
 M-2
 Reference: 329
- 523
 Higashiyata River / Ibaraki (1983-07)
 Unialgal, Clonal, S.Suda (1983-07)
 Identified by: S.Suda
 Culture conditions: AF-6, 20° C, 22 μ E/m²sec, 1M
 Characteristics: Freshwater, Homothallic
 USI-8
 References: 326, 329
- 524
 Izumi / Miyagi (1985-08)
 Axenic, Clonal, S.Suda (1985-08)
 Identified by: S.Suda
 Culture conditions: AF-6, 20° C, 22 μ E/m²sec, 1M
 Characteristics: Freshwater, Homothallic
 M-4
 Reference: 329
- Pseudopleurococcus printzii* Vischer
 var. *longissimus* S.Watanabe
- 159
 Kyoto (1975-03)
 Unialgal, Clonal, S.Watanabe (1975-03)
 Identified by: S.Watanabe
 Culture conditions: C(S), 20° C, 4 μ E/m²sec, 3M,
 (25° C, 30 μ E/m²sec)
 Characteristics: Indicator, Soil
 KUC6-2
 Reference: 416
- Pteromonas aculeata* Lemmermann
 738
 Shinobazu-no-ike, Ueno Park / Tokyo (1996-10)
 Unialgal, Clonal, S.Tanaka (1996-10)
 Identified by: S.Tanaka
 Culture conditions: AF-6, 20° C, 12 μ E/m²sec, 1M
 Characteristics: Freshwater
 970603-PtAcl
 Reference: 342
- Pteromonas angulosa* (Carter) Lemmermann
 739
 Shinobazu-no-ike, Ueno Park / Tokyo (1996-11)
 Axenic, Clonal, S.Tanaka (1996-11)
 Identified by: S.Tanaka
 Culture conditions: AF-6, 20° C, 12 μ E/m²sec, 1M
 Characteristics: Freshwater
 970603-PtAng
 Reference: 342
- Pteromonas multipyrenoidea* Iyenger
 740
 Shinobazu-no-ike, Ueno Park / Tokyo (1996-11)
 Axenic, Clonal, S.Tanaka (1996-11)
 Identified by: S.Tanaka
 Culture conditions: AF-6, 20° C, 12 μ E/m²sec, 1M
 Characteristics: Freshwater
 970603-PtMul
 Reference: 342
- Pterosperma cristatum* Schiller
 221
 Harima-Nada / Seto Inland Sea (1983-02)
 Axenic, Clonal, S.Suda (1983-09)
 Identified by: I.Inouye
 Culture conditions: f/2, ESM, 20° C, 32 μ E/m²sec,
 1M
 Characteristics: Red tide, Marine, Untransportable
 H-88-1
 References: 171, 361

- 626
Seto Inland Sea / Kagawa (1989-02)
Unialgal, Clonal, T.Sawaguchi (1989)
Identified by: I.Inouye
Culture conditions: ESM, 15° C, 20 µE/m²sec, 20D
Characteristics: Marine, Untransportable
89KGW-1
- Pyramimonas* aff. *amyliifera* Conrad
251
Yashima Bay / Kagawa (1982-10)
Axenic, Clonal, S.Yoshimatsu
Identified by: S.Suda
Culture conditions: f/2, ESM, 20° C, 32 µE/m²sec, 1M
Characteristics: Red tide, Marine
KGW-64-3
Reference: 361
- 320
Onagawa Bay / Miyagi (1984-08)
Axenic, Clonal, S.Suda (1984-09)
Identified by: S.Suda
Culture conditions: f/2, 20° C, 32 µE/m²sec, 1M
Characteristics: Red tide, Marine
8280G47-5
- Pyramimonas parkeae* Norris et Pearson
254
Hachijo Isl. / Tokyo (1984-04)
Axenic, Clonal, S.Suda (1984-04)
Identified by: S.Suda
Culture conditions: ESM, 20° C, 32 µE/m²sec, 1M
Characteristics: Indicator, Red tide, Marine, Tide pool, Collected from Senjo-jiki Yaene Hachijo 8-25-2
References: 135, 159, 160, 295
- Pyrocystis lunura* (Schütt) Schütt
609
Unialgal, Non-Clonal
Culture conditions: f/2, 20° C, 40 µE/m²sec, 1M
Characteristics: Marine
- Pyrophacus steinii* (Schiller) Wall et Dale
321
Matoya Bay / Mie (1984-09)
Unialgal, Clonal, T.Sawaguchi (1984-09)
Identified by: T.Sawaguchi
Culture conditions: ESM, 20° C, 40 µE/m²sec, 2M
Characteristics: Red tide, Marine, Untransportable
MB-D-27
- Rhodomonas atrorosea* Butcher ex Hill et Wetherbee
699
Isle of Wight / U.K.
CCAP 978/6a, Unialgal, Clonal, B.W.Butcher
Identified by: B.W.Butcher
Culture conditions: ESM, 15° C, 15 µE/m²sec, 1M
Characteristics: Marine, Type strain
Reference: 23
- Rhodomonas baltica* Karsten
700
Channel Islands / U.K.
CCAP 979/9, Unialgal, Clonal, B.W.Butcher (1961)
Identified by: B.W.Butcher
Culture conditions: ESM, 15° C, 15 µE/m²sec, 1M
Characteristics: Marine
Reference: 23
- Rhodomonas chrysoidea* Butcher ex Hill et Wetherbee
701
River Colne, Essex / U.K.
CCAP 979/8, Unialgal, Clonal, B.W.Butcher (1953)
Identified by: B.W.Butcher
Culture conditions: ESM, 15° C, 15 µE/m²sec, 1M
Characteristics: Brackish, Type strain
Reference: 23
- Rhodomonas duplex* Hill et Wetherbee
765
Yaka, Kin / Okinawa (1986)
Unialgal Clonal, I. Inouye (1986)
Identified by: M.Erata
Culture conditions: ESM, 15° C, 15 µE/m²sec, 1M
Characteristics: Marine
M014
- Rhodomonas falcata* Butcher ex Hill et Wetherbee
702
Aberystwyth, Wales / U.K.
CCAP 978/5a, Unialgal, Clonal, B.W.Butcher (1956)
Identified by: B.W.Butcher
Culture conditions: ESM, 15° C, 15 µE/m²sec, 1M
Characteristics: Marine, Type strain
Reference: 23
- Scenedesmus acuminatus* (Lageraeim) Chodat
var. *tetradesmoides* G.M.Smith
92
Lake Kasumigaura / Ibaraki (1983-08)
Axenic, Clonal, T.Hiwatari (1983-08)
Identified by: M.Watanabe
Culture conditions: CT, 20° C, 8 µE/m²sec, 2M

- Characteristics: Indicator, Freshwater
K-S-1
Reference: 417
- Scenedesmus acutus* Meyen
94
Kosaka River / Akita (1983-04)
Axenic, Clonal, A.Yuri (1983-05)
Identified by: M.Watanabe
Culture conditions: C(S), 20° C, 4 µE/m² sec, 3M,
(25° C, 30 µE/m² sec)
Characteristics: Indicator, Freshwater
2-2-3-1
Reference: 417
- 95
Tsukuba / Ibaraki (1983-05)
Axenic, Clonal, S.Suda (1983-05)
Identified by: M.Watanabe
Culture conditions: C(S), 20° C, 4 µE/m² sec, 3M,
(25° C, 30 µE/m² sec)
Characteristics: Indicator, Freshwater, Collected
from an artificial pond beside Aquatron at the
NIES
Aq-S-1
References: 70, 399
- 120
Tsukuba / Ibaraki (1983-05)
Axenic, Clonal, S.Suda (1983-05)
Identified by: M.Watanabe
Culture conditions: C(S), 20° C, 4 µE/m² sec, 3M,
(25° C, 30 µE/m² sec)
Characteristics: Indicator, Freshwater, Collected
from an artificial pond beside Aquatron at the
NIES
Aq-S-2
Reference: 399
- Scenedesmus dimorphus* (Turpin) Kützing
93
Lake Kasumigaura / Ibaraki (1983-07)
Axenic, Clonal, F.Kasai (1983-07)
Identified by: M.Watanabe
Culture conditions: C(S), 20° C, 4 µE/m² sec, 3M,
(25° C, 30 µE/m² sec)
Characteristics: Indicator, Freshwater
F-18-1
Reference: 399
- 119
Ozegahara / Gunma (1983-08)
Axenic, Clonal, S.Suda (1983-09)
Identified by: T.Hiwatari
Culture conditions: C, 20° C, 8 µE/m² sec, 2M
Characteristics: Indicator, Freshwater
OZ-29
- Scenedesmus gutwinskii* Chodat
var. *heterospina* Bodfoggözy
797
Lake Kasumigaura / Ibaraki (1992-06)
Unialgal, Clonal, F.Kasai (1992-07)
Identified by: T.Nakano
Culture conditions: C, 20° C, 4 µE/m² sec, 6M,
(25° C, 30 µE/m² sec)
Characteristics: Freshwater, Herbicide (simetryn)
susceptible
B8-7
Reference: 121
- 798
Lake Kasumigaura / Ibaraki (1992-06)
Unialgal, Clonal, F.Kasai (1992-07)
Identified by: T.Nakano
Culture conditions: C, 20° C, 4 µE/m² sec, 6M,
(25° C, 30 µE/m² sec)
Characteristics: Freshwater, Herbicide (simetryn)
susceptible
B8-16
Reference: 121
- 799
Lake Kasumigaura / Ibaraki (1992-06)
Unialgal, Clonal, F.Kasai (1992-07)
Identified by: T.Nakano
Culture conditions: C, 20° C, 4 µE/m² sec, 6M,
(25° C, 30 µE/m² sec)
Characteristics: Freshwater, Herbicide (simetryn)
susceptible
B8-23
Reference: 121
- 800
Lake Kasumigaura / Ibaraki (1992-06)
Unialgal, Clonal, F.Kasai (1992-07)
Identified by: T.Nakano
Culture conditions: C, 20° C, 4 µE/m² sec, 6M,
(25° C, 30 µE/m² sec)
Characteristics: Freshwater, Herbicide (simetryn)
tolerant

- B3-12
Reference: 121
- 801
Lake Kasumigaura / Ibaraki (1992-06)
Unialgal, Clonal, F.Kasai (1992-07)
Identified by: T.Nakano
Culture conditions: C, 20° C, 4 µE/m² sec, 6M,
(25° C, 30 µE/m² sec)
Characteristics: Freshwater, Herbicide (simetryn)
tolerant
B3-15
Reference: 121
- 802
Lake Kasumigaura / Ibaraki (1992-06)
Unialgal, Clonal, F.Kasai (1992-07)
Identified by: T.Nakano
Culture conditions: C, 20° C, 4 µE/m² sec, 6M,
(25° C, 30 µE/m² sec)
Characteristics: Freshwater, Herbicide (simetryn)
tolerant
B12-2
References: 120, 121
- Scenedesmus quadricauda*
(Turpin) Brébisson sensu Chodat
96
Lake Shoji / Yamanashi (1981-08)
TAC 51-3B, Axenic, Clonal, M.Watanabe
Identified by: M.Watanabe
Culture conditions: C, 20° C, 4 µE/m² sec, 3M,
(25° C, 30 µE/m² sec)
Characteristics: Indicator, Freshwater,
COXI gene (D63658, AB011524))
TAN-51-3B
References: 66, 374, 434
- Scenedesmus serratus* (Corda) Bohlin
97
Lake Shoji / Yamanashi (1981-08)
TAC 51-3C, Axenic, Clonal, M.Watanabe
Identified by: M.Watanabe
Culture conditions: C, 20° C, 4 µE/m² sec, 3M,
(25° C, 30 µE/m² sec)
Characteristics: Indicator, Freshwater
TAN-51-3C
- Schroederia setigera* (Schröder) Lemmermann
246
Lake Kasumigaura / Ibaraki (1983-08)
- Axenic, Clonal, F.Kasai (1983-08)
Identified by: M.Watanabe
Culture conditions: C, 25° C, 30 µE/m² sec, 20D
Characteristics: Indicator, Freshwater
F47-3
- Scrippsiella sweeneyae* Balech
684
Bisan-Seto / Seto Inland Sea (1982-07)
Unialgal, Clonal, S.Yoshimatsu (1982-07)
Identified by: S.Yoshimatsu
Culture conditions: ESM, 20° C, 40 µE/m² sec, 1M
Characteristics: Red tide, Marine, Untransportable
- Scrippsiella trochoidea* (Stein) Loeblich III
369
Hachinohe Harbor / Aomori (1985-08)
Axenic, Clonal, T.Sawaguchi (1985-08)
Identified by: T.Sawaguchi
Culture conditions: ESM, 20° C, 40 µE/m² sec, 1M
Characteristics: Red tide, Marine, Homothallic,
Unstable, Untransportable,
COXI gene (AB000135)
HHSS-1
References: 97, 194, 435
- Selenastrum capricornutum* Printz
Syn. *Monoraphidium capricornutum* (Printz) Nygaard
35
Nitelva River / Norway
Axenic, Clonal, O.M.Skulberg (1959)
Culture conditions: C(S), 20° C, 4 µE/m² sec, 3M,
(25° C, 30 µE/m² sec)
Characteristics: AGP, Freshwater
P-26
References: 59, 73, 116, 117, 122, 126, 131, 162,
197, 198, 218, 330, 352, 419, 420, 422
- Skeletonema costatum* (Greville) Cleve
16
Harima-Nada / Seto Inland Sea (1982-02)
Unialgal, Clonal, M.M.Watanabe (1982-05)
Identified by: M.M.Watanabe
Culture conditions: f/2, 5° C, 15 µE/m² sec, 1M
Characteristics: Red tide, Marine,
Collected from St. 53 Harima-Nada
H-53-3
References: 277, 300
- 17
Harima-Nada / Seto Inland Sea (1983-02)
Unialgal, Clonal, M.M.Watanabe (1983-05)

- Identified by: M.M.Watanabe
 Culture conditions: f/2, 5° C, 15 µE/m² sec, 1M
 Characteristics: Red tide, Marine,
 Collected from St. 90 Harima-Nada
 H-90-2
- 223
 Shodo Isl. / Kagawa (1979-07)
 Unialgal, Clonal, K.Yuki
 Culture conditions: f/2, 5° C, 15 µE/m² sec, 1M
 Characteristics: Red tide, Marine
 KGW-26
- 323
 Off Kishiwada / Osaka Bay (1985-01)
 Axenic, Clonal, S.Yamochi (1985-01)
 Identified by: S.Yamochi
 Culture conditions: f/2, 5° C, 15 µE/m² sec, 1M
 Characteristics: Red tide, Marine
 Sk-85w
 References: 108, 160
- 324
 Off Kobe / Osaka Bay (1985-07)
 Axenic, Clonal, S.Yamochi (1985-07)
 Identified by: S.Yamochi
 Culture conditions: f/2, 5° C, 15 µE/m² sec, 1M
 Characteristics: Red tide, Marine
 Sk-85su
 Reference: 265
- Spinoclosterium cuspidatum* (Bailey ex Ralfs) Hirano
 325
 Higashihiroshima / Hiroshima (1983-10)
 Unialgal, Clonal, T.Ichimura (1983-10)
 Identified by: T.Ichimura
 Culture conditions: SW(Bi), 20° C, 8 µE/m² sec, 4M,
 (25° C, 30 µE/m² sec)
 Characteristics: Freshwater, Homothallic
 83-24-19
 Reference: 83
- Spirulina platensis* (Gomont) Geitler
 Syn. *Arthrospira platensis* Gomont
 39
 Lake Chad / Chad
 IAM M-135, Axenic, Clonal
 Culture conditions: SOT, 20° C, 4 µE/m² sec, 4M,
 (25° C, 15 µE/m² sec)
 Characteristics: Salt water, Hydrogen evolution,
 Contains good quality of proteins
- References: 4, 80, 165, 177, 335, 387, 391, 399
- 45
 Lake Kasumigaura / Ibaraki (1975-11)
 IAM M-184, Unialgal, Clonal, M.M.Watanabe
 (1975-11)
 Identified by: M.M.Watanabe
 Culture conditions: MA, 25° C, 24 µE/m² sec, 1M,
 [Cryopreserved]
 Characteristics: Water bloom, Freshwater,
 Forming water bloom in Inbanuma
 KAS-6-50
 References: 80, 335, 387, 391, 399, 417
- 46
 Lake Texcoco / Mexico
 IAM M-185, Axenic, Clonal
 Culture conditions: SOT, 20° C, 4 µE/m² sec, 4M
 (25° C, 15 µE/m² sec)
 Characteristics: Water bloom, Salt water,
 Hydrogen evolution
 References: 2, 4, 7, 8, 80, 165, 335, 387, 391, 399
- 597
 Lake Teganuma / Chiba (1990-07)
 Unialgal, Non-clonal, T.Hagiwara (1990-07)
 Identified by: T.Hagiwara
 Culture conditions: MA, 20° C, 4 µE/m² sec, 2M,
 (25° C, 15 µE/m² sec), [Cryopreserved]
 Characteristics: Water bloom, Freshwater, Planktonic
 T-43
- Spirulina subsalsa* Oersted ex Gomont
 27
 IAM M-183, Axenic, Clonal
 Culture conditions: MA, 25° C, 24 µE/m² sec, 1M,
 [Cryopreserved]
 Characteristics: Freshwater
 References: 80, 169, 417
- 527
 Shikabe / Hokkaido (1976-04)
 IAM M-182, Unialgal, Clonal, M.M.Watanabe
 (1976-04)
 Identified by: M.M.Watanabe
 Culture conditions: f/2, 25° C, 24 µE/m² sec, 1M,
 [Cryopreserved]
 Characteristics: Indicator, Marine
 Reference: 80

- 598
Chiyoda-ku / Tokyo (1989-10)
Unialgal, Non-clonal, T.Hagiwara (1989-10)
Identified by: T.Hagiwara
Culture conditions: CB, 20° C, 4 $\mu\text{E}/\text{m}^2\text{sec}$, 2M,
(25° C, 15 $\mu\text{E}/\text{m}^2\text{sec}$)
Characteristics: Freshwater, Planktonic
KO-39
- Staurastrum dejectum* Brébisson ex Ralfs
224
Lake Yamanaka / Yamanashi (1981-10)
TAC 53-1, Axenic, Clonal, M.Watanabe
Identified by: M.Watanabe
Culture conditions: AF-6, 20° C, 8 $\mu\text{E}/\text{m}^2\text{sec}$, 2M,
(20° C, 25 $\mu\text{E}/\text{m}^2\text{sec}$)
Characteristics: Freshwater
TAN-53-1
- Staurastrum dorcidentiferum* W. et G.S.West
665
Lake Biwa / Shiga (1986-09)
Unialgal, Clonal, S.Ohara (1986-09)
Identified by: M.Nakanishi
Culture conditions: AF-6, 25° C, 30 $\mu\text{E}/\text{m}^2\text{sec}$, 2M
Characteristics: Freshwater
NB
- Staurastrum inconspicuum* Nordstedt
390
Oze / Gunma (1983-08)
Axenic, Clonal, F.Kasai (1983-09)
Culture conditions: CAM, 20° C, 8 $\mu\text{E}/\text{m}^2\text{sec}$, 3M,
(20° C, 12 $\mu\text{E}/\text{m}^2\text{sec}$)
Characteristics: Freshwater
34-10'
- Staurastrum levanderi* Grönblad
841
Namiki-ike Pond, Tsukuba / Ibaraki (1998-07)
Unialgal, Clonal, A.Gontcharov (1998-07)
Identified by: A.Gontcharov
Culture conditions: C, 20° C, 8 $\mu\text{E}/\text{m}^2\text{sec}$, 3M,
(25° C, 30 $\mu\text{E}/\text{m}^2\text{sec}$)
Characteristics: Freshwater
Reference: 48
- Staurastrum paradoxum* Meyen
528
Lake Kasumigaura / Ibaraki (1982-12)
Axenic, Clonal, M.H.Watanabe (1982-12)
Culture conditions: C, 20° C, 8 $\mu\text{E}/\text{m}^2\text{sec}$, 2M,
(20° C, 25 $\mu\text{E}/\text{m}^2\text{sec}$)
Characteristics: Indicator, Freshwater
Kas-K-3
- Staurastrum tsukubikum*
Gontcharov et M.M.Watanabe
842
Tsukuba / Ibaraki (1997-12)
Unialgal, Clonal, A.Gontcharov (1997-12)
Identified by: A.Gontcharov
Culture conditions: C, 20° C, 8 $\mu\text{E}/\text{m}^2\text{sec}$, 3M,
(25° C, 30 $\mu\text{E}/\text{m}^2\text{sec}$)
Characteristics: Freshwater, Type strain
Reference: 48
- Stephanopyxis palmeriana* (Greville) Grunow
327
Hachijo Isl. / Tokyo (1984-04)
Unialgal, Clonal, T.Sawaguchi (1984-04)
Identified by: T.Sawaguchi
Culture conditions: f/2, 10° C, 25 $\mu\text{E}/\text{m}^2\text{sec}$, 1M
Characteristics: Marine
8-B-2
- Stichococcus bacillaris* Nägeli
529
Watarase River / Gunma (1987-08)
Unialgal, Non-clonal, F.Kasai (1987-08)
Identified by: F.Kasai
Culture conditions: C, 15° C, 15 $\mu\text{E}/\text{m}^2\text{sec}$, 3M
Characteristics: Freshwater
AT2-16
Reference: 338
- 530
Watarase River / Gunma (1987-08)
Unialgal, Non-clonal, F.Kasai (1987-09)
Identified by: F.Kasai
Culture conditions: C, 15° C, 15 $\mu\text{E}/\text{m}^2\text{sec}$, 3M
Characteristics: Freshwater
AT5-17
References: 338, 339
- Stigeoclonium aestivale* (Hazen) Collins
531
Miyata River / Ibaraki (1987-03)
Unialgal, Non-clonal, F.Kasai (1987-04)
Identified by: F.Kasai
Culture conditions: C, 20° C, 8 $\mu\text{E}/\text{m}^2\text{sec}$, 3M
Characteristics: Freshwater

- 2st-3-12
References: 337, 338
- Stigeoclonium fasciculare* Kützing var. *fasciculare*
532
Lake Mashu / Hokkaido (1987-08)
Unialgal, Clonal, F.Kasai (1987-09)
Identified by: F.Kasai
Culture conditions: C, 10° C, 6 µE/m² sec, 3M,
(10° C, 15 µE / m² sec)
Characteristics: Freshwater
M-2
Reference: 338
- Synura petersenii* Korshikov
233
Higashiyata River / Ibaraki (1983-07)
Axenic, Clonal, S.Suda (1983-07)
Identified by: S.Suda
Culture conditions: C, 20° C, 12 µE/m² sec, 2M
Characteristics: Indicator, Freshwater
USI-10
Reference: 325
- Synura sphagnicola* (Korshikov) Korshikov
695
Miyatoko Mire / Fukushima (1992-04)
Axenic, Clonal, H.Nozaki (1992-04)
Identified by: H.Nozaki
Culture conditions: AF-6, 20° C, 12 µE/m² sec, 1M
Characteristics: Freshwater
92-520-s-6
696
Miyatoko Mire / Fukushima (1992-10)
Axenic, Clonal, H.Nozaki (1992-10)
Identified by: H.Nozaki
Culture conditions: AF-6, 20° C, 12 µE/m² sec, 1M
Characteristics: Freshwater
92-1001-s-2
- Synura spinosa* Korshikov
234
Tsuchiura / Ibaraki (1983-07)
Axenic, Clonal, S.Suda (1983-07)
Identified by: S.Suda
Culture conditions: C, 20° C, 12 µE/m² sec, 2M
Characteristics: Indicator, Freshwater
SIS-1
Reference: 325
- Tabellaria flocculosa* (Roth) Kützing
225
Oze / Fukushima (1983-08)
Unialgal, Clonal, M.M.Watanabe (1983-09)
Identified by: M.M.Watanabe
Culture conditions: CSi, 15° C, 20 µE/m² sec, 2M
Characteristics: Indicator, Freshwater
OZ-43-4
Reference: 277
- Tetrabaena socialis* (Dujardin) Nozaki et Ito
Syn. *Gonium sociale* (Dujardin) Warming
691
King George Isl. / Antarctic (1990-12)
Axenic, Clonal, S.Ohtani (1990-12)
Identified by: H.Nozaki
Culture conditions: AF-6, 10° C, 25 µE/m² sec, 1M
Characteristics: Freshwater, Psychrophilic
KG-4-8th
Reference: 256
- Tetrabaena socialis* (Dujardin) Nozaki et Ito
var. *socialis*
Syn. *Gonium sociale* (Dujardin) Warming var. *sociale*
571
Kohoku-ku / Yokohama / Kanagawa (1982-08)
Unialgal, Clonal, H.Nozaki (1982-10)
Identified by: H.Nozaki
Culture conditions: AF-6, 20° C, 12 µE/m² sec, 1M
Characteristics: Freshwater, Homothallic, Isogamy,
rbcL gene (D63443)
21028-4
References: 228, 242, 243, 256
- Tetracystis chlorococcoides* (Korshikov) S.Watanabe
155
Mt. Eboshidake / Nagasaki (1975-08)
Axenic, Clonal, S.Watanabe
Identified by: S.Watanabe
Culture conditions: C(S), 20° C, 4 µE/m² sec, 3M,
(25° C, 30 µE/m² sec)
Characteristics: Soil
3-EBO-1
Reference: 416
- Tetraëdron incus* (Teiling) G.M.Smith
392
Tsukuba / Ibaraki (1984-05)
Axenic, Clonal, F.Kasai (1984-05)
Identified by: M.Watanabe
Culture conditions: C, 20° C, 4 µE/m² sec, 3M,

- (25° C, 30 μ E/m² sec)
 Characteristics: Freshwater
 F115
 Reference: 325
- Tetraselmis cordiformis* (Carter) Stein
 18
 Oniishi / Gunma (1980-04)
 Axenic, Clonal, M.M.Watanabe (1980-04)
 Identified by: I.Inouye
 Culture conditions: C, 20° C, 32 μ E/m² sec, 1M
 Characteristics: Water bloom, Freshwater
 SM-6-9
 Reference: 399
- 533
 Mitsukaido / Ibaraki (1985-07)
 Axenic, Clonal, S.Suda (1985-07)
 Identified by: S.Suda
 Culture conditions: C, 20° C, 22 μ E/m² sec, 20D
 Characteristics: Freshwater
 KY-20-1
- Thalassionema nitzschioides* (Grunow) Hustedt
 534
 Matoya Bay / Mie (1984-09)
 Unialgal, Clonal, T.Sawaguchi (1984-09)
 Identified by: T.Sawaguchi
 Culture conditions: f/2, 15° C, 20 μ E/m² sec, 1M
 Characteristics: Marine
 MBB-6
 Reference: 277
- Thalassiosira pacifica* Gran et Angst
 535
 Hachinohe Harbor / Aomori (1987-03)
 Unialgal, Clonal, T.Sawaguchi (1987-03)
 Identified by: T.Sawaguchi
 Culture conditions: f/2, 10° C, 25 μ E/m² sec, 1M
 Characteristics: Marine
 87MHHB-1
- Tolypothrix tenuis* Kützing ex Bornet et Flahault
 37
 Borneo
 IAM M-29, Unialgal, Non-clonal, A.Watanabe
 Identified by: K.Negoro
 Culture conditions: MDM(S), 20° C, 4 μ E/m² sec,
 4M, (25° C, 30 μ E/m² sec), [Cryopreserved]
 Characteristics: Freshwater, Nitrogen fixation,
 Chromatic adaptation, Heterotrophic, Reidentified
- by M.M.Watanabe, Material for studying on
 phycobilin production
 References: 18, 30, 31, 32, 33, 34, 35, 37, 61, 62, 63,
 80, 141, 193, 298, 353, 360, 365, 366, 367, 368,
 369, 370, 371, 373, 436
- Treubaria triappendiculata* Bernard
 394
 Lake Kasumigaura / Ibaraki (1983-10)
 Axenic, Clonal, F.Kasai (1983-10)
 Identified by: Y.Niiyama
 Culture conditions: C, 20° C, 4 μ E/m² sec, 2M,
 (25° C, 30 μ E/m² sec)
 Characteristics: Freshwater
 F67-5
- * *Tribonema marinum* J.Feldmann
 See *Acinetospora crinita* (Carmichael) Sauvageau
- Triceratium dubium* Brightwell
 556
 Okinawa (1990)
 Unialgal, Clonal, S.Ono (1990)
 Identified by: S.Ono
 Culture conditions: f/2, 20° C, 40 μ E/m² sec, 1M
 Characteristics: Marine
 No.20
- Triploceras gracile* Bailey
 789
 2 km east of Melaka / Malaysia (1985-08)
 Unialgal, Clonal, T.Ichimura (1985)
 Identified by: T.Ichimura
 Culture conditions: MG, 20° C, 8 μ E/m² sec, 3M,
 (25° C, 30 μ E/m² sec)
 Characteristics: Freshwater, Homothallic
 85-28-1
 Reference: 85
- 790
 2 km east of Melaka / Malaysia (1985-08)
 Unialgal, Clonal, T.Ichimura (1985)
 Identified by: T.Ichimura
 Culture conditions: MG, 20° C, 8 μ E/m² sec, 3M,
 (25° C, 30 μ E/m² sec)
 Characteristics: Freshwater, Homothallic
 85-28-2
 Reference: 85
- 791
 2 km east of Melaka / Malaysia (1985-08)

- Unialgal, Clonal, T.Ichimura (1985)
 Identified by: T.Ichimura
 Culture conditions: MG, 20° C, 8 µE/m² sec, 3M,
 (25° C, 30 µE/m² sec)
 Characteristics: Freshwater, Homothallic
 85-28-3
 Reference: 85
- 792
 2 km east of Melaka / Malaysia (1985-08)
 Unialgal, Clonal, T.Ichimura (1985)
 Identified by: T.Ichimura
 Culture conditions: MG, 20° C, 8 µE/m² sec, 3M,
 (25° C, 30 µE/m² sec)
 Characteristics: Freshwater, Homothallic
 85-28-4
 Reference: 85
- 793
 Higashihiroshima / Hiroshima (1983-10)
 Unialgal, Clonal, T.Ichimura (1983)
 Identified by: T.Ichimura
 Culture conditions: MG, 20° C, 8 µE/m² sec, 3M,
 (25° C, 30 µE/m² sec)
 Characteristics: Freshwater, Heterothallic,
 Mating type +
 83-24-2
 Reference: 85
- 794
 Higashihiroshima / Hiroshima (1983-10)
 Unialgal, Clonal, T.Ichimura (1983)
 Identified by: T.Ichimura
 Culture conditions: MG, 20° C, 8 µE/m² sec, 3M,
 (25° C, 30 µE/m² sec)
 Characteristics: Freshwater, Heterothallic,
 Mating type +
 83-24-7
 Reference: 85
- 795
 Higashihiroshima / Hiroshima (1983-10)
 Unialgal, Clonal, T.Ichimura (1983)
 Identified by: T.Ichimura
 Culture conditions: MG, 20° C, 8 µE/m² sec, 3M,
 (25° C, 30 µE/m² sec)
 Characteristics: Freshwater, Heterothallic,
 Mating type –
 83-24-3
 Reference: 85
- 796
 Higashihiroshima / Hiroshima (1983-10)
 Unialgal, Clonal, T.Ichimura (1983)
 Identified by: T.Ichimura
 Culture conditions: MG, 20° C, 8 µE/m² sec, 3M,
 (25° C, 30 µE/m² sec)
 Characteristics: Freshwater, Heterothallic,
 Mating type –
 83-24-6
 Reference: 85
- Ulothrix variabilis* Kützing
 329
 Takatori River / Ibaraki (1984-12)
 Unialgal, Clonal, S.Suda (1984-12)
 Identified by: M.M.Watanabe
 Culture conditions: C, 20° C, 12 µE/m² sec, 3M
 Characteristics: Freshwater
 References: 337, 338
- Ulothrix zonata* (Weber et Mohr) Kützing
 536
 Hitachi / Ibaraki (1987-05)
 Unialgal, Non-clonal, F.Kasai (1987-06)
 Identified by: F.Kasai
 Culture conditions: C, 10° C, 6 µE/m² sec, 3M,
 (10° C, 15 µE/m² sec)
 Characteristics: Freshwater
 4st-1'-24
 Reference: 338
- 537
 Shirai River / Sapporo (1987-10)
 Unialgal, Non-clonal, F.Kasai (1987-10)
 Identified by: F.Kasai
 Culture conditions: C, 10° C, 15 µE/m² sec, 1M
 Characteristics: Freshwater
 2Tst-1-1
 Reference: 338
- Urnella terrestris* Playfair
 156
 Pokhara / Nepal (1975-10)
 Unialgal, Clonal, S.Watanabe
 Identified by: S.Watanabe
 Culture conditions: C(S), 20° C, 4 µE/m² sec, 3M,
 (25° C, 30 µE/m² sec)
 Characteristics: Soil
 NPL-111
 Reference: 415

Uroglena americana Calkins

395
Lake Biwa / Shiga (1978-05)
Unialgal, Clonal, Monoxenic, Y. Ishida (1978-05)
Identified by: Y. Ishida
Culture conditions: URO, 15° C, 20 µE/m² sec, 1M
Characteristics: Water bloom, Phagotrophic,
Freshwater, Untransportable
Strain 78
References: 99, 137, 138

Uronema confervicolum Lagerheim

538
Miyata River / Ibaraki (1987-05)
Unialgal, Non-clonal, F. Kasai (1987-05)
Identified by: F. Kasai
Culture conditions: C, 20° C, 8 µE/m² sec, 3M
Characteristics: Freshwater
4st-2-10
References: 337, 338

Uronema gigas Vischer

539
Miyata River / Ibaraki (1987-05)
Unialgal, Non-clonal, F. Kasai (1987-05)
Identified by: F. Kasai
Culture conditions: C, 20° C, 8 µE/m² sec, 3M
Characteristics: Freshwater
4st-3-5
Reference: 338

540

Miyata River / Ibaraki (1987-05)
Unialgal, Non-clonal, F. Kasai (1987-05)
Identified by: F. Kasai
Culture conditions: C, 20° C, 8 µE/m² sec, 3M
Characteristics: Freshwater
4st-0-16
Reference: 338

Volvox aureus Ehrenberg

241
Nagatoro / Saitama (1969-11)
IAM C-419, Axenic, Clonal, T. Ichimura
Identified by: T. Ichimura
Culture conditions: VT, 25° C, 30 µE/m² sec, 20D
Characteristics: Freshwater, Fertility lost,
Untransportable
S-9-8
Reference: 80

396

Koshokugun / Nagano (1983-08)
Axenic, Clonal, Y. Ogasawara (1983-08)
Identified by: Y. Ogasawara
Culture conditions: VT, 20° C, 12 µE/m² sec, 20D
Characteristics: Freshwater, Homothallic,
Untransportable

693

Meguro / Tokyo (1977-06)
Axenic, Clonal, H. Nozaki (1977-06)
Identified by: H. Nozaki
Culture conditions: VT, 20° C, 12 µE/m² sec, 1M
Characteristics: Freshwater, Water bloom,
Homothallic, Dioecious, Oogamy, Untransportable
k-5

694

Sakyo / Kyoto (1983-10)
Axenic, Clonal, H. Nozaki (1983-10)
Identified by: H. Nozaki
Culture conditions: VT, 20° C, 12 µE/m² sec, 1M
Characteristics: Freshwater, Water bloom,
Homothallic, Dioecious, Oogamy, Untransportable
31202-2-9

Volvox aureus Ehrenberg var. *aureus*

541
Lake Yamanaka / Yamanashi (1981)
Axenic, Clonal, H. Nozaki (1981-07)
Identified by: H. Nozaki
Culture conditions: VT, 20° C, 12 µE/m² sec, 1M
Characteristics: Freshwater, *rbcl* gene (D63445),
Untransportable
1706-2
References: 225, 238, 242, 243

542

Lake Yamanaka / Yamanashi (1981)
Axenic, Clonal, H. Nozaki (1981-07)
Identified by: H. Nozaki
Culture conditions: VT, 20° C, 12 µE/m² sec, 1M
Characteristics: Freshwater, Untransportable
1706-4
Reference: 225

Volvox barberi Shaw

730
California / USA (1965-09)
UTEX 804, Unialgal, Clonal, J. Stein (1957-04)
Culture conditions: AF-6, 20° C, 12 µE/m² sec, 1M
Characteristics: Freshwater, *rbcl* gene (D86835),

- Untransportable
Reference: 244
- Volvox carteri* Stein
397
Ichinomiya / Aichi (1983-06)
Axenic, Clonal, Y.Ogasawara (1983-06)
Culture conditions: VT, 25° C, 30 μ E/m² sec, 20D
Characteristics: Freshwater, Heterothallic, Female,
Crosses with NIES-398, Untransportable
V-4
- 398
Ichinomiya / Aichi (1983-06)
Axenic, Clonal, Y.Ogasawara (1983-06)
Culture conditions: VT, 25° C, 30 μ E/m² sec, 20D
Characteristics: Freshwater, Heterothallic, Male,
Crosses with NIES-397, Untransportable
V-11
- Volvox carteri* Stein f. *kawasakiensis* Nozaki
580
Kawasaki / Kanagawa (1984-01)
Unialgal, Clonal, H.Nozaki (1986-06)
Identified by: H.Nozaki
Culture conditions: VTAC, 20° C, 22 μ E/m² sec,
1M
Characteristics: Freshwater, Heterothallic,
Dioecious, Oogamy, Female, Crosses with
NIES-581, Untransportable
6823- ♀ -2
Reference: 231
- 581
Kawasaki / Kanagawa (1990-10)
Unialgal, Clonal, H.Nozaki (1990-11)
Identified by: H.Nozaki
Culture conditions: VTAC, 20° C, 22 μ E/m² sec, 1M
Characteristics: Freshwater, Heterothallic, Dioecious,
Oogamy, Male, Crosses with NIES-580,
Untransportable
90-1111-5
Reference: 231
- 732
Kawasaki / Kanagawa
Axenic, Clonal, H.Nozaki
Identified by: H.Nozaki
Culture conditions: AF-6, 20° C, 12 μ E/m² sec, 1M
Characteristics: Freshwater, Heterothallic, Dioecious,
Oogamy, Female, Crosses with NIES-733, Type
strain, *rbcL* gene (D63446), Untransportable
- KK-3
References: 231, 238, 243
- 733
Kawasaki / Kanagawa
Axenic, Clonal, H.Nozaki
Identified by: H.Nozaki
Culture conditions: AF-6, 20° C, 12 μ E/m² sec, 1M
Characteristics: Freshwater, Heterothallic, Dioecious,
Oogamy, Male, Crosses with NIES-732,
Untransportable
KK-5
Reference: 231
- Volvox dissipatrix* (Shaw) Printz
731
UTEX 2184, Unialgal, Clonal, R.C.Starr
Culture conditions: AF-6, 20° C, 12 μ E/m² sec, 1M
Characteristics: Freshwater, *rbcL* gene (D63447),
Unstable, Untransportable
Reference: 243
- Volvox prolificus* Iyengar
543
Axenic, Clonal, Y.Ogasawara
Identified by: S.Suda
Culture conditions: VT, 25° C, 30 μ E/m² sec, 1M
Characteristics: Freshwater, Untransportable
V-sp
- Volvox rousseletii* G.S.West
734
UTEX 1862, Unialgal, Clonal, R.C.Starr
Culture conditions: AF-6, 20° C, 12 μ E/m² sec, 1M
Characteristics: Freshwater, *rbcL* gene (D63448),
Untransportable
Reference: 243
- Volvox tertius* Meyer
544
Kisofukushima / Nagano (1986-08)
Axenic, Clonal, Y.Ogasawara (1986-08)
Identified by: Y.Ogasawara
Culture conditions: MG, 20° C, 12 μ E/m² sec, 20D
Characteristics: Freshwater, Homothallic,
Untransportable
- Volvulina compacta* Nozaki
582
Birtamod / Nepal (1988-10)
Axenic, Clonal, H.Nozaki (1989-08)
Identified by: H.Nozaki

- Culture conditions: VT, 20° C, 12 µE/m² sec, 1M
 Characteristics: Freshwater, Heterothallic,
 Mating type +, Crosses with NIES-583,
rbcL gene (D86832)
 89-804-4
 References 244, 251
- 583
 Birtamod / Nepal (1988-10)
 Axenic, Clonal, H.Nozaki (1989-08)
 Identified by: H.Nozaki
 Culture conditions: VT, 20° C, 12 µE/m² sec, 1M
 Characteristics: Freshwater, Heterothallic,
 Mating type –, Crosses with NIES-582
 89-804-7
 Reference: 251
- Volvulina steinii* Playfair
 545
 Hayama / Kanagawa (1980-12)
 Axenic, Clonal, H.Nozaki (1981-01)
 Identified by: H.Nozaki
 Culture conditions: VTAC, 20° C, 12 µE/m² sec,
 1M
 Characteristics: Freshwater, Heterothallic,
 Mating type –, Crosses with NIES-546
 1107-5 (–)
 References: 221, 241
- 546
 Hayama / Kanagawa (1980-12)
 Axenic, Clonal, H.Nozaki (1981-01)
 Identified by: H.Nozaki
 Culture conditions: VTAC, 20° C, 12 µE/m² sec, 1M
 Characteristics: Freshwater, Heterothallic,
 Mating type +, Crosses with NIES-545
 1107-8 (+)
 Reference: 221
- 584
 Bahrabise / Nepal (1988-09)
 Unialgal, Clonal, H.Nozaki (1989-03)
 Identified by: H.Nozaki
 Culture conditions: VTAC, 20° C, 22 µE/m² sec, 1M
 Characteristics: Freshwater, Heterothallic, Isogamy,
 Mating type +, Crosses with NIES-585
 89-306-1
 Reference: 233
- 585
 Bahrabise / Nepal (1988-09)
 Unialgal, Clonal, H.Nozaki (1989-04)
- Identified by: H.Nozaki
 Culture conditions: VTAC, 20° C, 22 µE/m² sec, 1M
 Characteristics: Freshwater, Heterothallic, Isogamy,
 Mating type –, Crosses with NIES-584
 89-423-1
 Reference: 233
- Woloszynskia leopoliense* (Woloszynska)Thompson
 619
 Mitsukaidou / Ibaraki (1985-04)
 Axenic, Clonal, T.Sawaguchi (1985-04)
 Identified by: T.Sawaguchi
 Culture conditions: MW1/5, 20° C, 40 µE/m² sec, 1M
 Characteristics: Freshwater, Homothallic,
 Untransportable
 KRYZ-3
- Yamagishiella unicocca* (Raybarn et Starr) Nozaki
 Syn. *Pandorina unicocca* Rayburn et Starr
 578
 Kamogawa / Chiba (1980-10)
 Unialgal, Clonal, H.Nozaki (1980-12)
 Identified by: H.Nozaki
 Culture conditions: VTAC, 20° C, 22 µE/m² sec, 1M
 Characteristics: Freshwater, Heterothallic, Isogamy,
 Mating type +, Crosses with NIES-579,
rbcL gene (AB000811)
 01209-1
 Reference: 257
- 579
 Kamogawa / Chiba (1980-10)
 Unialgal, Clonal, H.Nozaki (1980-12)
 Identified by: H.Nozaki
 Culture conditions: VTAC, 20° C, 22 µE/m² sec,
 1M
 Characteristics: Freshwater, Heterothallic, Isogamy,
 Mating type –, Crosses with NIES-578
 01209-7
- 666
 Nobi / Kanagawa (1979-05)
 UTEX 2428, Unialgal, Clonal, S.Kato (1979-05)
 Identified by: H.Nozaki
 Culture conditions: VT, 20° C, 22 µE/m² sec, 1M
 Characteristics: Freshwater, Heterothallic, Isogamy,
 Mating type +, Crosses with NIES-667,
rbcL gene (D86823)
 X-441
 References: 135, 220, 244, 254

667

Nobi / Kanagawa (1979-05)
UTEX 2429, Unialgal, Clonal, S.Kato (1979-05)
Identified by: H.Nozaki
Culture conditions: VT, 20° C, 22 μ E/m²sec, 1M
Characteristics: Freshwater, Heterothallic, Isogamy,
Mating type -, Crosses with NIES-666
X-443
Reference: 220

762

China
CCFA 646, Unialgal, Clonal
Reidentified by: H.Nozaki
Culture conditions: AF-6, 20° C, 12 μ E/m²sec, 1M
Characteristics: Freshwater, Formerly identified as
Eudorina sp., *rbcL* gene (AB000810)
Reference: 257

PROTOZOA

Paramecium bursaria Forke

668
Miyatoko Mire / Fukushima (1993-05)
Xenic, Clonal, H.Nozaki (1993-05)
Identified by: H.Nozaki
Culture conditions: AF-6, 20° C, 22 μ E/m²sec, 1M
Characteristics: Freshwater, Symbiotic
93-527-Pa-1

Tetrahymena pyriformis Ehrenberg

403
Tsuchiura Harbor / Lake Kasumigaura / Ibaraki
(1976-08)
Xenic, Non-clonal, R.Sudo (1976-08)
Identified by: R.Sudo
Culture conditions: LE, 10° C, 20D, (20° C)
Characteristics: Freshwater, Water bloom,
Untransportable
Tetra-1

IX. INDEXES

1. Numerical index

- | | | | |
|----|--|----|---|
| 1 | <i>Chattonella antiqua</i> | 47 | <i>Euglena gracilis</i> |
| 2 | <i>Chattonella antiqua</i> | 48 | <i>Euglena gracilis</i> |
| 3 | <i>Chattonella marina</i> | 49 | <i>Euglena gracilis</i> var. <i>bacillaris</i> |
| 4 | <i>Heterosigma akashiwo</i> | 50 | <i>Aulosira laxa</i> |
| 5 | <i>Heterosigma akashiwo</i> | 51 | <i>Closterium peracerosum-strigosum-littorale</i> complex |
| 6 | <i>Heterosigma akashiwo</i> | 52 | <i>Closterium peracerosum-strigosum-littorale</i> complex |
| 7 | <i>Heterocapsa triquetra</i> | 53 | <i>Closterium peracerosum-strigosum-littorale</i> complex |
| 8 | <i>Cricosphaera roscoffensis</i> | 54 | <i>Closterium peracerosum-strigosum-littorale</i> complex |
| 9 | <i>Heterosigma akashiwo</i> | 55 | <i>Closterium peracerosum-strigosum-littorale</i> complex |
| 10 | <i>Heterosigma akashiwo</i> | 56 | <i>Closterium peracerosum-strigosum-littorale</i> complex |
| 12 | <i>Prorocentrum micans</i> | 57 | <i>Closterium peracerosum-strigosum-littorale</i> complex |
| 14 | <i>Chattonella marina</i> | 58 | <i>Closterium peracerosum-strigosum-littorale</i> complex |
| 15 | <i>Olisthodiscus luteus</i> | 59 | <i>Closterium peracerosum-strigosum-littorale</i> complex |
| 16 | <i>Skeletonema costatum</i> | 60 | <i>Closterium peracerosum-strigosum-littorale</i> complex |
| 17 | <i>Skeletonema costatum</i> | 61 | <i>Closterium peracerosum-strigosum-littorale</i> complex |
| 18 | <i>Tetraselmis cordiformis</i> | 62 | <i>Closterium peracerosum-strigosum-littorale</i> complex |
| 19 | <i>Anabaena cylindrica</i> | 63 | <i>Closterium peracerosum-strigosum-littorale</i> complex |
| 21 | <i>Anabaenopsis circularis</i> | 64 | <i>Closterium peracerosum-strigosum-littorale</i> complex |
| 22 | <i>Calothrix brevissima</i> | 65 | <i>Closterium peracerosum-strigosum-littorale</i> complex |
| 23 | <i>Anabaena variabilis</i> | 66 | <i>Closterium peracerosum-strigosum-littorale</i> complex |
| 24 | <i>Nostoc commune</i> | 67 | <i>Closterium peracerosum-strigosum-littorale</i> complex |
| 25 | <i>Nostoc linckia</i> | 68 | <i>Closterium peracerosum-strigosum-littorale</i> complex |
| 26 | <i>Nostoc minutum</i> | 69 | <i>Closterium peracerosum-strigosum-littorale</i> complex |
| 27 | <i>Spirulina subsalsa</i> | | |
| 28 | <i>Nostoc linckia</i> var. <i>arvense</i> | | |
| 29 | <i>Nostoc minutum</i> | | |
| 30 | <i>Phormidium tenue</i> | | |
| 31 | <i>Oscillatoria laetevirens</i> | | |
| 32 | <i>Phormidium foveolarum</i> | | |
| 33 | <i>Oscillatoria tenuis</i> | | |
| 34 | <i>Phormidium foveolarum</i> | | |
| 35 | <i>Selenastrum capricornutum</i> | | |
| 36 | <i>Oscillatoria limnetica</i> | | |
| 37 | <i>Tolypothrix tenuis</i> | | |
| 38 | <i>Nostoc commune</i> | | |
| 39 | <i>Spirulina platensis</i> | | |
| 40 | <i>Anabaena affinis</i> | | |
| 41 | <i>Anabaena circinalis</i> | | |
| 42 | <i>Microcystis elabens</i> var. <i>minor</i> | | |
| 43 | <i>Microcystis holsatica</i> | | |
| 44 | <i>Microcystis aeruginosa</i> f. <i>aeruginosa</i> | | |
| 45 | <i>Spirulina platensis</i> | | |
| 46 | <i>Spirulina platensis</i> | | |

- 70 *Closterium peracerosum-strigosum-littorale* complex
71 *Achnanthes minutissima*
73 *Anabaena flos-aquae* f. *flos-aquae*
74 *Anabaena flos-aquae* f. *flos-aquae*
75 *Anabaena flos-aquae* f. *flos-aquae*
76 *Anabaena spiroides*
77 *Anabaena spiroides* f. *spiroides*
78 *Anabaena spiroides* f. *crassa*
79 *Anabaena spiroides* f. *spiroides*
80 *Anabaena solitaria* f. *solitaria*
81 *Aphanizomenon flos-aquae* f. *gracile*
83 *Chattonella antiqua*
84 *Chattonella antiqua*
85 *Chattonella antiqua*
86 *Chattonella antiqua*
87 *Microcystis aeruginosa* f. *aeruginosa*
88 *Microcystis aeruginosa* f. *aeruginosa*
89 *Microcystis aeruginosa* f. *aeruginosa*
90 *Microcystis aeruginosa* f. *aeruginosa*
91 *Microcystis aeruginosa* f. *aeruginosa*
92 *Scenedesmus acuminatus* var. *tetradesmoides*
93 *Scenedesmus dimorphus*
94 *Scenedesmus acutus*
95 *Scenedesmus acutus*
96 *Scenedesmus quadricauda*
97 *Scenedesmus serratus*
98 *Microcystis aeruginosa* f. *flos-aquae*
99 *Microcystis aeruginosa* f. *aeruginosa*
100 *Microcystis aeruginosa* f. *aeruginosa*
101 *Microcystis aeruginosa* f. *aeruginosa*
102 *Microcystis viridis*
103 *Microcystis viridis*
104 *Microcystis wesenbergii*
105 *Microcystis wesenbergii*
106 *Microcystis wesenbergii*
107 *Microcystis wesenbergii*
108 *Microcystis wesenbergii*
109 *Microcystis wesenbergii*
110 *Microcystis wesenbergii*
111 *Microcystis wesenbergii*
112 *Microcystis wesenbergii*
113 *Chattonella antiqua*
114 *Chattonella antiqua*
115 *Chattonella marina*
116 *Chattonella marina*
117 *Chattonella marina*
118 *Chattonella marina*
119 *Scenedesmus dimorphus*
120 *Scenedesmus acutus*
121 *Chattonella marina*
122 *Chlamydomonas pulsatilla*
123 *Chlorogonium fusiforme*
124 *Closterium acerosum*
125 *Closterium acerosum*
127 *Closterium acerosum*
128 *Closterium calosporum* var. *galiciense*
129 *Coelastrum astroideum*
130 *Coelastrum astroideum*
131 *Coelastrum proboscideum*
132 *Coelastrum reticulatum*
133 *Cosmarium contractum*
134 *Dimorphococcus lunatus*
135 *Dimorphococcus lunatus*
136 *Fibrocapsa japonica*
137 *Echinosphaeridium nordstedtii*
138 *Gonatozygon brebissonii*
139 *Gonatozygon brebissonii*
143 *Gyrodinium instriatum*
144 *Haematococcus lacustris*
145 *Heterosigma akashiwo*
146 *Heterosigma akashiwo*
147 *Hyalotheca dissiliens*
148 *Hyalotheca dissiliens*
149 *Hyalotheca dissiliens*
150 *Hyalotheca dissiliens*
151 *Micractinium pusillum*
152 *Micrasterias crux-melitensis*
153 *Chlorosarcinopsis delicata*
154 *Characium maximum*
155 *Tetracystis chlorococcoides*
156 *Urnella terrestris*
157 *Chlamydomonas monticola*
158 *Chlamydomonas augustae* var. *ellipsoidea*
159 *Pseudopleurococcus printzii* var. *longissimus*
160 *Chlorosarcinopsis caeca*
161 *Chattonella antiqua*
162 *Closterium calosporum* var. *galiciense*
163 *Closterium calosporum* var. *galiciense*
164 *Closterium calosporum* var. *galiciense*
165 *Closterium calosporum* var. *galiciense*
166 *Closterium calosporum* var. *galiciense*
167 *Closterium calosporum* var. *galiciense*
168 *Closterium calosporum* var. *galiciense*
169 *Closterium calosporum* var. *himalayense*
170 *Closterium calosporum* var. *himalayense*
171 *Closterium calosporum* var. *himalayense*
172 *Closterium moniliferum* var. *moniliferum*
173 *Closterium moniliferum* var. *moniliferum*
174 *Closterium moniliferum* var. *moniliferum*

- 175 *Closterium navicula*
176 *Closterium navicula*
177 *Closterium navicula*
178 *Closterium navicula*
179 *Closterium gracile*
180 *Closterium gracile*
181 *Closterium incurvum*
182 *Closterium moniliferum* var. *submoniliferum*
183 *Closterium moniliferum* var. *submoniliferum*
185 *Closterium pusillum* var. *maius*
186 *Closterium spinosporum* var. *crassum*
187 *Closterium spinosporum* var. *crassum*
188 *Closterium spinosporum* var. *malaysiense*
189 *Closterium spinosporum* var. *malaysiense*
191 *Closterium spinosporum* var. *ryukyuense*
192 *Closterium spinosporum* var. *ryukyuense*
193 *Closterium spinosporum* var. *ryukyuense*
194 *Closterium spinosporum* var. *spinosporum*
195 *Closterium spinosporum* var. *spinosporum*
196 *Closterium spinosporum* var. *spinosporum*
197 *Closterium spinosporum* var. *spinosporum*
198 *Closterium tumidum*
199 *Closterium venus*
200 *Closterium wallichii*
201 *Closterium wallichii*
202 *Closterium wallichii*
203 *Oedogonium obesum*
204 *Oscillatoria agardhii*
205 *Oscillatoria agardhii*
206 *Oscillatoria animalis*
207 *Oscillatoria raciborskii*
208 *Oscillatoria rosea*
209 *Pediastrum boryanum*
210 *Pediastrum duplex* var. *duplex*
211 *Pediastrum duplex* var. *gracillimum*
212 *Pediastrum duplex*
213 *Pediastrum duplex* var. *duplex*
214 *Pediastrum duplex* var. *gracillimum*
215 *Pediastrum simplex*
216 *Pediastrum tetras*
217 *Penium margaritaceum*
218 *Prorocentrum micans*
219 *Prorocentrum triestinum*
220 *Alexandrium catenella*
221 *Pterosperma cristatum*
223 *Skeletonema costatum*
224 *Staurastrum dejectum*
225 *Tabellaria flocculosa*
226 *Graesiella emersonii*
227 *Chlorella vulgaris* var. *vulgaris*
228 *Closterium ehrenbergii*
229 *Closterium ehrenbergii*
230 *Merismopedia tenuissima*
231 *Coelastrum morus*
232 *Polyedriopsis spinulosa*
233 *Synura petersenii*
234 *Synura spinosa*
235 *Heterocapsa triquetra*
237 *Prorocentrum minimum*
238 *Prorocentrum minimum*
241 *Volvox aureus*
242 *Pandorina morum*
243 *Pandorina morum*
244 *Coelastrum astroideum*
245 *Coelastrum reticulatum* var. *reticulatum*
246 *Schroederia setigera*
247 *Gonatozygon monotaenium*
248 *Cosmocladium constrictum*
250 *Galdieria sulphuraria*
251 *Pyramimonas* aff. *amylifera*
252 *Nephroselmis astigmatica*
253 *Euglena clara*
254 *Pyramimonas parkeae*
255 *Monomastix minuta*
256 *Monomastix minuta*
257 *Hafniomonas montana*
258 *Closterium aciculare* var. *subpronum*
259 *Closterium aciculare* var. *subpronum*
261 *Closterium peracerosum-strigosum-littorale* complex
262 *Closterium peracerosum-strigosum-littorale* complex
263 *Anabaena spiroides* f. *spiroides*
265 *Asterionella glacialis*
266 *Calothrix crustacea*
267 *Calothrix parasitica*
268 *Calothrix scopulorum*
271 *Closterium calosporum* var. *calosporum*
274 *Cryptomonas ovata*
275 *Cryptomonas ovata*
276 *Cryptomonas platyuris*
277 *Cryptomonas rostratiformis*
278 *Cryptomonas rostratiformis*
279 *Cryptomonas tetrapyrenoidosa*
280 *Cryptomonas tetrapyrenoidosa*
281 *Cryptomonas tetrapyrenoidosa*
282 *Cryptomonas tetrapyrenoidosa*
284 *Dinobryon divergens*
285 *Docidium undulatum* var. *undulatum*
286 *Euglena mutabilis*

- 287 *Gonatozygon monotaenium*
288 *Gonium viridistellatum*
289 *Gonium viridistellatum*
290 *Gonium viridistellatum*
293 *Heterosigma akashiwo*
294 *Hyalotheca dissiliens*
 var. *dissiliens* f. *tridentula*
295 *Hydrodictyon reticulatum*
296 *Mesostigma viride*
297 *Micrasterias foliacea* var. *foliacea*
298 *Microcystis aeruginosa* f. *aeruginosa*
299 *Microcystis aeruginosa* f. *aeruginosa*
300 *Pediastrum angulosum* var. *angulosum*
301 *Pediastrum boryanum*
302 *Pediastrum simplex*
303 *Penium margaritaceum*
304 *Peridinium willei*
305 *Phormidium ramosum*
306 *Pleurotaenium cylindricum* var. *stuhlmannii*
307 *Pleurotaenium ehrenbergii* var. *curtum*
308 *Pleurotaenium ehrenbergii* var. *curtum*
309 *Pleurotaenium ehrenbergii* var. *ehrenbergii*
310 *Pleurotaenium ehrenbergii* var. *ehrenbergii*
311 *Pleurotaenium ehrenbergii* var. *curtum*
312 *Pleurotaenium nodosum* var. *nodosum*
313 *Pleurotaenium ovatum*
315 *Prorocentrum gracile*
316 *Prorocentrum micans*
317 *Prorocentrum mexicanum*
318 *Protoceratium reticulatum*
319 *Protoceratium reticulatum*
320 *Pyramimonas* aff. *amylifera*
321 *Pyrophacus steinii*
323 *Skeletonema costatum*
324 *Skeletonema costatum*
325 *Spinoclosterium cuspidatum*
327 *Stephanopyxis palmeriana*
329 *Ulothrix variabilis*
330 *Achnanthes longipes*
331 *Amphidinium carterae*
333 *Melosira granulata*
 var. *angustissima* f. *spiralis*
334 *Calothrix parasitica*
336 *Closterium calosporum* var. *himalayense*
337 *Closterium incurvum*
338 *Closterium rostratum* var. *subrostratum*
339 *Closterium selenastrum*
340 *Closterium selenastrum*
341 *Closterium spinosporum* var. *crassum*
342 *Coelastrum astroideum*
343 *Coolia monotis*
344 *Cryptomonas platyuris*
345 *Cryptomonas rostratiformis*
346 *Cryptomonas tetrapyrenoidosa*
347 *Cryptomonas tetrapyrenoidosa*
348 *Cryptomonas tetrapyrenoidosa*
349 *Cylindrocystis brebissonii* var. *brebissonii*
350 *Ditylum brightwellii*
351 *Eudorina elegans*
353 *Gephyrocapsa oceanica*
356 *Katodinium rotundatum*
359 *Oltmannsiellopsis unicellularis*
360 *Oltmannsiellopsis viridis*
361 *Oscillatoria amphibia*
362 *Pandorina morum*
363 *Pedinomonas minor*
364 *Peridinium bipes* var. *occultatum*
365 *Peridinium volzii*
366 *Peridinium willei*
369 *Scrippsiella trochoidea*
372 *Achnanthes minutissima* var. *saprophila*
375 *Brachiomonas submarina*
376 *Ceratium hirundinella*
377 *Chaetoceros sociale*
378 *Dictyochloropsis irregularis*
379 *Eremosphaera gigas*
380 *Eremosphaera viridis*
381 *Eutreptiella gymnastica*
382 *Lagerheimia ciliata*
384 *Monoraphidium contortum*
385 *Monoraphidium griffithii*
387 *Phacus agilis*
388 *Phaeocystis pouchetii*
390 *Staurastrum inconspicuum*
391 *Fragilaria capucina*
392 *Tetraëdron incus*
394 *Treubaria triappendiculata*
395 *Uroglena americana*
396 *Volvox aureus*
397 *Volvox carteri*
398 *Volvox carteri*
403 *Tetrahymena pyriformis*
405 *Amphidinium britannicum*
407 *Achnanthes minutissima*
408 *Achnanthes minutissima*
409 *Achnanthes minutissima*
410 *Achnanthes minutissima*
411 *Achnanthes minutissima*
412 *Achnanthes minutissima*
413 *Achnanthes minutissima*

- 414 *Achnanthes minutissima*
415 *Actinastrum hantzschii*
416 *Aphanocapsa montana*
417 *Asterionella glacialis*
418 *Astrephomene gubernaculifera*
419 *Astrephomene gubernaculifera*
420 *Cachonina niei*
421 *Carteria crucifera*
422 *Carteria inversa*
423 *Carteria inversa*
424 *Carteria cerasiformis*
425 *Carteria cerasiformis*
426 *Carteria klebsii*
427 *Carteria multifilis*
428 *Carteria obtusa*
429 *Carteria obtusa*
430 *Carteria obtusa*
431 *Carteria obtusa*
432 *Carteria radiosa*
433 *Chamaesiphon polymorphus*
434 *Chamaesiphon subglobosus*
436 *Characium polymorphum*
437 *Chlamydomonas fasciata*
438 *Chlamydomonas monadina* var. *monadina*
439 *Chlorogonium neglectum*
440 *Chlamydomonas parkeae*
441 *Chlamydomonas parkeae*
446 *Chlamydomonas tetragama*
447 *Chloromonas insignis*
448 *Closterium acerosum*
449 *Closterium pleurodermatum*
450 *Closterium praelongum* var. *brevius*
451 *Closterium praelongum* var. *brevius*
452 *Cosmarium hians*
453 *Dictyosphaerium pulchellum*
454 *Draparnaldia plumosa*
455 *Errerella bornhemiensis*
456 *Eudorina elegans* var. *elegans*
457 *Eudorina elegans* var. *elegans*
458 *Eudorina elegans* var. *synoica*
459 *Eudorina illinoisensis*
460 *Eudorina illinoisensis*
461 *Eunotia pectinalis* var. *minor*
462 *Fibrocapsa japonica*
463 *Glenodiniopsis uliginosa*
464 *Gloeomonas lateperforata*
465 *Gomphonema gracile* var. *gracile*
466 *Gomphonema parvulum* var. *parvulum*
467 *Gomphonema parvulum* var. *parvulum*
468 *Gonium pectorale* var. *pectorale*
469 *Gonium pectorale* var. *pectorale*
470 *Gymnodinium fuscum*
471 *Hemidinium nasutum*
472 *Heterocapsa pygmaea*
473 *Heterocapsa pygmaea*
474 *Lobomonas monstruosa*
475 *Mesostigma viride*
476 *Mesostigma viride*
477 *Mesostigma viride*
478 *Microcystis aeruginosa* f. *flos-aquae*
479 *Microthamnion kützingianum*
480 *Monoraphidium circinale*
481 *Myxosarsina burmensis*
483 *Nephroselmis olivacea*
484 *Nephroselmis olivacea*
485 *Nephroselmis olivacea*
486 *Nephroselmis viridis*
487 *Nitzschia palea*
488 *Nitzschia palea*
489 *Nitzschia palea*
494 *Oxyrrhis marina*
495 *Peridinium bipes* f. *globosum*
496 *Peridinium bipes* f. *occultatum*
497 *Peridinium bipes* f. *occultatum*
499 *Peridinium inconspicuum* subsp. *remotum*
500 *Peridinium polonicum*
501 *Peridinium volzii*
502 *Peridinium wierzejskii*
503 *Phormidium foveolarum*
504 *Phormidium foveolarum*
505 *Phormidium foveolarum*
506 *Phormidium jenkelianum*
507 *Phormidium jenkelianum*
509 *Phormidium molle*
510 *Phormidium mucicola*
512 *Phormidium tenue*
514 *Planctonema lauterbornii*
515 *Plectonema radiosum*
519 *Alexandrium catenella*
520 *Alexandrium catenella*
522 *Pseudocarteria mucosa*
523 *Pseudocarteria mucosa*
524 *Pseudocarteria mucosa*
527 *Spirulina subsalsa*
528 *Staurastrum paradoxum*
529 *Stichococcus bacillaris*
530 *Stichococcus bacillaris*
531 *Stigeoclonium aestivale*
532 *Stigeoclonium fasciculare* var. *fasciculare*
533 *Tetraselmis cordiformis*

- 534 *Thalassionema nitzschioides*
535 *Thalassiosira pacifica*
536 *Ulothrix zonata*
537 *Ulothrix zonata*
538 *Uronema confervicolum*
539 *Uronema gigas*
540 *Uronema gigas*
541 *Volvox aureus* var. *aureus*
542 *Volvox aureus* var. *aureus*
543 *Volvox prolificus*
544 *Volvox tertius*
545 *Volvulina steinii*
546 *Volvulina steinii*
547 *Cyanophora paradoxa*
548 *Acinetospora crinita*
549 *Cyanidioschyzon merolae*
550 *Galdieria sulphuraria*
551 *Cyanidium caldarium*
553 *Chaetoceros sociale*
556 *Triceratium dubium*
557 *Chattonella antiqua*
558 *Chattonella antiqua*
559 *Chattonella marina*
560 *Fibrocapsa japonica*
561 *Heterosigma akashiwo*
562 *Chrysochromulina parva*
564 *Astrephomene perforata*
565 *Astrephomene perforata*
566 *Basichlamys sacculifera*
567 *Characiochloris sasae*
568 *Eudorina elegans* var. *synoica*
569 *Gonium pectorale* var. *pectorale*
570 *Gonium pectorale* var. *pectorale*
571 *Tetrabaena socialis* var. *socialis*
572 *Pandorina colemaniae*
573 *Pandorina colemaniae*
574 *Pandorina morum* var. *morum*
575 *Pandorina morum* var. *morum*
576 *Pleodorina californica*
577 *Pleodorina japonica*
578 *Yamagishiella unicocca*
579 *Yamagishiella unicocca*
580 *Volvox carteri* f. *kawasakiensis*
581 *Volvox carteri* f. *kawasakiensis*
582 *Volvulina compacta*
583 *Volvulina compacta*
584 *Volvulina steinii*
585 *Volvulina steinii*
586 *Chaetoceros didymus*
587 *Hantzschia amphioxys* var. *compacta*
588 *Lithodesmium variabile*
589 *Odontella aurita*
590 *Odontella longicuris*
592 *Fischerella major*
593 *Hydrococcus rivularis*
594 *Oscillatoria agardhii*
595 *Oscillatoria agardhii*
596 *Oscillatoria agardhii*
597 *Spirulina platensis*
598 *Spirulina subsalsa*
600 *Peridinium bipes* var. *tabulatum*
601 *Prorocentrum micans*
603 *Chattonella ovata*
604 *Mycrocystis wesenbergii*
605 *Fibrocapsa japonica*
608 *Prorocentrum micans*
609 *Pyrocystis lunura*
610 *Oscillatoria rubescens*
611 *Phormidium tenue*
612 *Alexandrium hiranoi*
613 *Amphidinium klebsii*
614 *Cachonina niei*
615 *Coolia monotis*
617 *Prorocentrum lima*
618 *Prorocentrum mexicanum*
619 *Woloszynskia leopoliense*
620 *Gomphonema angustatum* var. *obtusatum*
621 *Botrydiopsis arrhiza*
622 *Botrydium granulatum*
623 *Pavlova gyrans*
624 *Chlorarachnion reptans*
626 *Pterosperma cristatum*
628 *Astrephomene gubernaculifera*
629 *Auxenochlorella protothecoides*
630 *Carteria crucifera*
631 *Carteria eugametos*
632 *Carteria eugametos*
633 *Carteria eugametos*
634 *Carteria eugametos*
635 *Carteria eugametos*
636 *Carteria eugametos*
637 *Characiochloris acuminata*
638 *Characiochloris sasae*
639 *Characium angustum*
640 *Chlorella saccharophila*
641 *Chlorella vulgaris* var. *vulgaris*
642 *Chlorella vulgaris* var. *vulgaris*
643 *Eremosphaera viridis*
644 *Eremosphaera viridis*
645 *Gonium pectorale* var. *pectorale*

- 646 *Gonium pectorale* var. *pectorale*
647 *Gonium quadratum*
648 *Gonium quadratum*
649 *Gonium quadratum*
650 *Gonium quadratum*
651 *Gonium quadratum*
652 *Gonium quadratum*
653 *Gonium quadratum*
654 *Gonium viridistellatum*
655 *Gonium viridistellatum*
656 *Hafniomonas montana*
657 *Mesotaenium kramstae*
658 *Mesotaenium kramstae*
659 *Oocystis borgei*
660 *Oocystis lacustris*
661 *Oocystis lacustris*
662 *Oocystis lacustris*
663 *Pleurotaenium nodosum* var. *borgei*
664 *Pleurotaenium nodosum* var. *borgei*
665 *Staurastrum dorcidentiferum*
666 *Yamagishiella unicocca*
667 *Yamagishiella unicocca*
668 *Paramecium bursaria*
670 *Chattonella verruculosa*
671 *Chattonella ovata*
672 *Oltmannsiellopsis geminata*
673 *Alexandrium affine*
674 *Alexandrium catenella*
675 *Alexandrium catenella*
677 *Alexandrium catenella*
678 *Alexandrium insuetum*
680 *Gymnodinium mikimotoi*
682 *Prorocentrum dentatum*
683 *Prorocentrum sigmoides*
684 *Scrippsiella sweeneyae*
685 *Chlorella fusca* var. *fusca*
686 *Chlorella vulgaris* var. *vulgaris*
687 *Graesiella emersonii*
688 *Graesiella emersonii*
689 *Graesiella emersonii*
690 *Graesiella emersonii*
691 *Tetrabaena socialis*
692 *Chlorogonium capillatum*
693 *Volvox aureus*
694 *Volvox aureus*
695 *Synura sphagnicola*
696 *Synura sphagnicola*
697 *Cryptomonas acuta*
698 *Cryptomonas irregularis*
699 *Rhodomonas atrorosea*
700 *Rhodomonas baltica*
701 *Rhodomonas chrysoidea*
702 *Rhodomonas falcata*
703 *Chroomonas collegionis*
704 *Chroomonas dispersa*
705 *Chroomonas placoidea*
706 *Chroomonas nordstedtii*
707 *Chroomonas nordstedtii*
708 *Chroomonas nordstedtii*
709 *Chroomonas nordstedtii*
710 *Chroomonas nordstedtii*
711 *Chroomonas nordstedtii*
712 *Chroomonas caudata*
713 *Chroomonas coerulea*
714 *Chroomonas coerulea*
715 *Chilomonas paramecium*
716 *Haramonas dimorpha*
717 *Eudorina elegans* var. *elegans*
718 *Eudorina elegans* var. *elegans*
719 *Eudorina elegans* var. *elegans*
720 *Eudorina elegans* var. *elegans*
721 *Eudorina elegans* var. *carteri*
722 *Eudorina cylindrica*
723 *Eudorina illinoisensis*
724 *Eudorina unicocca* var. *unicocca*
725 *Eudorina unicocca* var. *unicocca*
726 *Eudorina unicocca* var. *peripheralis*
727 *Paulschulzia pseudovolvox*
728 *Platydorina caudata*
729 *Platydorina caudata*
730 *Volvox barberi*
731 *Volvox dissipatrix*
732 *Volvox carteri* f. *kawasakiensis*
733 *Volvox carteri* f. *kawasakiensis*
734 *Volvox rousseletii*
735 *Pleodorina californica*
736 *Pleodorina indica*
737 *Gonium multicocum*
738 *Pteromonas aculeata*
739 *Pteromonas angulosa*
740 *Pteromonas multiplyrenoidea*
741 *Chrysochromulina hirta*
742 *Chlorogonium capillatum*
743 *Chlorogonium capillatum*
744 *Chlorogonium capillatum*
745 *Chlorogonium capillatum*
746 *Chlorogonium capillatum*
747 *Chlorogonium capillatum*
748 *Chlorogonium capillatum*
749 *Chlorogonium capillatum*

- 750 *Chlorogonium capillatum*
751 *Chlorogonium elongatum*
752 *Chlorogonium elongatum*
753 *Chlorogonium elongatum*
754 *Chlorogonium euchlorum*
755 *Chlorogonium euchlorum*
756 *Chlorogonium euchlorum*
757 *Chlorogonium euchlorum*
758 *Chlorogonium euchlorum*
759 *Chlorogonium euchlorum*
760 *Chlorogonium euchlorum*
761 *Chlorogonium kasakii*
762 *Yamagishiella unicocca*
763 *Cyanophora paradoxa*
764 *Cyanophora tetracyanea*
765 *Rhodomonas duplex*
766 *Chilomonas paramecium*
767 *Chilomonas paramecium*
768 *Cosmarium askenasyi*
769 *Cosmarium askenasyi*
770 *Cosmarium askenasyi*
771 *Cosmarium askenasyi*
772 *Euastrum turgidum*
773 *Euastrum turgidum*
774 *Micrasterias anomala*
775 *Micrasterias anomala*
776 *Micrasterias anomala*
777 *Micrasterias foliacea*
778 *Micrasterias foliacea*
779 *Micrasterias mahabuleshwarensis*
780 *Micrasterias mahabuleshwarensis*
781 *Micrasterias thomasiana* var. *notata*
782 *Micrasterias thomasiana* var. *notata*
783 *Micrasterias truncata* var. *pusilla*
784 *Micrasterias truncata* var. *pusilla*
785 *Pleurotaenium nodosum* var. *nodosum*
786 *Pleurotaenium nodosum* var. *nodosum*
787 *Pleurotaenium nodosum* var. *gutwinskii*
788 *Pleurotaenium nodosum* var. *gutwinskii*
789 *Triploceras gracile*
790 *Triploceras gracile*
791 *Triploceras gracile*
792 *Triploceras gracile*
793 *Triploceras gracile*
794 *Triploceras gracile*
795 *Triploceras gracile*
796 *Triploceras gracile*
797 *Scenedesmus gutwinskii* var. *heterospina*
798 *Scenedesmus gutwinskii* var. *heterospina*
799 *Scenedesmus gutwinskii* var. *heterospina*
800 *Scenedesmus gutwinskii* var. *heterospina*
801 *Scenedesmus gutwinskii* var. *heterospina*
802 *Scenedesmus gutwinskii* var. *heterospina*
803 *Cyclotella meneghiniana*
804 *Cyclotella meneghiniana*
805 *Cyclotella meneghiniana*
806 *Anabaena compacta*
807 *Anabaena kisseleviana*
808 *Anabaena mendotae*
809 *Anabaena mucosa*
810 *Anabaena planktonica*
811 *Anabaena planktonica*
812 *Anabaena planktonica*
813 *Anabaena planktonica*
814 *Anabaena planktonica*
815 *Anabaena planktonica*
816 *Anabaena planktonica*
817 *Anabaena planktonica*
818 *Anabaena smithii*
819 *Anabaena smithii*
820 *Anabaena smithii*
821 *Anabaena smithii*
822 *Anabaena smithii*
823 *Anabaena smithii*
824 *Anabaena smithii*
825 *Anabaena ucrainica*
826 *Anabaena ucrainica*
827 *Anabaena viguieri*
828 *Anabaena ellipsoides*
829 *Anabaena oumiana*
830 *Anabaena smithii*
831 *Anabaena smithii*
832 *Anabaena ucrainica*
833 *Anabaena lemmermannii*
834 *Anabaena planktonica*
835 *Anabaena compacta*
836 *Botryococcus braunii*
837 *Emiliana huxleyi*
838 *Gephyrocapsa oceanica*
839 *Cosmarium dilatatum*
840 *Euastrum biverrucosum*
841 *Staurastrum levanderi*
842 *Staurastrum tsukubicum*

2. Systematic index

ALGAE

CYANOPHYTA

Cyanophyceae

<i>Anabaena affinis</i>	40	<i>Anabaena spiroides</i> f. <i>crassa</i>	78
<i>Anabaena circinalis</i>	41	<i>Anabaena spiroides</i> f. <i>spiroides</i>	77
<i>Anabaena compacta</i>	806	<i>Anabaena spiroides</i> f. <i>spiroides</i>	79
<i>Anabaena compacta</i>	835	<i>Anabaena spiroides</i> f. <i>spiroides</i>	263
<i>Anabaena cylindrica</i>	19	<i>Anabaena variabilis</i>	23
<i>Anabaena ellipsoides</i>	828	<i>Anabaenopsis circularis</i>	21
<i>Anabaena flos-aquae</i> f. <i>flos-aquae</i>	73	<i>Aphanizomenon flos-aquae</i> f. <i>gracile</i>	81
<i>Anabaena flos-aquae</i> f. <i>flos-aquae</i>	74	<i>Aphanocapsa montana</i>	416
<i>Anabaena flos-aquae</i> f. <i>flos-aquae</i>	75	<i>Aulosira laxa</i>	50
<i>Anabaena kisseleviana</i>	807	<i>Calothrix brevissima</i>	22
<i>Anabaena lemmermannii</i>	833	<i>Calothrix crustacea</i>	266
<i>Anabaena mendotae</i>	808	<i>Calothrix parasitica</i>	267
<i>Anabaena mucosa</i>	809	<i>Calothrix parasitica</i>	334
<i>Anabaena oumiana</i>	829	<i>Calothrix scopulorum</i>	268
<i>Anabaena planktonica</i>	810	<i>Chamaesiphon polymorphus</i>	433
<i>Anabaena planktonica</i>	811	<i>Chamaesiphon subglobosus</i>	434
<i>Anabaena planktonica</i>	812	<i>Fischerella major</i>	592
<i>Anabaena planktonica</i>	813	<i>Hydrococcus rivularis</i>	593
<i>Anabaena planktonica</i>	814	<i>Merismopedia tenuissima</i>	230
<i>Anabaena planktonica</i>	815	<i>Microcystis aeruginosa</i> f. <i>aeruginosa</i>	44
<i>Anabaena planktonica</i>	816	<i>Microcystis aeruginosa</i> f. <i>aeruginosa</i>	87
<i>Anabaena planktonica</i>	817	<i>Microcystis aeruginosa</i> f. <i>aeruginosa</i>	88
<i>Anabaena planktonica</i>	834	<i>Microcystis aeruginosa</i> f. <i>aeruginosa</i>	89
<i>Anabaena smithii</i>	818	<i>Microcystis aeruginosa</i> f. <i>aeruginosa</i>	90
<i>Anabaena smithii</i>	819	<i>Microcystis aeruginosa</i> f. <i>aeruginosa</i>	91
<i>Anabaena smithii</i>	820	<i>Microcystis aeruginosa</i> f. <i>aeruginosa</i>	99
<i>Anabaena smithii</i>	821	<i>Microcystis aeruginosa</i> f. <i>aeruginosa</i>	100
<i>Anabaena smithii</i>	822	<i>Microcystis aeruginosa</i> f. <i>aeruginosa</i>	101
<i>Anabaena smithii</i>	823	<i>Microcystis aeruginosa</i> f. <i>aeruginosa</i>	101
<i>Anabaena smithii</i>	824	<i>Microcystis aeruginosa</i> f. <i>aeruginosa</i>	101
<i>Anabaena smithii</i>	830	<i>Microcystis aeruginosa</i> f. <i>aeruginosa</i>	101
<i>Anabaena smithii</i>	831	<i>Microcystis aeruginosa</i> f. <i>flos-aquae</i>	98
<i>Anabaena ucrainica</i>	825	<i>Microcystis aeruginosa</i> f. <i>flos-aquae</i>	478
<i>Anabaena ucrainica</i>	826	<i>Microcystis elabens</i> var. <i>minor</i>	42
<i>Anabaena ucrainica</i>	832	<i>Microcystis holsatica</i>	43
<i>Anabaena viguieri</i>	827	<i>Microcystis viridis</i>	102
<i>Anabaena solitaria</i> f. <i>solitaria</i>	80	<i>Microcystis viridis</i>	103
<i>Anabaena spiroides</i>	76	<i>Microcystis wesenbergii</i>	104
		<i>Microcystis wesenbergii</i>	105
		<i>Microcystis wesenbergii</i>	106
		<i>Microcystis wesenbergii</i>	107
		<i>Microcystis wesenbergii</i>	108
		<i>Microcystis wesenbergii</i>	109
		<i>Microcystis wesenbergii</i>	110
		<i>Microcystis wesenbergii</i>	111

<i>Microcystis wesenbergii</i>	112
<i>Microcystis wesenbergii</i>	604
<i>Myxosarcina burmensis</i>	481
<i>Nostoc commune</i>	24
<i>Nostoc commune</i>	38
<i>Nostoc linckia</i>	25
<i>Nostoc linckia</i> var. <i>arvense</i>	28
<i>Nostoc minutum</i>	26
<i>Nostoc minutum</i>	29
<i>Oscillatoria agardhii</i>	204
<i>Oscillatoria agardhii</i>	205
<i>Oscillatoria agardhii</i>	594
<i>Oscillatoria agardhii</i>	595
<i>Oscillatoria agardhii</i>	596
<i>Oscillatoria amphibia</i>	361
<i>Oscillatoria animalis</i>	206
<i>Oscillatoria laetevirens</i>	31
<i>Oscillatoria limnetica</i>	36
<i>Oscillatoria raciborskii</i>	207
<i>Oscillatoria rosea</i>	208
<i>Oscillatoria rubescens</i>	610
<i>Oscillatoria tenuis</i>	33
<i>Phormidium foveolarum</i>	32
<i>Phormidium foveolarum</i>	34
<i>Phormidium foveolarum</i>	503
<i>Phormidium foveolarum</i>	504
<i>Phormidium foveolarum</i>	505
<i>Phormidium jenkelianum</i>	506
<i>Phormidium jenkelianum</i>	507
<i>Phormidium molle</i>	509
<i>Phormidium mucicola</i>	510
<i>Phormidium ramosum</i>	305
<i>Phormidium tenue</i>	30
<i>Phormidium tenue</i>	512
<i>Phormidium tenue</i>	611
<i>Plectonema radiosum</i>	515
<i>Spirulina platensis</i>	39
<i>Spirulina platensis</i>	45
<i>Spirulina platensis</i>	46
<i>Spirulina platensis</i>	597
<i>Spirulina subsalsa</i>	27
<i>Spirulina subsalsa</i>	527
<i>Spirulina subsalsa</i>	598
<i>Tolypothrix tenuis</i>	37

GLAUCOPHYTA

Glaucoephyceae

<i>Cyanophora paradoxa</i>	547
<i>Cyanophora paradoxa</i>	763
<i>Cyanophora tetracyanea</i>	764

RHODOPHYTA

Rhodophyceae

<i>Cyanidioschyzon merolae</i>	549
<i>Cyanidium caldarium</i>	551
<i>Galdieria sulphuraria</i>	250
<i>Galdieria sulphuraria</i>	550

CRYPTOPHYTA

Cryptophyceae

<i>Chilomonas paramecium</i>	715
<i>Chilomonas paramecium</i>	766
<i>Chilomonas paramecium</i>	767
<i>Chroomonas caudata</i>	712
<i>Chroomonas coerulea</i>	713
<i>Chroomonas coerulea</i>	714
<i>Chroomonas collegionis</i>	703
<i>Chroomonas dispersa</i>	704
<i>Chroomonas nordstedtii</i>	706
<i>Chroomonas nordstedtii</i>	707
<i>Chroomonas nordstedtii</i>	708
<i>Chroomonas nordstedtii</i>	709
<i>Chroomonas nordstedtii</i>	710
<i>Chroomonas nordstedtii</i>	711
<i>Chroomonas placoidea</i>	705
<i>Cryptomonas acuta</i>	697
<i>Cryptomonas irregularis</i>	698
<i>Cryptomonas ovata</i>	274
<i>Cryptomonas ovata</i>	275
<i>Cryptomonas platyuris</i>	276
<i>Cryptomonas platyuris</i>	344
<i>Cryptomonas rostratiformis</i>	277
<i>Cryptomonas rostratiformis</i>	278
<i>Cryptomonas rostratiformis</i>	345
<i>Cryptomonas tetrapyrenoidosa</i>	279
<i>Cryptomonas tetrapyrenoidosa</i>	280
<i>Cryptomonas tetrapyrenoidosa</i>	281
<i>Cryptomonas tetrapyrenoidosa</i>	282
<i>Cryptomonas tetrapyrenoidosa</i>	346
<i>Cryptomonas tetrapyrenoidosa</i>	347
<i>Cryptomonas tetrapyrenoidosa</i>	348

<i>Rhodomonas atrorosea</i>	699
<i>Rhodomonas baltica</i>	700
<i>Rhodomonas chrysoidea</i>	701
<i>Rhodomonas duplex</i>	765
<i>Rhodomonas falcata</i>	702

HETEROKONTOPHYTA

Chrysophyceae

<i>Dinobryon divergens</i>	284
<i>Synura petersenii</i>	233
<i>Synura sphagnicola</i>	695
<i>Synura sphagnicola</i>	696
<i>Synura spinosa</i>	234
<i>Uroglena americana</i>	395

Raphidophyceae

<i>Chattonella antiqua</i>	1
<i>Chattonella antiqua</i>	2
<i>Chattonella antiqua</i>	83
<i>Chattonella antiqua</i>	84
<i>Chattonella antiqua</i>	85
<i>Chattonella antiqua</i>	86
<i>Chattonella antiqua</i>	113
<i>Chattonella antiqua</i>	114
<i>Chattonella antiqua</i>	161
<i>Chattonella antiqua</i>	557
<i>Chattonella antiqua</i>	558
<i>Chattonella marina</i>	3
<i>Chattonella marina</i>	14
<i>Chattonella marina</i>	115
<i>Chattonella marina</i>	116
<i>Chattonella marina</i>	117
<i>Chattonella marina</i>	118
<i>Chattonella marina</i>	121
<i>Chattonella marina</i>	559
<i>Chattonella ovata</i>	603
<i>Chattonella ovata</i>	671
<i>Chattonella verruculosa</i>	670
<i>Fibrocapsa japonica</i>	136
<i>Fibrocapsa japonica</i>	462
<i>Fibrocapsa japonica</i>	560
<i>Fibrocapsa japonica</i>	605
<i>Haramonas dimorpha</i>	716
<i>Heterosigma akashiwo</i>	4
<i>Heterosigma akashiwo</i>	5

<i>Heterosigma akashiwo</i>	6
<i>Heterosigma akashiwo</i>	9
<i>Heterosigma akashiwo</i>	10
<i>Heterosigma akashiwo</i>	145
<i>Heterosigma akashiwo</i>	146
<i>Heterosigma akashiwo</i>	293
<i>Heterosigma akashiwo</i>	561
<i>Olisthodiscus luteus</i>	15

Bacillariophyceae

<i>Achnanthes longipes</i>	330
<i>Achnanthes minutissima</i>	71
<i>Achnanthes minutissima</i>	407
<i>Achnanthes minutissima</i>	408
<i>Achnanthes minutissima</i>	409
<i>Achnanthes minutissima</i>	410
<i>Achnanthes minutissima</i>	411
<i>Achnanthes minutissima</i>	412
<i>Achnanthes minutissima</i>	413
<i>Achnanthes minutissima</i>	414
<i>Achnanthes minutissima</i> var. <i>saprophila</i>	372
<i>Asterionella glacialis</i>	265
<i>Asterionella glacialis</i>	417
<i>Chaetoceros didymus</i>	586
<i>Chaetoceros sociale</i>	377
<i>Chaetoceros sociale</i>	553
<i>Cyclotella meneghiniana</i>	803
<i>Cyclotella meneghiniana</i>	804
<i>Cyclotella meneghiniana</i>	805
<i>Ditylum brightwellii</i>	350
<i>Eunotia pectinalis</i> var. <i>minor</i>	461
<i>Fragilaria capucina</i>	391
<i>Gomphonema angustatum</i> var. <i>obtusatum</i>	620
<i>Gomphonema gracile</i> var. <i>gracile</i>	465
<i>Gomphonema parvulum</i> var. <i>parvulum</i>	466
<i>Gomphonema parvulum</i> var. <i>parvulum</i>	467
<i>Hantzschia amphioxys</i> var. <i>compacta</i>	587
<i>Lithodesmium variabile</i>	588
<i>Melosira granulata</i>	333
var. <i>angustissima</i> f. <i>spiralis</i>	
<i>Nitzschia palea</i>	487
<i>Nitzschia palea</i>	488
<i>Nitzschia palea</i>	489
<i>Odontella aurita</i>	589
<i>Odontella longicruris</i>	590
<i>Skeletonema costatum</i>	16
<i>Skeletonema costatum</i>	17
<i>Skeletonema costatum</i>	223

<i>Skeletonema costatum</i>	323	<i>Amphidinium klebsii</i>	613
<i>Skeletonema costatum</i>	324	<i>Cachonina niei</i>	420
<i>Stephanopyxis palmeriana</i>	327	<i>Cachonina niei</i>	614
<i>Tabellaria flocculosa</i>	225	<i>Ceratium hirundinella</i>	376
<i>Thalassionema nitzschioides</i>	534	<i>Coolia monotis</i>	343
<i>Thalassiosira pacifica</i>	535	<i>Coolia monotis</i>	615
<i>Triceratium dubium</i>	556	<i>Glenodiniopsis uliginosa</i>	463
		<i>Gymnodinium fuscum</i>	470
		<i>Gymnodinium mikimotoi</i>	680
		<i>Gyrodinium instriatum</i>	143
		<i>Hemidinium nasutum</i>	471
		<i>Heterocapsa pygmaea</i>	472
		<i>Heterocapsa pygmaea</i>	473
		<i>Heterocapsa triquetra</i>	7
		<i>Heterocapsa triquetra</i>	235
		<i>Katodinium rotundatum</i>	356
		<i>Oxyrrhis marina</i>	494
		<i>Peridinium bipes</i> f. <i>globosum</i>	495
		<i>Peridinium bipes</i> f. <i>occultatum</i>	364
		<i>Peridinium bipes</i> f. <i>occultatum</i>	496
		<i>Peridinium bipes</i> f. <i>occultatum</i>	497
		<i>Peridinium bipes</i> var. <i>tabulatum</i>	600
		<i>Peridinium inconspicuum</i> subsp. <i>remotum</i>	499
		<i>Peridinium polonicum</i>	500
		<i>Peridinium volzii</i>	365
		<i>Peridinium volzii</i>	501
		<i>Peridinium wierzejskii</i>	502
		<i>Peridinium willei</i>	304
		<i>Peridinium willei</i>	366
		<i>Prorocentrum dentatum</i>	682
		<i>Prorocentrum gracile</i>	315
		<i>Prorocentrum lima</i>	617
		<i>Prorocentrum mexicanum</i>	317
		<i>Prorocentrum mexicanum</i>	618
		<i>Prorocentrum micans</i>	12
		<i>Prorocentrum micans</i>	218
		<i>Prorocentrum micans</i>	316
		<i>Prorocentrum micans</i>	601
		<i>Prorocentrum micans</i>	608
		<i>Prorocentrum minimum</i>	237
		<i>Prorocentrum minimum</i>	238
		<i>Prorocentrum sigmoides</i>	683
		<i>Prorocentrum triestinum</i>	219
		<i>Protoceratium reticulatum</i>	318
		<i>Protoceratium reticulatum</i>	319
		<i>Pyrocystis lunura</i>	609
		<i>Pyrophacus steinii</i>	321
		<i>Scrippsiella sweeneyae</i>	684
		<i>Scrippsiella trochoidea</i>	369
		<i>Woloszynskia leopoliense</i>	619
Phaeophyceae			
<i>Acinetospora crinita</i>	548		
Xanthophyceae			
<i>Botrydiopsis arrhiza</i>	621		
<i>Botrydium granulatum</i>	622		
HAPTOPHYTA			
Haptophyceae			
<i>Chrysochromulina hirta</i>	741		
<i>Chrysochromulina parva</i>	562		
<i>Cricosphaera roscoffensis</i>	8		
<i>Emiliana huxleyi</i>	837		
<i>Gephyrocapsa oceanica</i>	353		
<i>Gephyrocapsa oceanica</i>	838		
<i>Pavlova gyrans</i>	623		
<i>Phaeocystis pouchetii</i>	388		
DINOPHYTA			
Dinophyceae			
<i>Alexandrium affine</i>	673		
<i>Alexandrium catenella</i>	220		
<i>Alexandrium catenella</i>	519		
<i>Alexandrium catenella</i>	520		
<i>Alexandrium catenella</i>	674		
<i>Alexandrium catenella</i>	675		
<i>Alexandrium catenella</i>	677		
<i>Alexandrium hiranoi</i>	612		
<i>Alexandrium insuetum</i>	678		
<i>Amphidinium britannicum</i>	405		
<i>Amphidinium carterae</i>	331		

EUGLENOPHYTA

Euglenophyceae

<i>Euglena clara</i>	253
<i>Euglena gracilis</i>	47
<i>Euglena gracilis</i>	48
<i>Euglena gracilis</i> var. <i>bacillaris</i>	49
<i>Euglena mutabilis</i>	286
<i>Eutreptiella gymnastica</i>	381
<i>Phacus agilis</i>	387

CHLORARACHNIOPHYTA

Chlorarachniophyceae

<i>Chlorarachnion reptans</i>	624
-------------------------------	-----

CHLOROPHYTA

Prasinophyceae

<i>Mesostigma viride</i>	296
<i>Mesostigma viride</i>	475
<i>Mesostigma viride</i>	476
<i>Mesostigma viride</i>	477
<i>Monomastix minuta</i>	255
<i>Monomastix minuta</i>	256
<i>Nephroselmis astigmatica</i>	252
<i>Nephroselmis olivacea</i>	483
<i>Nephroselmis olivacea</i>	484
<i>Nephroselmis olivacea</i>	485
<i>Nephroselmis viridis</i>	486
<i>Pterosperma cristatum</i>	221
<i>Pterosperma cristatum</i>	626
<i>Pyramimonas</i> aff. <i>amylifera</i>	251
<i>Pyramimonas</i> aff. <i>amylifera</i>	320
<i>Pyramimonas parkeae</i>	254
<i>Tetraselmis cordiformis</i>	18
<i>Tetraselmis cordiformis</i>	533

Chlorophyceae

<i>Actinastrum hantzschii</i>	415
<i>Astrephomene gubernaculifera</i>	418
<i>Astrephomene gubernaculifera</i>	419

<i>Astrephomene gubernaculifera</i>	628
<i>Astrephomene perforata</i>	564
<i>Astrephomene perforata</i>	565
<i>Basichlamys sacculifera</i>	566
<i>Botryococcus braunii</i>	836
<i>Brachiomonas submarina</i>	375
<i>Carteria cerasiformis</i>	424
<i>Carteria cerasiformis</i>	425
<i>Carteria crucifera</i>	421
<i>Carteria crucifera</i>	630
<i>Carteria eugametos</i>	631
<i>Carteria eugametos</i>	632
<i>Carteria eugametos</i>	633
<i>Carteria eugametos</i>	634
<i>Carteria eugametos</i>	635
<i>Carteria eugametos</i>	636
<i>Carteria inversa</i>	422
<i>Carteria inversa</i>	423
<i>Carteria klebsii</i>	426
<i>Carteria multifilis</i>	427
<i>Carteria obtusa</i>	428
<i>Carteria obtusa</i>	429
<i>Carteria obtusa</i>	430
<i>Carteria obtusa</i>	431
<i>Carteria radiosa</i>	432
<i>Characiochloris acuminata</i>	637
<i>Characiochloris sasae</i>	567
<i>Characiochloris sasae</i>	638
<i>Characium angustum</i>	639
<i>Characium maximum</i>	154
<i>Characium polymorphum</i>	436
<i>Chlamydomonas augustae</i> var. <i>ellipsoidea</i>	158
<i>Chlamydomonas fasciata</i>	437
<i>Chlamydomonas monadina</i> var. <i>monadina</i>	438
<i>Chlamydomonas monticola</i>	157
<i>Chlamydomonas parkeae</i>	440
<i>Chlamydomonas parkeae</i>	441
<i>Chlamydomonas pulsatilla</i>	122
<i>Chlamydomonas tetragama</i>	446
<i>Chlorogonium capillatum</i>	692
<i>Chlorogonium capillatum</i>	742
<i>Chlorogonium capillatum</i>	743
<i>Chlorogonium capillatum</i>	744
<i>Chlorogonium capillatum</i>	745
<i>Chlorogonium capillatum</i>	746
<i>Chlorogonium capillatum</i>	747
<i>Chlorogonium capillatum</i>	748
<i>Chlorogonium capillatum</i>	749
<i>Chlorogonium capillatum</i>	750
<i>Chlorogonium elongatum</i>	751

<i>Chlorogonium elongatum</i>	752	<i>Eudorina unicocca</i> var. <i>unicocca</i>	725
<i>Chlorogonium elongatum</i>	753	<i>Gloeomonas lateperforata</i>	464
<i>Chlorogonium euchlorum</i>	754	<i>Gonium multicocum</i>	737
<i>Chlorogonium euchlorum</i>	755	<i>Gonium pectorale</i> var. <i>pectorale</i>	468
<i>Chlorogonium euchlorum</i>	756	<i>Gonium pectorale</i> var. <i>pectorale</i>	469
<i>Chlorogonium euchlorum</i>	757	<i>Gonium pectorale</i> var. <i>pectorale</i>	569
<i>Chlorogonium euchlorum</i>	758	<i>Gonium pectorale</i> var. <i>pectorale</i>	570
<i>Chlorogonium euchlorum</i>	759	<i>Gonium pectorale</i> var. <i>pectorale</i>	645
<i>Chlorogonium euchlorum</i>	760	<i>Gonium pectorale</i> var. <i>pectorale</i>	646
<i>Chlorogonium fusiforme</i>	123	<i>Gonium quadratum</i>	647
<i>Chlorogonium kasakii</i>	761	<i>Gonium quadratum</i>	648
<i>Chlorogonium neglectum</i>	439	<i>Gonium quadratum</i>	649
<i>Chloromonas insignis</i>	447	<i>Gonium quadratum</i>	650
<i>Chlorosarcinopsis caeca</i>	160	<i>Gonium quadratum</i>	651
<i>Chlorosarcinopsis delicata</i>	153	<i>Gonium quadratum</i>	652
<i>Coelastrum astroideum</i>	129	<i>Gonium quadratum</i>	653
<i>Coelastrum astroideum</i>	130	<i>Gonium viridistellatum</i>	288
<i>Coelastrum astroideum</i>	244	<i>Gonium viridistellatum</i>	289
<i>Coelastrum astroideum</i>	342	<i>Gonium viridistellatum</i>	290
<i>Coelastrum morus</i>	231	<i>Gonium viridistellatum</i>	654
<i>Coelastrum proboscideum</i>	131	<i>Gonium viridistellatum</i>	655
<i>Coelastrum reticulatum</i>	132	<i>Graesiella emersonii</i>	226
<i>Coelastrum reticulatum</i> var. <i>reticulatum</i>	245	<i>Graesiella emersonii</i>	687
<i>Dictyochloropsis irregularis</i>	378	<i>Graesiella emersonii</i>	688
<i>Dictyosphaerium pulchellum</i>	453	<i>Graesiella emersonii</i>	689
<i>Dimorphococcus lunatus</i>	134	<i>Graesiella emersonii</i>	690
<i>Dimorphococcus lunatus</i>	135	<i>Haematococcus lacustris</i>	144
<i>Draparnaldia plumosa</i>	454	<i>Hafniomonas montana</i>	257
<i>Echinospaeridium nordstedtii</i>	137	<i>Hafniomonas montana</i>	656
<i>Eremosphaera gigas</i>	379	<i>Hydrodictyon reticulatum</i>	295
<i>Eremosphaera viridis</i>	380	<i>Lagerheimia ciliata</i>	382
<i>Eremosphaera viridis</i>	643	<i>Lobomonas monstrosa</i>	474
<i>Eremosphaera viridis</i>	644	<i>Micractinium pusillum</i>	151
<i>Errerella bornhemiensis</i>	455	<i>Monoraphidium circinale</i>	480
<i>Eudorina cylindrica</i>	722	<i>Monoraphidium contortum</i>	384
<i>Eudorina elegans</i>	351	<i>Monoraphidium griffithii</i>	385
<i>Eudorina elegans</i> var. <i>carteri</i>	721	<i>Oedogonium obesum</i>	203
<i>Eudorina elegans</i> var. <i>elegans</i>	456	<i>Oocystis borgei</i>	659
<i>Eudorina elegans</i> var. <i>elegans</i>	457	<i>Oocystis lacustris</i>	660
<i>Eudorina elegans</i> var. <i>elegans</i>	717	<i>Oocystis lacustris</i>	661
<i>Eudorina elegans</i> var. <i>elegans</i>	718	<i>Oocystis lacustris</i>	662
<i>Eudorina elegans</i> var. <i>elegans</i>	719	<i>Pandorina colemaniae</i>	572
<i>Eudorina elegans</i> var. <i>elegans</i>	720	<i>Pandorina colemaniae</i>	573
<i>Eudorina elegans</i> var. <i>synoica</i>	458	<i>Pandorina morum</i>	242
<i>Eudorina elegans</i> var. <i>synoica</i>	568	<i>Pandorina morum</i>	243
<i>Eudorina illinoisensis</i>	459	<i>Pandorina morum</i>	362
<i>Eudorina illinoisensis</i>	460	<i>Pandorina morum</i> var. <i>morum</i>	574
<i>Eudorina illinoisensis</i>	723	<i>Pandorina morum</i> var. <i>morum</i>	575
<i>Eudorina unicocca</i> var. <i>peripheralis</i>	726	<i>Paulschulzia pseudovolvox</i>	727
<i>Eudorina unicocca</i> var. <i>unicocca</i>	724	<i>Pediastrum angulosum</i> var. <i>angulosum</i>	300

<i>Pediastrum boryanum</i>	209	<i>Treubaria triappendiculata</i>	394
<i>Pediastrum boryanum</i>	301	<i>Urnella terrestris</i>	156
<i>Pediastrum duplex</i>	212	<i>Uronema confervicolum</i>	538
<i>Pediastrum duplex</i> var. <i>duplex</i>	210	<i>Uronema gigas</i>	539
<i>Pediastrum duplex</i> var. <i>duplex</i>	213	<i>Uronema gigas</i>	540
<i>Pediastrum duplex</i> var. <i>gracillimum</i>	211	<i>Volvox aureus</i>	241
<i>Pediastrum duplex</i> var. <i>gracillimum</i>	214	<i>Volvox aureus</i>	396
<i>Pediastrum simplex</i>	215	<i>Volvox aureus</i>	693
<i>Pediastrum simplex</i>	302	<i>Volvox aureus</i>	694
<i>Pediastrum tetras</i>	216	<i>Volvox aureus</i> var. <i>aureus</i>	541
<i>Planctonema lauterbornii</i>	514	<i>Volvox aureus</i> var. <i>aureus</i>	542
<i>Platydorina caudata</i>	728	<i>Volvox barberi</i>	730
<i>Platydorina caudata</i>	729	<i>Volvox carteri</i>	397
<i>Pleodorina californica</i>	576	<i>Volvox carteri</i>	398
<i>Pleodorina californica</i>	735	<i>Volvox carteri</i> f. <i>kawasakiensis</i>	580
<i>Pleodorina indica</i>	736	<i>Volvox carteri</i> f. <i>kawasakiensis</i>	581
<i>Pleodorina japonica</i>	577	<i>Volvox carteri</i> f. <i>kawasakiensis</i>	732
<i>Polyedriopsis spinulosa</i>	232	<i>Volvox carteri</i> f. <i>kawasakiensis</i>	733
<i>Pseudocarteria mucosa</i>	522	<i>Volvox dissipatrix</i>	731
<i>Pseudocarteria mucosa</i>	523	<i>Volvox prolificus</i>	543
<i>Pseudocarteria mucosa</i>	524	<i>Volvox rousseletii</i>	734
<i>Pseudopleurococcus printzii</i>	159	<i>Volvox tertius</i>	544
var. <i>longissimus</i>		<i>Volvulina compacta</i>	582
<i>Pteromonas aculeata</i>	738	<i>Volvulina compacta</i>	583
<i>Pteromonas angulosa</i>	739	<i>Volvulina steinii</i>	545
<i>Pteromonas multipyrenoidea</i>	740	<i>Volvulina steinii</i>	546
<i>Scenedesmus acuminatus</i>	92	<i>Volvulina steinii</i>	584
var. <i>tetradesmoides</i>		<i>Volvulina steinii</i>	585
<i>Scenedesmus acutus</i>	94	<i>Yamagishiella unicocca</i>	578
<i>Scenedesmus acutus</i>	95	<i>Yamagishiella unicocca</i>	579
<i>Scenedesmus acutus</i>	120	<i>Yamagishiella unicocca</i>	666
<i>Scenedesmus dimorphus</i>	93	<i>Yamagishiella unicocca</i>	667
<i>Scenedesmus dimorphus</i>	119	<i>Yamagishiella unicocca</i>	762
<i>Scenedesmus gutwinskii</i> var. <i>heterospina</i>	797		
<i>Scenedesmus gutwinskii</i> var. <i>heterospina</i>	798		
<i>Scenedesmus gutwinskii</i> var. <i>heterospina</i>	799		
<i>Scenedesmus gutwinskii</i> var. <i>heterospina</i>	800		
<i>Scenedesmus gutwinskii</i> var. <i>heterospina</i>	801		
<i>Scenedesmus gutwinskii</i> var. <i>heterospina</i>	802		
<i>Scenedesmus quadricauda</i>	96		
<i>Scenedesmus serratus</i>	97		
<i>Schroederia setigera</i>	246		
<i>Selenastrum capricornutum</i>	35		
<i>Stigeoclonium aestivale</i>	531		
<i>Stigeoclonium fasciculare</i>	532		
var. <i>fasciculare</i>			
<i>Tetrabaena socialis</i>	691		
<i>Tetrabaena socialis</i> var. <i>socialis</i>	571		
<i>Tetracystis chlorococcoides</i>	155		
<i>Tetraëdron incus</i>	392		
		Trebouxiophyceae	
		<i>Auxenochlorella protothecoides</i>	629
		<i>Chlorella fusca</i> var. <i>fusca</i>	685
		<i>Chlorella saccharophila</i>	640
		<i>Chlorella vulgaris</i> var. <i>vulgaris</i>	227
		<i>Chlorella vulgaris</i> var. <i>vulgaris</i>	641
		<i>Chlorella vulgaris</i> var. <i>vulgaris</i>	642
		<i>Chlorella vulgaris</i> var. <i>vulgaris</i>	686
		<i>Microthamnion kützingianum</i>	479
		<i>Stichococcus bacillaris</i>	529
		<i>Stichococcus bacillaris</i>	530

Charophyceae

<i>Closterium acerosum</i>	124	<i>Closterium peracerosum-</i> <i>strigosum-littorale</i> complex	57
<i>Closterium acerosum</i>	125	<i>Closterium peracerosum-</i> <i>strigosum-littorale</i> complex	58
<i>Closterium acerosum</i>	127	<i>Closterium peracerosum-</i> <i>strigosum-littorale</i> complex	59
<i>Closterium acerosum</i>	448	<i>Closterium peracerosum-</i> <i>strigosum-littorale</i> complex	60
<i>Closterium aciculare</i> var. <i>subpronum</i>	258	<i>Closterium peracerosum-</i> <i>strigosum-littorale</i> complex	61
<i>Closterium aciculare</i> var. <i>subpronum</i>	259	<i>Closterium peracerosum-</i> <i>strigosum-littorale</i> complex	62
<i>Closterium calosporum</i> var. <i>calosporum</i>	271	<i>Closterium peracerosum-</i> <i>strigosum-littorale</i> complex	63
<i>Closterium calosporum</i> var. <i>galiciense</i>	128	<i>Closterium peracerosum-</i> <i>strigosum-littorale</i> complex	64
<i>Closterium calosporum</i> var. <i>galiciense</i>	162	<i>Closterium peracerosum-</i> <i>strigosum-littorale</i> complex	65
<i>Closterium calosporum</i> var. <i>galiciense</i>	163	<i>Closterium peracerosum-</i> <i>strigosum-littorale</i> complex	66
<i>Closterium calosporum</i> var. <i>galiciense</i>	164	<i>Closterium peracerosum-</i> <i>strigosum-littorale</i> complex	67
<i>Closterium calosporum</i> var. <i>galiciense</i>	165	<i>Closterium peracerosum-</i> <i>strigosum-littorale</i> complex	68
<i>Closterium calosporum</i> var. <i>galiciense</i>	166	<i>Closterium peracerosum-</i> <i>strigosum-littorale</i> complex	69
<i>Closterium calosporum</i> var. <i>galiciense</i>	167	<i>Closterium peracerosum-</i> <i>strigosum-littorale</i> complex	70
<i>Closterium calosporum</i> var. <i>galiciense</i>	168	<i>Closterium peracerosum-</i> <i>strigosum-littorale</i> complex	261
<i>Closterium calosporum</i> var. <i>himalayense</i>	169	<i>Closterium peracerosum-</i> <i>strigosum-littorale</i> complex	262
<i>Closterium calosporum</i> var. <i>himalayense</i>	170	<i>Closterium pleurodermatum</i>	449
<i>Closterium calosporum</i> var. <i>himalayense</i>	171	<i>Closterium praelongum</i> var. <i>brevius</i>	450
<i>Closterium calosporum</i> var. <i>himalayense</i>	336	<i>Closterium praelongum</i> var. <i>brevius</i>	451
<i>Closterium ehrenbergii</i>	228	<i>Closterium pusillum</i> var. <i>maius</i>	185
<i>Closterium ehrenbergii</i>	229	<i>Closterium rostratum</i> var. <i>subrostratum</i>	338
<i>Closterium gracile</i>	179	<i>Closterium selenastrum</i>	339
<i>Closterium gracile</i>	180	<i>Closterium selenastrum</i>	340
<i>Closterium incurvum</i>	181	<i>Closterium spinosporum</i> var. <i>crassum</i>	186
<i>Closterium incurvum</i>	337	<i>Closterium spinosporum</i> var. <i>crassum</i>	187
<i>Closterium moniliferum</i> var. <i>moniliferum</i>	172	<i>Closterium spinosporum</i> var. <i>crassum</i>	341
<i>Closterium moniliferum</i> var. <i>moniliferum</i>	173	<i>Closterium spinosporum</i> var. <i>malaysiense</i>	188
<i>Closterium moniliferum</i> var. <i>moniliferum</i>	174	<i>Closterium spinosporum</i> var. <i>malaysiense</i>	189
<i>Closterium moniliferum</i>	182	<i>Closterium spinosporum</i> var. <i>ryukyuense</i>	191
var. <i>submoniliferum</i>		<i>Closterium spinosporum</i> var. <i>ryukyuense</i>	192
<i>Closterium moniliferum</i>	183	<i>Closterium spinosporum</i> var. <i>ryukyuense</i>	193
var. <i>submoniliferum</i>		<i>Closterium spinosporum</i> var. <i>spinosporum</i>	194
<i>Closterium navicula</i>	175	<i>Closterium spinosporum</i> var. <i>spinosporum</i>	195
<i>Closterium navicula</i>	176	<i>Closterium spinosporum</i> var. <i>spinosporum</i>	196
<i>Closterium navicula</i>	177		
<i>Closterium navicula</i>	178		
<i>Closterium peracerosum-</i> <i>strigosum-littorale</i> complex	51		
<i>Closterium peracerosum-</i> <i>strigosum-littorale</i> complex	52		
<i>Closterium peracerosum-</i> <i>strigosum-littorale</i> complex	53		
<i>Closterium peracerosum-</i> <i>strigosum-littorale</i> complex	54		
<i>Closterium peracerosum-</i> <i>strigosum-littorale</i> complex	55		
<i>Closterium peracerosum-</i> <i>strigosum-littorale</i> complex	56		

PROTOZOA

Oligohymenophorea *

<i>Paramecium bursaria</i>	668
<i>Tetrahymena pyriformis</i>	403

* See Ref. 168.

X. REFERENCES

- 1 Allen, M. B. 1959 Studies with *Cyanidium caldarium*, an anomalously pigmented chlorophyte. *Arch. Mikrobiol.*, **32**, 270-277.
- 2 Aoyama, K., Uemura, I., Miyake, J. & Asada, Y. 1997 Fermentative metabolism to produce hydrogen gas and organic compounds in a cyanobacterium, *Spirulina platensis*. *J. Ferment. Bioeng.*, **83**, 17-20.
- 3 Aruga, Y. 1965 Ecological studies of photosynthesis and matter production of phytoplankton II. Photosynthesis of algae in relation to light intensity and temperature. *Bot. Mag. Tokyo*, **78**, 360-365.
- 4 Asada, Y. & Kawamura, S. 1986 Screening for cyanobacteria that evolve molecular hydrogen under dark and anaerobic conditions. *J. Ferment. Technol.*, **64**, 553-556.
- 5 Asada, Y., Kawamura, S. & Ho, K.-K. 1987 Hydrogenase from the unicellular cyanobacterium, *Microcystis aeruginosa*. *Phytochem.*, **26**, 637-640.
- 6 Asai, R., Matsukawa, R., Ikebukuro, K. & Karube, I. 1998 Chemiluminescence flow-injection system for rapid detection of red tide phytoplankton, *Chattonella antiqua*. *Anal. Lett.*, **31**, 2279-2288.
- 7 Baldia, S. F., Fukami, K., Nishijima, T. & Hata, T. 1995 Growth responses of *Spirulina platensis* to some physico-chemical factors and the kinetics of phosphorus utilization. *Fish. Sci.*, **61**, 331-335.
- 8 Baldia, S. F., Nishijima, T., Fukami, K. & Hata, Y. 1994 Studies on the chemical composition of a cyanophyte, *Spirulina platensis*. In *The Third Asian Fisheries Forum*, Eds. by Chou, L. M., et al., Asian Fisheries Society, Manila, Philippines, 1994, p.940-943.
- 9 Ban, S. 1992 Effects of photoperiod, temperature, and population density on induction of diapause egg production in *Eurytemora affinis* (Copepoda: Calanoida) in lake Ohnuma, Hokkaido, Japan. *J. Crustacean Biol.*, **12**, 361-367.
- 10 Ban, S. 1994 Effect of temperature and food concentration on post-embryonic development, egg production and adult body size of calanoid copepod *Eurytemora affinis*. *J. Plankton R.*, **16**, 721-735.
- 11 Ban, S. & Minoda, T. 1994 Induction of diapause egg production in *Eurytemora affinis* by their own metabolites. *Hydrobiologia*, **292/293**, 185-189.
- 12 Canini, A., Civitareale, P., Marini, S., Grilli Caiola, M. & Rotilio, G. 1992 Purification of iron superoxide dismutase from the cyanobacterium *Anabaena cylindrica* Lemm. and localization of the enzyme in heterocysts by immunogold labeling. *Planta*, **187**, 438-444.
- 13 Carefoot, J. R. 1968 Culture and heterotrophy of the freshwater dinoflagellate, *Peridinium cinctum* fa. *ovoplanum* Lindeman. *J. Phycol.*, **4**, 129-131.
- 14 Chihara, M., Inouye, I. & Takahata, N. 1986 *Oltmannsiellopsis*, a new genus of marine flagellate (Dunaliellaceae, Chlorophyceae). *Arch. Protistenk.*, **132**, 313-324.
- 15 Chu, S. P. 1942 The influence of the mineral composition of the medium on the growth of planktonic algae. Part I. Methods and culture media. *J. Ecol.*, **30**, 284-325.
- 16 Coesel, P. F. M. 1991 Ammonium dependency in *Closterium aciculare* T. West, a planktonic desmid from alkaline, eutrophic waters. *J. Plank. Res.*, **13**, 913-922.
- 17 Connell, L. & Cattolico, R. A. 1996 Fragile algae: axenic culture of field-collected samples of *Heterosigma carterae*. *Mar. Biol.*, **125**, 421-426.
- 18 De Halperin, D. R., Mendoza, M. L. & De Caire, G. Z. 1973 Axenic cultures of blue-green algae (Cyanophyta). *Physis secc. B. Buenos Aires.*, **32**, 67-84 (in Spanish with English summary).
- 19 Ehara, M., Kitayama, T., Watanabe, K. I., Inagaki, Y., Hayashi-Ishimaru, Y. & Ohama, T. 1999 Comprehensive molecular phylogenetic analysis of a heterokont alga (NIES 548) using

- genes from all three cellular compartments. *Phycol. Res.*, **47**, 225-231.
- 20 Ehara, M., Watanabe, K. I., Kawai, H., Inagaki, Y., Hayashi-Ishimaru, Y. & Ohama, T. 1998 Distribution of the mitochondrial deviant genetic code AUA for methionine in heterokont algae. *J. Phycol.*, **34**, 1005-1008.
- 21 Ehara, T., Shihira-Ishikawa, I., Osafune, T., Hase, E. & Ohkuro, I. 1975 Some structural characteristics of chloroplast degeneration in cells of *Euglena gracilis* Z during their heterotrophic growth in darkness. *J. Electron Microscopy*, **24**, 253-261.
- 22 Erata, M. & Chihara, M. 1987 Cryptomonads from the Sugadaira-Moor, Central Japan. *Bull. Sugadaira Mont. Res. Center, Univ. Tsukuba*, **8**, 57-69 (in Japanese with English summary).
- 23 Erata, M. & Chihara, M. 1989 Re-examination of *Pyrenomonas* and *Rhodomonas* (Class Cryptophyceae) through ultrastructural survey of red pigmented cryptomonads. *Bot. Mag. Tokyo*, **102**, 429-443.
- 24 Erata, M. & Chihara, M. 1991 Cryptomonads of Japan (2) three species of *Chroomonas*. *J. Jpn. Bot.*, **66**, 7-11.
- 25 Erata, M., Kubota, M., Takahashi, T., Inouye, I. & Watanabe, M. 1995 Ultrastructure and phototactic action spectra of two genera of cryptophyte flagellate algae, *Cryptomonas* and *Chroomonas*. *Protoplasma*, **188**, 258-266.
- 26 Foy, R. H. & Gibson, C. E. 1982 Photosynthetic characteristics of planktonic blue-green algae: the response of twenty strains grown under high and low light. *Br. phycol. J.*, **17**, 169-182.
- 27 Fujita, Y. 1974 The light-induced oxidation-reduction reaction of cytochrome *b*-559 in membrane fragments of the blue-green alga *Anabaena variabilis*. *Plant & Cell Physiol.*, **15**, 861-874.
- 28 Fujita, Y. 1975 Further investigation of the light-induced cytochrome *b*-559 reaction in membrane fragments of the blue-green alga *Anabaena variabilis*. *Plant & Cell Physiol.*, **16**, 1037-1048.
- 29 Fujita, Y. 1976 The C550 photoresponse at room temperature observed in membrane fragments of the blue-green alga *Anabaena variabilis*. *Plant & Cell Physiol.*, **17**, 187-191.
- 30 Fujita, Y. & Hattori, A. 1960 Formation of phycoerythrin in pre-illuminated cells of *Tolypothrix tenuis* with special reference to nitrogen metabolism. *Plant & Cell Physiol.*, **1**, 281-292.
- 31 Fujita, Y. & Hattori, A. 1960 Effect of chromatic lights on phycobilin formation in a blue-green alga, *Tolypothrix tenuis*. *Plant & Cell Physiol.*, **1**, 293-303.
- 32 Fujita, Y. & Hattori, A. 1962 Preliminary note on a new phycobilin pigment isolated from blue-green algae. *J. Biochem.*, **51**, 89-91.
- 33 Fujita, Y. & Hattori, A. 1962 Changes in composition of cellular material during formation of phycobilin chromoproteids in a blue-green alga, *Tolypothrix tenuis*. *J. Biochem.*, **52**, 38-42.
- 34 Fujita, Y. & Hattori, A. 1962 Photochemical interconversion between precursors of phycobilin chromoproteids in *Tolypothrix tenuis*. *Plant & Cell Physiol.*, **3**, 209-220.
- 35 Fujita, Y. & Hattori, A. 1963 Occurrence of a purple bile pigment in phycoerythrin-rich cells of the blue-green alga, *Tolypothrix tenuis*. *J. Gen. Appl. Microbiol.*, **9**, 253-256.
- 36 Fujita, Y. & Hattori, A. 1963 Action spectrum of light-induced nitrite reduction in *Anabaena cylindrica*. *J. Gen. Appl. Microbiol.*, **9**, 257-265.
- 37 Fujita, Y. & Hattori, A. 1963 Effects of second chromatic illumination on phycobilin chromoprotein formation in chromatically preilluminated cells of *Tolypothrix tenuis*. *Plant & Cell Physiol., Spec. Iss. (Studies on Microalgae and Photosynthetic Bacteria)*, 431-440.
- 38 Fujita, Y. & Murano, F. 1968 Occurrence of back flow of electrons against the action of photochemical system I in sonicated lamellar fragments. In *Comparative Biochemistry and Biophysics of Photosynthesis*, Eds. by Shibata, K., et al., University of Tokyo Press, Tokyo, p. 161-169.
- 39 Fujita, Y. & Myers, J. 1966 Some properties of the cytochrome *C* reducing substance, a factor

- for light-induced redox reaction of cytochrome C in photosynthetic lamellae. *Plant & Cell Physiol.*, **7**, 599-606.
- 40 Fujita, Y., Ohama, H. & Hattori, A. 1964 Hydrogenase activity of cell-free preparation obtained from the blue-green alga, *Anabaena cylindrica*. *Plant & Cell Physiol.*, **5**, 305-314.
- 41 Fujita, Y., Pjon, C.-J. & Suzuki, R. 1974 Studies on the Hill reaction of membrane fragments of blue-green algae V. A correlation between the solubilization of a photochemically active chromoprotein (ACP) and the inactivation of photosystem II activity in membrane fragments of *Anabaena cylindrica*. *Plant & Cell Physiol.*, **15**, 779-787.
- 42 Fujita, Y. & Suzuki, R. 1971 Studies on the Hill reaction of membrane fragments of blue-green algae I. Stabilizing effect of various media on the 2,6-dichlorophenol indophenol-Hill activity of membrane fragments obtained from *Anabaena cylindrica* and *Anabaena variabilis*. *Plant & Cell Physiol.*, **12**, 641-651.
- 43 Fujita, Y. & Suzuki, R. 1973 Studies on the Hill reaction of membrane fragments of blue-green algae III. Fluorescence characteristics of membrane fragments of *Anabaena variabilis* and *Anabaena cylindrica*. *Plant & Cell Physiol.*, **14**, 249-260.
- 44 Fujita, Y. & Suzuki, R. 1973 Studies on the Hill reaction of membrane fragments of blue-green algae IV. Carotenoid photobleaching induced by photosystem II action. *Plant & Cell Physiol.*, **14**, 261-273.
- 45 Fujita, Y. & Tsuji, T. 1968 Photochemically active chromoprotein isolated from the blue-green alga *Anabaena cylindrica*. *Nature*, **219**, 1270-1271.
- 46 Fukumoto, R., Fujii, T. & Sekimoto, H. 1997 Detection and evaluation of a novel sexual pheromone that induces sexual cell division of *Closterium ehrenbergii* (Chlorophyta). *J. Phycol.*, **33**, 441-445.
- 47 Fukumoto, R., Fujii, T. & Sekimoto, H. 1998 A newly identified chemotactic sexual pheromone from *Closterium ehrenbergii*. *Sex. Plant. Reprod.*, **11**, 81-85.
- 48 Gontcharov, A. A. & Watanabe, M. M. 1999 Rare and new desmids (Desmidiaceae, Chlorophyta) from Japan. *Phycol. Res.*, **47**, 233-240.
- 49 Grilli Caiola, M., Canini, A., Galiazzo, F. & Rotilio, G. 1991 Superoxide dismutase in vegetative cells, heterocysts and akinetes of *Anabaena cylindrica* Lemm. *FEMS Microbiol. Letters*, **80**, 161-166.
- 50 Guillard, R. R. L. & Ryther, J. H. 1962 Studies of marine planktonic diatoms. I. *Cyclotella nana* Hustedt, and *Detonula confervacea* (Cleve) Gran. *Can. J. Microbiol.*, **8**, 229-239.
- 51 Hagiwara, T., Yagi, O., Takamura, Y. & Sudo, R. 1984 Isolation of bacteria-free *Microcystis aeruginosa* from Lake Kasumigaura. *Jpn. J. Water Poll. Res.*, **7**, 437-442 (in Japanese with English summary).
- 52 Hanazato, T. 1991 Interrelations between *Microcystis* and Cladocera in the Highly Eutrophic Lake Kasumigaura, Japan. *Int. Revue ges. Hydrobiol.*, **76**, 21-36.
- 53 Hara, Y. 1990 *Chattonella ovata* Y. Hara et Chihara mss. In *Red Tide Organisms in Japan - An Illustrated Taxonomic Guide*, Eds. by Fukuyo, Y., Takano, H., Chihara, M. & Matsuoka, K., Uchida Rokakuho, Tokyo, p. 340-341 (in Japanese with English summary).
- 54 Hara, Y. & Chihara, M. 1982 Ultrastructure and taxonomy of *Chattonella* (Class Raphidophyceae) in Japan. *Jpn. J. Phycol.*, **30**, 47-56 (in Japanese with English summary).
- 55 Hara, Y., Doi, K. & Chihara, M. 1994 Four new species of *Chattonella* (Raphidophyceae, Chromophyta) from Japan. *Jpn. J. Phycol. (Sôru)*, **42**, 407-420.
- 56 Hara, Y., Inouye, I. & Chihara, M. 1985 Morphology and ultrastructure of *Olisthodiscus luteus* (Raphidophyceae) with special reference to the taxonomy. *Bot. Mag. Tokyo*, **98**, 251-262.
- 57 Harashima, A., Watanabe, M. & Fujishiro, I. 1988 Evolution of bioconvection patterns in a culture of motile flagellates. *Phys. Fluids*, **31**, 764-775.
- 58 Hashimoto, H. 1997 Electron-opaque annular structure girdling the constricting isthmus of the

- dividing chloroplasts of *Heterosigma akashiwo* (Raphidophyceae, Chromophyta). *Protoplasma*, **197**, 210-216.
- 59 Hatakeyama, S., Fukushima, S., Kasai, F. & Shiraishi, H. 1994 Assessment of herbicide effects on algal production in the Kokai River (Japan) using a model stream and *Selenastrum* bioassay. *Ecotoxicology*, **3**, 143-156.
- 60 Hatano, S., Hara, Y. & Takahashi, M. 1983 Preliminary study on the effects of photoperiod and nutrients on the vertical migratory behavior of a red tide flagellate, *Heterosigma akashiwo*. *Jpn. J. Phycol.*, **31**, 263-269 (in Japanese with English summary).
- 61 Hattori, A. & Fujita, Y. 1959 Crystalline phycobilin chromoproteids obtained from a blue-green alga, *Tolypothrix tenuis*. *J. Biochem.*, **46**, 633-644.
- 62 Hattori, A. & Fujita, Y. 1959 Spectroscopic studies on the phycobilin pigments obtained from blue-green and red algae. *J. Biochem.*, **46**, 903-909.
- 63 Hattori, A. & Fujita, Y. 1959 Effect of pre-illumination on the formation of phycobilin pigments in a blue-green alga, *Tolypothrix tenuis*. *J. Biochem.*, **46**, 1259-1261.
- 64 Hattori, A. & Uesugi, I. 1968 Ferredoxin-dependent photoreduction of nitrate and nitrite by subcellular preparations of *Anabaena cylindrica*. In *Comparative Biochemistry and Biophysics of Photosynthesis*, Eds. by Shibata, K., et al., University of Tokyo Press, Tokyo, p. 201-205.
- 65 Hayashi-Ishimaru, Y., Ehara, M., Inagaki, Y. & Ohama, T. 1997 A deviant mitochondrial genetic code in prymnesiophytes (yellow-algae): UGA codon for tryptophan. *Curr Genet*, **32**, 296-299.
- 66 Hayashi-Ishimaru, Y., Ohama, T., Kawatsu, Y., Nakamura, K. & Osawa, S. 1996 UAG is a sense codon in several chlorophycean mitochondria. *Curr Genet*, **30**, 29-33.
- 67 Hino, S. 1991 An examination of algal bioassay method for agricultural chemicals and heavy metals in a water environment. *Hokkaido Kankyō Kagaku Kenkyū Sentā Syohō*, **18**, 41-49 (in Japanese with English summary).
- 68 Hiroishi, S., & Uchida, A., Nagasaki, K. & Ishida, Y. 1988 A new method for identification of inter- and intra-species of the red tide algae *Chattonella antiqua* and *Chattonella marina* (Raphidophyceae) by means of monoclonal antibodies. *J. Phycol.*, **24**, 442-444.
- 69 Hiroishi, S., Uchida, A., Nagasaki, K. & Ishida, Y. 1989 A new method for inter- and intra-species identification of red tide algae *Chattonella antiqua* and *Chattonella marina* by means of monoclonal antibodies. In *Red Tides: Biology, Environmental Science, and Toxicology*, Eds. by Okaichi, T., Anderson, D. M. & Nemoto, T., Elsevier Science Pub., New York, p. 299-300.
- 70 Hiwatari, T., Kasai, F., Watanabe, M. M. & Nei, T. 1984 Cryopreservation of *Scenedesmus acutus* and *Pediastrum duplex* in liquid nitrogen - Survival and growth after freezing. *Jpn. J. Freez. Dry.*, **30**, 27-31 (in Japanese with English title).
- 71 Holm-Hansen, O., Gerloff, G. C. & Skoog, F. 1954 Cobalt as an essential element for blue-green algae. *Physiol. Planta.*, **7**, 665-675.
- 72 Horiguchi, T. 1996 *Haramonas dimorpha* gen. et sp. nov. (Raphidophyceae), a new marine raphidophyte from Australian mangrove. *Phycol. Res.*, **44**, 143-150.
- 73 Hosomi, M. & Sudo, R. 1979 Studies on the effects of sediments on algal growth - Algal growth potential of sediments. *Res. Rep. Natl. Inst. Environ. Stud.*, **No. 6**, 115-121 (in Japanese with English summary).
- 74 Hutner, S. H., Zahalsky, A. C., Aaronson, S., Baker, H. & Frank, O. 1966 Culture media for *Euglena gracilis*. In *Methods in Cell Physiology*. Vol. 2, Ed. by Prescott, D. M., Academic Press, New York, p. 217-228.
- 75 Ichimura, T. 1971 Sexual cell division and conjugation-papilla formation in sexual reproduction of *Closterium strigosum*. In *Proceedings of the Seventh International Seaweed Symposium*, University of Tokyo Press, Tokyo, p. 208-214.
- 76 Ichimura, T. 1973 The life cycle and its control in some species of *Closterium*, with special reference to the biological species problems. *Thesis D. Sci.*, University of Tokyo, 69 pp., 11 tables, 40 figs.

- 77 Ichimura, T. 1979 2. Isolation and culture methods of algae. 2.5.B. Freshwater algae [2. Sôruï no bunri to baiyôhô. 2.5.B. Tansui sôruï]. In *Methods in Phycological Studies* [Sôruï Kenkyûhô], Eds. by Nishizawa, K. & Chihara, M., Kyoritsu Shuppan, Tokyo, p. 294-305 (in Japanese without English title).
- 78 Ichimura, T. 1981 Mating types and reproductive isolation in *Closterium ehrenbergii* Meneghini. *Bot. Mag. Tokyo*, **94**, 325-334.
- 79 Ichimura, T. 1983 Hybrid inviability and predominant survival of mating type minus progeny in laboratory crosses between two closely related mating groups of *Closterium ehrenbergii*. *Evolution*, **37**, 252-260.
- 80 Ichimura, T. & Itoh, T. 1977 17. Preservation methods of microalgae (I) [17. Bisaisôruï no hozonhô (I)]. In *Preservation methods of microorganisms* [Biseibutsu Hozonhô], Ed. by Nei, T., University of Tokyo Press, Tokyo, p. 355-373 (in Japanese without English title).
- 81 Ichimura, T. & Kasai, F. 1984 Post-zygotic isolation between allopatric mating groups of *Closterium ehrenbergii* Meneghini (Conjugatophyceae). *Phycologia*, **23**, 77-85.
- 82 Ichimura, T. & Kasai, F. 1984 Time lapse analyses of sexual reproduction in *Closterium ehrenbergii* (Conjugatophyceae). *J. Phycol.*, **20**, 258-265.
- 83 Ichimura, T. & Kasai, F. 1985 Studies on the life cycle of *Spinoclosterium cuspidatum* (Bailey) Hirano (Conjugatophyceae). *Phycologia*, **24**, 205-216.
- 84 Ichimura, T. & Kasai, F. 1987 Time-lapse analyses of sexual isolation between two closely related mating groups of the *Closterium ehrenbergii* species complex (Chlorophyta). *J. Phycol.*, **23**, 523-534.
- 85 Ichimura, T. & Kasai, F. 1989 Life cycles of homothallic and heterothallic clones of *Triploceras gracile* Bailey (Desmidiaceae, Chlorophyta). *Phycologia*, **28**, 212-221.
- 86 Ichimura, T. & Watanabe, M. 1974 The *Closterium calosporum* complex from the Ryukyu Islands - Variation and taxonomical problems. *Mem. Natn. Sci. Mus. Tokyo*, **7**, 89-102, pls. 13-14.
- 87 Ichimura, T. & Watanabe, M. M. 1977 An axenic clone of *Microcystis aeruginosa* Kütz. emend. Elenkin from Lake Kasumigaura. *Bull. Jpn. Soc. Phycol.*, **25**, 177-181.
- 88 Ikushima, T. 1975 Action of furylfuramide (AF-2) on chloroplasts of *Euglena gracilis*. *Annu. Rep. Res. Reactor Inst. Kyoto Univ.*, **8**, 83-85.
- 89 Imai, A., Fukushima, T. & Matsushige, K. 1999 Effects of aquatic humic substances on the growth of the cyanobacterium *Microcystis aeruginosa*. *J. Jpn. Soc. Wat. Environ.* [Mizukankyô Gakkai Shi], **22**, 555-560 (in Japanese with English summary).
- 90 Imai, I., Ishida, Y. & Hata, Y. 1993 Killing of marine phytoplankton by a gliding bacterium *Cytophaga* sp., isolated from the coastal sea of Japan. *Mar. Biol.*, **116**, 527-532.
- 91 Imai, I., Ishida, Y., Sakaguchi, K. & Hata, Y. 1995 Algicidal marine bacteria isolated from northern Hiroshima Bay, Japan. *Fish. Sci.*, **61**, 628-636.
- 92 Imai, I., Ishida, Y., Sawayama, S. & Hata, Y. 1991 Isolation of a marine gliding bacterium that kills *Chattonella antiqua* (Raphidophyceae). *Nippon Suisan Gakkaishi*, **57**, 1409.
- 93 Imai, I., Kim, M. C., Nagasaki, K., Itakura, S. & Ishida, Y. 1998 Detection and enumeration of microorganisms that are lethal to harmful phytoplankton in coastal waters. *Plankton Biol. Ecol.*, **45**, 19-29.
- 94 Imai, I., Kim, M. C., Nagasaki, K., Itakura, S. & Ishida, Y. 1998 Relationships between dynamics of red tide-causing raphidophycean flagellates and algicidal micro-organisms in the coastal sea of Japan. *Phycol. Res.*, **46**, 139-146.
- 95 Imai, I., Nakagiri, S., Nagai, K., Nagasaki, K., Itakura, S. & Yamaguchi, M. 1998 Fluctuations of algicidal microorganisms against the harmful dinoflagellate *Heterocapsa circularisquama* in Ago Bay, Mie Prefecture, Japan. *Bull. Nansei Natl. Fish. Res. Inst.*, **31**, 53-61 (in Japanese with English summary).
- 96 Inagaki, Y., Ehara, M., Watanabe, K. I., Hayashi-Ishimaru, Y. & Ohama, T. 1998 Directionally evolving genetic code: the UGA codon from stop to tryptophan in mitochondria. *J. Mol. Evol.*, **47**,

- 378-384.
- 97 Inagaki, Y., Hayashi-Ishimaru, Y., Ehara, M., Igarashi, I. & Ohama, T. 1997 Algae or protozoa: phylogenetic position of euglenophytes and dinoflagellates as inferred from mitochondrial sequences. *J. Mol. Evol.*, **45**, 295-300.
- 98 Inamori, Y., Ohno, Y., Kaya, K., Watanabe, M. M. & Sudo, R. 1990 Studies on the removal and decomposition of *Microcystis viridis* toxic substance in the bio-film reactor. *Jpn. J. Water Poll. Res.*, **13**, 525-530 (in Japanese with English summary).
- 99 Ishibashi, Y., Ichianagi, J., Gotoh, M. & Konno, H. 1993 Culture conditions and characteristics of offensive odor-producing substance in the genus *Uroglena* [*Uroglena zoku no baiyôjôken to syûkibusshitsu tokusei*]. In *Proceedings of 27th Annual Meeting of Japan Society on Water Environment* [Dai 27kai Nippon Mizukankyô-Gakkai Nenkaikôensyû], p. 282-283 (in Japanese with English title).
- 100 Ishida, K., Matsuda, H., Murakami, M. & Yamaguchi, K. 1996 Kawaguchipeptin A, a novel cyclic undecapeptide from cyanobacterium *Microcystis aeruginosa* (NIES-88). *Tetrahedron*, **52**, 9025-9030.
- 101 Ishida, K., Matsuda, H., Murakami, M. & Yamaguchi, K. 1996 The absolute stereochemistry of micropeptin 90. *Tetrahedron Letters*, **37**, 9225-9226.
- 102 Ishida, K., Matsuda, H., Murakami, M. & Yamaguchi, K. 1997 Kawaguchipeptin B, an antibacterial cyclic undecapeptide from the cyanobacterium *Microcystis aeruginosa*. *J. Nat. Prod.*, **60**, 724-726.
- 103 Ishida, K., Matsuda, H., Murakami, M. & Yamaguchi, K. 1997 Microginins 299-A and -B, leucine aminopeptidase inhibitors from the cyanobacterium *Microcystis aeruginosa* (NIES-299). *Tetrahedron*, **53**, 10281-10288.
- 104 Ishida, K., Matsuda, H., Murakami, M. & Yamaguchi, K. 1997 Micropeptins 478-A and -B, plasmin inhibitors from the cyanobacterium *Microcystis aeruginosa*. *J. Nat. Prod.*, **60**, 184-187.
- 105 Ishida, K., Murakami, M., Matsuda, H. & Yamaguchi, K. 1995 Micropeptin 90, a plasmin and trypsin inhibitor from the blue-green alga *Microcystis aeruginosa* (NIES-90). *Tetrahedron Letters*, **36**, 3535-3538.
- 106 Ishimitsu, M. & Chihara, M. 1984 Four species of *Cryptomonas* (Class Cryptophyceae) in Japan. *J. Jpn. Bot.*, **59**, 161-169.
- 107 Ishitsuka, M. O., Kusumi, T., Kakisawa, H., Kaya, K. & Watanabe, M. M. 1990 Microviridin: a novel tricyclic depsipeptide from the toxic cyanobacterium *Microcystis viridis*. *J. Am. Chem. Soc.*, **112**, 8180-8182.
- 108 Itoh, Y., Murakami, H. & Ikeda, R. 1993 Studies on the growth limiting nutrients of red tide plankton in the tidal area. [Kanchô-iki ni okeru akashio gen'in sôru no zôshoku yokusei inshi no suitei.] *Proc. Coast. Engineering, JSCE* [*Kaigan Kôgaku Ronbun-shû*], **40**, 1001-1005 (in Japanese without English title).
- 109 Iwami, N., Inamori, Y., Sugiura, N. & Matsumura, M. 1996 Degradation characteristics of toxic *Microcystis viridis* by Mastigophora, *Monas guttula* using aquatic large scale microcosm. *J. Jpn. Soc. Wat. Environ.* **19**, 140-146.
- 110 Iwasaki, H., Fujiyama, T. & Yamashita, E. 1968 Studies on the red tide dinoflagellates - I. On *Entomosigma* sp. appeared in coastal area of Fukuyama. *J. Fac. Fish. Anim. Husb. Hiroshima Univ.*, **7**, 259-267 (in Japanese with English summary).
- 111 Iwasaki, H. & Sasada, K. 1969 Studies on the red tide dinoflagellates - II. On *Heterosigma inlandica* appeared in Gokasho Bay, Shima Peninsula. *Bull. Jpn. Soc. Sci. Fish.*, **35**, 943-947 (in Japanese with English summary).
- 112 Iwata, H., et al. 1991 Methods for producing heat-unsoluble β -1,3-glucan [Kanetsu gyôkosei β -1,3-glucan no seizôhō]. In *Nippon-koku Tokkyo-chô (JP), Kôkai Tokkyo Kôhō (A), Hei3-163102*, 5 pp (in Japanese without English title).
- 113 Johnson, L.M., Hoagland, K.D. & Gretz, M.R. 1995 Effects of bromide and iodine on stalk secretion in the biofouling diatom *Achnanthes longipes* (Bacillariophyceae). *J. Phycol.*, **31**, 401-412.

- 114 Kajiwara, S., Kakizono, T., Saito, T., Kondo, K., Ohtani, T., Nishio, N., Nagai, S. & Misawa, N. 1995 Isolation and functional identification of a novel cDNA for astaxanthin biosynthesis from *Haematococcus pluvialis*, and astaxanthin synthesis in *Escherichia coli*. *Plant Mol. Biol.*, **29**, 343-352.
- 115 Kakizono, T., Kobayashi, M. & Nagai, S. 1992 Effect of carbon/nitrogen ratio on encystment accompanied with astaxanthin formation in a green alga, *Haematococcus pluvialis*. *J. Ferment. Bioengineering*, **74**, 403-405.
- 116 Kaneko, H. & Yamaguchi, M. 1995 Toxicity evaluation of municipal solid waste incinerator fly ash by algal assay. *Reports Facul. Eng. Yamanashi Univ.*, **46**, 84-89 (in Japanese with English summary).
- 117 Kaneko, H., Yamaguchi, M. & Ikeda, F. 1996 Toxicity of municipal solid waste incinerator fly ash and the contribution of Cadmium. *J. Jpn. Soc. Wat. Environ. [Mizukankyō Gakkai Shi]*, **19**, 563-568 (in Japanese with English summary).
- 118 Kang, B.-H., Tan, S., & Ho, K.-K. 1988 Isolation of water-soluble cytochromes from cyanobacteria by adsorption chromatography. *Archiv. Biochem. Biophysics*, **263**, 387-393.
- 119 Kang, C. B. H. & Ho, K.-K. 1991 Characterization of a soluble inorganic pyrophosphatase from *Microcystis aeruginosa*. *Archiv. Biochem. Biophysics*, **289**, 281-288.
- 120 Kasai, F. & Arts, M. T. 1998 The interactive effects of UV-B radiation and a herbicide on uptake and allocation of carbon in two strains of the green alga *Scenedesmus*. *Aqua. Ecol.*, **31**, 261-272.
- 121 Kasai, F. & Hanazato, T. 1995 Genetic changes in phytoplankton communities exposed to the herbicide simetryn in outdoor experimental ponds. *Arch. Environ. Contam. Toxicol.*, **28**, 154-160.
- 122 Kasai, F. & Hatakeyama, S. 1993 Herbicide susceptibility in two green algae, *Chlorella vulgaris* and *Selenastrum capricornutum*. *Chemosphere*, **27**, 899-904.
- 123 Kasai, F. & Hatakeyama, S. 1996 Changes in herbicide susceptibility of algae in a river running through an agricultural region. *Ecol. Chem. (St. Petersburg)*, **5**, 292-296.
- 124 Kasai, F. & Ichimura, T. 1986 Morphological variabilities of three closely related mating groups of *Closterium ehrenbergii* Meneghini (Chlorophyta). *J. Phycol.*, **22**, 158-168.
- 125 Kasai, F. & Ichimura, T. 1990 Temperature optima of three closely related mating groups of the *Closterium ehrenbergii* (Chlorophyta) species complex. *Phycologia*, **29**, 396-402.
- 126 Kasai, F., Takamura, N. & Hatakeyama, S. 1993 Effects of simetryn on growth of various freshwater algal taxa. *Environ. Poll.*, **79**, 77-83.
- 127 Kato, A., Obokata, J. & Sasaki, K. 1981 Mating type interaction in *Closterium peracerosum-strigosum-littorale*: mating induced protoplast release. *Plant & Cell Physiol.*, **22**, 1215-1222.
- 128 Kato, A., Ohmura, K., Kanazawa, H. & Sasaki, K. 1983 Natural and artificial production of protoplasts from heterothallic and homothallic *Closterium*. *J. Fac. Sci., Hokkaido Univ., Ser. V (Bot.)*, **13**, 7-15.
- 129 Kato, A. & Sasaki, K. 1983 Effects of tunica-mycin on sexual reproduction in heterothallic strains of *Closterium*. *J. Fac. Sci., Hokkaido Univ., Ser. V (Bot.)*, **13**, 1-6.
- 130 Kato, S. 1982 Laboratory culture and morphology of *Colacium vesiculosum* Ehrb. (Euglenophyceae). *Jpn. J. Phycol.*, **30**, 63-67 (in Japanese with English summary).
- 131 Kawanobe, K. 1990 A report in the completion of the research work [Chōsa kenkyū syūryō no hōkoku]. Unpublished report in Kurashiki-shi Kankyō Hozen Kyōkai, 3 pp. (in Japanese without English title).
- 132 Kaya, K. & Sano, T. 1998 A photodetoxification mechanism of the cyanobacterial hepatotoxin microcystin-LR by ultraviolet irradiation. *Chem. Res. Toxicol.*, **11**, 159-163.
- 133 Kaya, K. & Watanabe, M. M. 1990 Microcystin composition of an axenic clonal strain of *Microcystis viridis* and *Microcystis viridis* containing waterblooms in Japanese freshwaters. *J. Appl. Phycol.*, **2**, 173-178.
- 134 Kikuchi, T., Ishida, M. R., Matsubara, T., Tsushimoto, G. & Mizuma, N. 1973 Some

- features of RNA and protein synthesis in thermophilic alga *Cyanidium caldarium*: the effects of antibiotics on *in vivo* incorporation of ¹⁴C-uracil and ¹⁴C-leucine. *Annu. Rep. Res. Reactor Inst. Kyoto Univ.*, **6**, 29-37.
- 135 Kim, Y.-S., Oyaizu, H., Matsumoto, S., Watanabe, M. M. & Nozaki, H. 1994 Chloroplast small-subunit ribosomal RNA gene sequence from *Chlamydomonas parkeae* (Chlorophyta): molecular phylogeny of a green alga with a peculiar pigment composition. *Eur. J. Phycol.*, **29**, 213-217.
- 136 Kimura, B. & Ishida, Y. 1985 Photophagotrophy in *Uroglena americana*, Chrysophyceae. *Jpn. J. Limnol.*, **46**, 315-318.
- 137 Kimura, B. & Ishida, Y. 1986 Possible phagotrophic feeding of bacteria in a freshwater red tide Chrysophyceae *Uroglena americana*. *Bull. Jpn. Soc. Sci. Fish.*, **52**, 697-701.
- 138 Kimura, B., Ishida, Y. & Kadota, H. 1986 Effect of naturally collected bacteria on growth of *Uroglena americana*, a freshwater red tide Chrysophyceae. *Bull. Jpn. Soc. Sci. Fish.*, **52**, 691-696.
- 139 Kita, T. & Fukuyo, Y. 1988 Description of the Gonyaulacoid dinoflagellate *Alexandrium hiranoi* sp. nov. inhabiting tidepools on Japanese Pacific coast. *Bull. Plank. Soc. Jpn.*, **35**, 1-7.
- 140 Kita, T., Fukuyo, Y., Tokuda, H. & Hirano, R. 1993 Sexual reproduction of *Alexandrium hiranoi* (Dinophyceae). *Bull. Plank. Soc. Jpn.*, **39**, 79-85.
- 141 Kiyohara, T., Fujita, Y., Hattori, A. & Watanabe, A. 1962 Effect of light on glucose assimilation in *Tolypothrix tenuis*. *J. Gen. Appl. Microbiol.*, **8**, 165-168.
- 142 Kobayashi, M., Hirai, N., Kurimura, Y., Ohigashi, H. & Tsuji, Y. 1997 Abscisic acid-dependent algal morphogenesis in the unicellular green alga *Haematococcus pluvialis*. *Plant Growth Regulation*, **22**, 79-85.
- 143 Kobayashi, M., Kakizono, T. & Nagai, S. 1991 Astaxanthin production by a green alga, *Haematococcus pluvialis* accompanied with morphological changes in acetate media. *J. Ferment. Bioengineering*, **71**, 335-339.
- 144 Kobayashi, M., Kakizono, T. & Nagai, S. 1993 Enhanced carotenoid biosynthesis by oxidative stress in acetate-induced cyst cells of a green unicellular alga, *Haematococcus pluvialis*. *Appl. Environ. Microbiol.*, **59**, 867-873.
- 145 Kobayashi, M., Kakizono, T. & Nagai, S. 1993 Photo-dependent astaxanthin biosynthesis in a green alga, *Haematococcus pluvialis*. *Seibutsu-Kôgaku*, **71**, 233-237 (in Japanese with English summary).
- 146 Kobayashi, M., Kakizono, T., Nishio, N. & Nagai, S. 1992 Effects of light intensity, light quality, and illumination cycle on astaxanthin formation in a green alga, *Haematococcus pluvialis*. *J. Ferment. Bioengineering*, **74**, 61-63.
- 147 Kobayashi, M., Kakizono, T., Nishio, N., Nagai, S., Kurimura, Y. & Tsuji, Y. 1997 Antioxidant role of astaxanthin in the green alga *Haematococcus pluvialis*. *Appl. Microbiol. Biotechnol.*, **48**, 351-356.
- 148 Kobayashi, M., Kakizono, T., Yamaguchi, K., Nishio, N. & Nagai, S. 1992 Growth and astaxanthin formation of *Haematococcus pluvialis* in heterotrophic and mixotrophic conditions. *J. Ferment. Bioengineering*, **74**, 17-20.
- 149 Kobayashi, M., Kurimura, Y., Kakizono, T., Nishio, N. & Tsuji, Y. 1997 Morphological changes in the life cycle of the green alga *Haematococcus pluvialis*. *J. Ferment. Bioeng.*, **84**, 94-97.
- 150 Kobayashi, M., Kurimura, Y., Sakamoto, Y. & Tsuji, Y. 1997 Selective extraction of astaxanthin and chlorophyll from the green alga *Haematococcus pluvialis*. *Biotech. Tech.*, **11**, 657-660.
- 151 Kobayashi, M., Kurimura, Y. & Tsuji, Y. 1997 Light-independent, astaxanthin production by the green microalga *Haematococcus pluvialis* under salt stress. *Biotech. Lett.*, **19**, 507-509.
- 152 Kobayashi, M., Todoroki, Y., Hirai, N., Kurimura, Y., Ohigashi, H. & Tsuji, Y. 1998 Biological activities of abscisic acid analogs in the morphological change of the green alga *Haematococcus pluvialis*. *J. Ferment. Bioeng.*, **85**, 529-531.

- 153 Kohata, K. & Watanabe, M. 1984 The use of a controlled experimental ecosystem (Microcosm) in studies of mechanism of red tide outbreaks (II) - Growth of *Heterosigma akashiwo*, red tide flagellate, in Microcosm. *Res. Rep. Natl. Inst. Environ. Stud.*, No. 63, 111-121 (in Japanese with English summary).
- 154 Kohata, K. & Watanabe, M. 1986 Synchronous division and the pattern of diel vertical migration of *Heterosigma akashiwo* (Hada) Hada (Raphido-phyceae) in a laboratory culture tank. *J. Exp. Mar. Biol. Ecol.*, 100, 209-224.
- 155 Kohata, K. & Watanabe, M. 1987 The use of a controlled experimental ecosystem (Microcosm) in studies of mechanism of red tide outbreaks (V) - Growth and cell volume change of *Heterosigma akashiwo*, a red tide flagellate. *Res. Rep. Natl. Inst. Environ. Stud.*, No. 110, 57-66 (in Japanese with English summary).
- 156 Kohata, K. & Watanabe, M. 1987 The use of a controlled experimental ecosystem (Microcosm) in studies of mechanism of red tide outbreaks (VI) - Diurnal changes of C/N ratio and distribution of chloroplast pigments in the culture of *Chattonella antiqua*, a red tide flagellate. *Res. Rep. Natl. Inst. Environ. Stud.*, No. 110, 67-77 (in Japanese with English summary).
- 157 Kohata, K. & Watanabe, M. 1987 The use of a controlled experimental ecosystem (Microcosm) in studies of mechanism of red tide outbreaks (VII) - Growth and its mathematical model of *Heterosigma akashiwo*, a red tide flagellate, in light-limited synchronous cultures. *Res. Rep. Natl. Inst. Environ. Stud.*, No. 110, 79-88 (in Japanese with English summary).
- 158 Kohata, K. & Watanabe, M. 1988 Diel changes in the composition of photosynthetic pigments and cellular carbon and nitrogen in *Chattonella antiqua* (Raphidophyceae). *J. Phycol.*, 24, 58-66.
- 159 Kohata, K. & Watanabe, M. 1989 Diel changes in the composition of photosynthetic pigments and cellular carbon and nitrogen in *Pyramimonas parkeae* (Prasinophyceae). *J. Phycol.*, 25, 377-385.
- 160 Kohata, K., Watanabe, M. & Yamanaka, K. 1991 Highly sensitive determination of photosynthetic pigments in marine *in situ* samples by high-performance liquid chromatography. *J. Chromatography*, 558, 131-140.
- 161 Kohata, K., Watanabe, M. M., Nakamura, Y. & Watanabe, M. 1982 Growth and phosphate uptake kinetics in *Olisthodiscus luteus*: grown on synchronized batch cultures. *Res. Rep. Natl. Inst. Environ. Stud.*, No. 30, 95-111 (in Japanese with English summary).
- 162 Konno, R. & Wakabayashi, A. 1987 Effects of chemical substance on the growth of algae [Sôru no zôsyoku ni oyobosu kagakubusshitsu no eikyô]. *Tokyo-to Kankyô Kagaku Kenkyûjo Nenpô*. 1987, 4 pp. (in Japanese without English title).
- 163 Kumagai, M., Takamura, Y., Yagi, O. & Sudo, R. 1986 Interactions between *Microcystis aeruginosa* and *Moraxella nonliquefaciens*. *Jpn. J. Limnol.*, 47, 219-228 (in Japanese with English summary).
- 164 Kusumi, T., Ooi, T., Watanabe, M. M., Takahashi, H. & Kakisawa, H. 1987 Cyanoviridin RR, a toxin from the cyanobacterium (blue-green alga) *Microcystis viridis*. *Tetrahedron Letters*, 28, 4695-4698.
- 165 Lee, K. Y., Park, J. H. & Park, B. S. 1996 CO₂ fixation and SCP production using blue green algae. *Theories and Applications of Chem. Eng. (Korea)*, 2, 489-492 (in Korean with English title).
- 166 Lee, T., Tsuzuki, M., Takeuchi, T., Yokoyama, K. & Karube, I. 1994 *In vivo* fluorometric method for early detection of cyanobacterial waterblooms. *J. Appl. Phycol.*, 6, 489-495.
- 167 Lee, T., Tsuzuki, M., Takeuchi, T., Yokoyama, K. & Karube, I. 1995 Quantitative determination of cyanobacteria in mixed phytoplankton assemblages by an *in vivo* fluorimetric method. *Anal. Chem. Acta*, 302, 81-87.
- 168 Levine, N. D., Corliss, J. O., Cox, F. E. G., Deroux, G., Grain, J., Honigberg, B. M., Leedale, G. F., Loeblich, III, A. R., Lom, J., Lynn, D., Merinfeld, E. G., Page, F. C., Poljansky, G., Sprague, V., Vavra, J. & Wallace, F. G. 1980 A newly revised classification of the Protozoa. *J. Protozool.*, 27, 37-58.
- 169 Li, R., Yokota, A., Sugiyama, J., Watanabe, M., Hiroki, M. & Watanabe, M. M. 1998

- Chemotaxonomy of planktonic cyanobacteria based on non-polar and 3-hydroxy fatty acid composition. *Phycol. Res.*, **46**, 21-28.
- 170 Liu, C. B., Lin, L. P. & Su, Y. C. 1996 Utilization of *Chlorella vulgaris* for uptake of nitrogen, phosphorus and heavy metals. *J. Chin. Agr. Chem. Soc.*, **34**, 331-343 (in Chinese with English summary).
- 171 Marin, B. & Melkonian, M. 1994 Flagellar hairs in prasinophytes (Chlorophyta): ultrastructure and distribution on the flagellar surface. *J. Phycol.*, **30**, 659-678.
- 172 Matsuda, H., Okino, T., Murakami, M. & Yamaguchi, K. 1996 Aeruginosins 102-A and B, new thrombin inhibitors from the cyanobacterium *Microcystis viridis* (NIES-102). *Tetrahedron*, **52**, 14501-14506.
- 173 Matsuda, H., Okino, T., Murakami, M. & Yamaguchi, K. 1996 Radiosumin, a trypsin inhibitor from the blue-green alga *Plectonema radiosum*. *J. Org. Chem.*, **61**, 8648-8650.
- 174 Matsunaga, S., Hori, T., Takahashi, T., Kubota, M., Watanabe, M., Okamoto, K., Masuda, K. & Sugai, M. 1998 Discovery of signaling effect of UV-B/C light in the extended UV-A/blue-type action spectra for step-down and step-up photophobic responses in the unicellular flagellate alga *Euglena gracilis*. *Protoplasma*, **201**, 45-52.
- 175 Merola, A., Castaldo, R., De Luca, P., Gambardella, R., Musacchio, A. & Taddei, R. 1981 Revision of *Cyanidium caldarium*. Three species of acidophilic algae. *Giorn. Bot. Ital.*, **115**, 189-195.
- 176 Mihara, S. & Hase, E. 1976 A short note on the effects of 6-methyl purine on the revolution of the cell cycle of *Chlorella pyrenoidosa* in synchronous culture. *Plant & Cell Physiol.*, **17**, 403-406.
- 177 Miya, A., Adachi, T., Tazawa, R., Suzuki, S., Kanki, R., Toyobe, M. & Oguchi, M. 1994 Gas revitalization by microalgae II. CO₂/O₂ gas exchange system. *CELLS J.*, **7**, 29-34 (in Japanese with English summary).
- 178 Miyagi, N., Satoh, S. & Fujii, T. 1989 Effects of *n*-alkylamines on the motility and viability of *Heterosigma akashiwo* cells. *Plant & Cell Physiol.*, **30**, 637-642.
- 179 Miyagi, N., Satoh, S. & Fujii, T. 1992 A nitrate-inducible plasma membrane protein of a marine alga, *Heterosigma akashiwo*. *Plant & Cell Physiol.*, **33**, 971-976.
- 180 Miyoshi, Y. & Tsubo, Y. 1969 Permanent bleaching of *Euglena* by chloramphenicol. *Plant & Cell Physiol.*, **10**, 221-225.
- 181 Mori, S., Nakamura, Y., Watanabe, M. M., Yamochi, S. & Watanabe, M. 1982 The effect of various environmental factors on the growth yield of red tide algae II. *Olisthodiscus luteus*. *Res. Rep. Natl. Inst. Environ. Stud.*, No. **30**, 71-86 (in Japanese with English summary).
- 182 Morimoto, T., Murakami, N., Nagatsu, A. & Sakakibara, J. 1993 Studies on glycolipids. VII. Isolation of two new sulfoquinovosyl diacylglycerols from the green alga *Chlorella vulgaris*. *Chem. Pharm. Bull.*, **41**, 1545-1548.
- 183 Morita, E., Abe, T., Tsuzuki, M., Fujiwara, S., Sato, N., Hirata, A., Sonoike, K. & Nozaki, H. 1999 Role of pyrenoids in the CO₂-concentrating mechanism: comparative morphology, physiology and molecular phylogenetic analysis of closely related strains of *Chlamydomonas* and *Chloromonas* (Volvocales). *Planta*, **208**, 365-372.
- 184 Murakami, M., Ishida, K., Okino, T., Okita, Y., Matsuda, H. & Yamaguchi, K. 1995 Aeruginosins 98-A and B, trypsin inhibitors from the blue-green alga *Microcystis aeruginosa* (NIES-98). *Tetrahedron Letters*, **36**, 2785-2788.
- 185 Murakami, M., Kodani, S., Ishida, K., Matsuda, H. & Yamaguchi, K. 1997 Micropeptin 103, a chymotrypsin inhibitor from the cyanobacterium *Microcystis viridis* (NIES-103). *Tetrahedron Letters*, **38**, 3035-3038.
- 186 Murakami, M., Makabe, K., Yamaguchi, K. & Konosu, S. 1988 Goniodomin A, a novel polyether macrolide from the dinoflagellate *Goniodoma pseudogoniaulax*. *Tetrahedron Letters*, **29**, 1149-1152.
- 187 Murakami, M., Okita, Y., Matsuda, H., Okino, T. & Yamaguchi, K. 1994 Aeruginosin 298-A, a thrombin and trypsin inhibitor from the blue-green alga *Microcystis aeruginosa* (NIES-298). *Tetrahedron Letters*, **35**, 3129-3132.

- 188 Murakami, M., Shin, H. J., Matsuda, H., Ishida, K. & Yamaguchi, K. 1997 A cyclic peptide, anabaenopeptin B, from the cyanobacterium *Oscillatoria agardhii*. *Phytochemistry*, **44**, 449-452.
- 189 Murakami, N., Shirahashi, H., Nagatsu, A. & Sakakibara, J. 1992 Two unsaturated 9R-hydroxy fatty acids from the cyanobacterium *Anabaena flos-aquae* f. *flos-aquae*. *Lipids*, **27**, 776-778.
- 190 Murakami, N., Shirahashi, H., Nagatsu, A. & Sakakibara, J. 1993 Studies on glycolipids. VI. New acyl-distributed glyceroglycolipids from the nitrogen-fixing cyanobacterium *Anabaena flos-aquae* f. *flos-aquae*. *Chem. Pharm. Bull.*, **41**, 1177-1179.
- 191 Murakami, N., Shirahashi, H., Sakakibara, J. & Tsuchida, Y. 1992 A novel glyceroglycolipid from the nitrogen-fixing cyanobacterium *Anabaena flos-aquae* f. *flos-aquae*. *Chem. Pharm. Bull.*, **40**, 285-287.
- 192 Murakami, M., Sun, Q., Ishida, K., Matsuda, H., Okino, T. & Yamaguchi, K. 1997 Microviridins, elastase inhibitors from the cyanobacterium *Nostoc minutum* (NIES-26). *Phytochemistry*, **45**, 1197-1202.
- 193 Murano, F. & Fujita, Y. 1967 Comparative studies of photochemical oxidation-reduction reactions in lamellar fragments of various algae and spinach. *Plant & Cell Physiol.*, **8**, 673-682.
- 194 Nagai, H., Satake, M., Murata, M. & Yasumoto, T. 1990 Screening of marine phytoplankton for antifungal substances. In *Toxic Marine Phytoplankton*, Eds. by Graneli, E., Sundstrom, B., Edler, L. & Anderson, D. M., Elsevier Science Pub., New York, p. 385-390.
- 195 Nakagawa, M., Takamura, Y. & Yagi, O. 1987 Isolation and characterization of the slime from a cyanobacterium, *Microcystis aeruginosa* K-3A. *Agric. Biol. Chem.*, **51**, 329-337.
- 196 Nakahara, H. & Sako, Y. 1987 2. Life history of freshwater phytoplankton [2. Tansui syokubutsu purankuton no seikatsushi]. In *Freshwater red tide [Tansui Akashio]*, Ed. by Kadota, H., Kôseisyu-Kôseikaku, Tokyo, p. 21-77 (in Japanese without English title).
- 197 Nakahara, M., Ito, H., Yano, H. & Harimaya, K. 1988 Eutrophication in three reservoirs as the sources of the waterworks in Kobe City. *Jpn. J. Water Treat. Biol.*, **24**, 19-24 (in Japanese with English title).
- 198 Nakamura, H. 1993 Results of AGP experiments [AGP, T-P, PO₄-P no shiken oyobi bunsekikekka]. Unpublished report in Ebara Yujiraito Co. Ltd., 32pp. (in Japanese without English title).
- 199 Nakamura, M., Yamagishi, M., Yoshizaki, F. & Sugimura, Y. 1992 The syntheses of plastocyanin and cytochrome *c*-553 are regulated by copper at the pre-translational level in a green alga, *Pediastrum boryanum*. *J. Biochem.*, **111**, 219-224.
- 200 Nakamura, Y. 1987 Effects of growth conditions on nitrate, ammonium and phosphate uptake by *Chattonella antiqua*. *Res. Rep. Natl. Inst. Environ. Stud.*, No. **110**, 115-126 (in Japanese with English summary).
- 201 Nakamura, Y., Sawai, K., Mochida, M. & Watanabe, M. 1987 An approach to estimate the limiting nutrient of *Chattonella antiqua* in the Seto Inland Sea - Semicontinuous culture using natural seawater. *Res. Rep. Natl. Inst. Environ. Stud.*, No. **110**, 197-212 (in Japanese with English summary).
- 202 Nakamura, Y., Sawai, K. & Watanabe, M. 1987 Growth inhibition of a red tide flagellate, *Chattonella antiqua* by cupric ion. *Res. Rep. Natl. Inst. Environ. Stud.*, No. **110**, 127-138 (in Japanese with English summary).
- 203 Nakamura, Y. & Watanabe, M. M. 1983 Growth characteristics of *Chattonella antiqua* (Raphidophyceae). Part 1. Effects of temperature, salinity, light intensity and pH on growth. *J. Oceanogr. Soc. Japan*, **39**, 110-114.
- 204 Nakamura, Y. & Watanabe, M. M. 1983 Growth characteristics of *Chattonella antiqua*. Part 2. Effects of nutrients on growth. *J. Oceanogr. Soc. Japan*, **39**, 151-155.
- 205 Nakamura, Y. & Watanabe, M. M. 1983 Nitrate and phosphate uptake kinetics of *Chattonella antiqua* grown in light/dark cycles. *J. Oceanogr. Soc. Japan*, **39**, 167-170.

- 206 Nakamura, Y. & Watanabe, M. M. 1984 Effects of temperature, salinity, light intensity and pH on the growth of *Chattonella antiqua*. *Res. Rep. Natl. Inst. Environ. Stud.*, No. 63, 79-85 (in Japanese with English summary).
- 207 Nakamura, Y. & Watanabe, M. M. 1984 Effects of nutrients on the growth of *Chattonella antiqua*. *Res. Rep. Natl. Inst. Environ. Stud.*, No. 63, 87-95 (in Japanese with English summary).
- 208 Nakamura, Y. & Watanabe, M. M. 1984 Diurnal vertical migration of a red tide flagellate, *Chattonella antiqua* with special reference to the ecological role. *Res. Rep. Natl. Inst. Environ. Stud.*, No. 63, 97-102 (in Japanese with English summary).
- 209 Nakamura, Y. & Watanabe, M. M. 1984 Nitrate and phosphate uptake kinetics of *Chattonella antiqua* grown in light/dark cycles. *Res. Rep. Natl. Inst. Environ. Stud.*, No. 63, 103-109 (in Japanese with English summary).
- 210 Nakamura, Y., Watanabe, M. M. & Watanabe, M. 1982 The effect of various environmental factors on the growth yield of red tide algae. I. *Chattonella antiqua*. *Res. Rep. Natl. Inst. Environ. Stud.*, No. 30, 53-70 (in Japanese with English summary).
- 211 Nakano, T. & Isagi, Y. 1987 *Dictyochloropsis irregularis* sp. nov. (Chlorococcales, Chlorophyceae) isolated from the surface of bark. *Phycologia*, 26, 222-227.
- 212 Nakayama, T., Ishii, E. & Kanazawa, H. 1987 The study on the isolation and culture of the protoplast from heterothallic *Closterium*. *J. Fac. Sci., Hokkaido Univ., Ser. V (Bot.)*, 14, 95-114.
- 213 Nakayama, T., Watanabe, S. & Inouye, I. 1996 Phylogeny of wall-less green flagellates inferred from 18SrDNA sequence data. *Phycol. Res.*, 44, 151-161.
- 214 Naohara, T. & Ishikawa, H. 1975 Assay of the activity of herbicides using *Chlorella*. *Zassô-Kenkyû*, 20, 82-86 (in Japanese with English summary).
- 215 Neilan, B.A., Hawkins, P.R., Cox, P.T. & Goodman, A.E. 1994 Towards a molecular taxonomy for the bloom-forming cyanobacteria. *Aust. J. Mar. Freshwater Res.*, 45, 869-873.
- 216 Neilan, B.A., Jacobs, D. & Goodman, A.E. 1995 Genetic diversity and phylogeny of toxic cyanobacteria determined by DNA polymorphisms within the phycocyanin locus. *Appl. Environ. Microbiol.*, 61, 3875-3883.
- 217 Nichols, P. D., Volkman, J. K., Hallegraeff, G. M. & Blackburn, S. I. 1987 Sterols and fatty acids of the red tide flagellates *Heterosigma akashiwo* and *Chattonella antiqua* (Raphidophyceae). *Phytochem.*, 26, 2537-2541.
- 218 Noike, T. & Souma, Y. 1993 Survey on the influence factors of eutrophication of lakes and the necessity of advanced waste treatment. [Koshô no fu-eiyôka no kanôsei to kôdo shori no hitsuyôsei ni kansuru chôsa.] *J. Jpn. Soc. Sewage Works [Gesuidô-Kyôkai Shi]*, 30, 38-44 (in Japanese without English title).
- 219 Nojiri, T., Fujii, T. & Sekimoto, H. 1995 Purification and characterization of a novel sex pheromone that induces the release of another sex pheromone during sexual reproduction of the heterothallic *Closterium peracerosum-strigosum-littorale* complex. *Plant Cell Physiol.*, 36, 79-84.
- 220 Nozaki, H. 1981 The life history of Japanese *Pandorina unicocca* (Chlorophyta, Volvocales). *J. Jpn. Bot.*, 56, 65-74.
- 221 Nozaki, H. 1982 Morphology and reproduction of Japanese *Volvolina steinii* (Chlorophyta, Volvocales). *J. Jpn. Bot.*, 57, 105-113, pls. 3-4.
- 222 Nozaki, H. 1983 Morphology and taxonomy of two species of *Astrephomene* (Chlorophyta, Volvocales) in Japan. *J. Jpn. Bot.*, 58, 345-352, pls. 6-8.
- 223 Nozaki, H. 1983 Sexual reproduction in *Eudorina elegans* (Chlorophyta, Volvocales). *Bot. Mag. Tokyo*, 96, 103-110.
- 224 Nozaki, H. 1984 Newly found facets in the asexual and sexual reproduction of *Gonium pectorale* (Chlorophyta, Volvocales). *Jpn. J. Phycol.*, 32, 130-133.
- 225 Nozaki, H. 1984 *Volvox aureus* Ehrenberg var. *aureus*. In *Photomicrog. Fresh-w. Alg.* [Tansui Sôru Syashinsyû] Vol. 1, Eds. by Yamagishi, T. & Akiyama, M., Uchida Rokakuho, Tokyo, p. 97 (in Japanese & English).

- 226 Nozaki, H. 1986 Notes on microalgae in Japan (9). *Basichlamys sacculifera* (Chlorophyta, Volvocales). *Jpn. J. Phycol.*, **34**, 143 (in Japanese with English title).
- 227 Nozaki, H. 1986 Notes on microalgae in Japan (10). *Eudorina illinoisensis* (Chlorophyta, Volvocales). *Jpn. J. Phycol.*, **34**, 144 (in Japanese with English title).
- 228 Nozaki, H. 1986 Sexual reproduction in *Gonium sociale* (Chlorophyta, Volvocales). *Phycologia*, **25**, 29-35.
- 229 Nozaki, H. 1986 Zygote germination in *Eudorina elegans* var. *synoica* (Chlorophyta, Volvocales). *J. Jpn. Bot.*, **61**, 316-320, pl. 8.
- 230 Nozaki, H. 1988 Colonial Volvocales (Chlorophyta) from Kathmandu, Nepal. In *Cryptogams of the Himalayas. Vol. 1. The Kathmandu Valley*, Eds. by Watanabe, M. & Malla, S. B., Department of Botany, National Science Museum, Tsukuba, p. 39-46.
- 231 Nozaki, H. 1988 Morphology, sexual reproduction and taxonomy of *Volvox carteri* f. *kawasakensis* f. nov. (Chlorophyta) from Japan. *Phycologia*, **27**, 209-220.
- 232 Nozaki, H. 1989 Morphological variation and reproduction in *Gonium viridistellatum* (Volvocales, Chlorophyta). *Phycologia*, **28**, 77-88.
- 233 Nozaki, H. 1990 Colonial Volvocales (Chlorophyta) from Central and East Nepal. In *Cryptogams of the Himalayas. Vol. 2. Central and East Nepal*, Eds. by Watanabe, M. & Malla, S.B., Department of Botany, National Science Museum, Tsukuba, p. 41-47.
- 234 Nozaki, H. 1990 Ultrastructure of the extracellular matrix of *Gonium* (Volvocales, Chlorophyta). *Phycologia*, **29**, 1-8.
- 235 Nozaki, H. 1993 Asexual and sexual reproduction in *Gonium quadratum* (Chlorophyta) with a discussion of phylogenetic relationships within the Goniaceae. *J. Phycol.*, **29**, 369-376.
- 236 Nozaki, H. 1993 Morphology, reproduction and taxonomy of *Characiochloris sasae* sp. nov. (Chlorophyta) from Japan. *Phycologia*, **32**, 129-135.
- 237 Nozaki, H. 1994 Aplanogamous sexual reproduction in *Carteria eugametos* (Volvocales, Chlorophyta). *Eur. J. Phycol.*, **29**, 135-139.
- 238 Nozaki, H. 1994 Unequal flagellar formation in *Volvox* (Volvocaceae, Chlorophyta). *Phycologia*, **33**, 58-61.
- 239 Nozaki, H., Aizawa, K. & Watanabe, M.M. 1994 A taxonomic study of four species of *Carteria* (Volvocales, Chlorophyta) with cruciate anterior papillae, based on cultured material. *Phycologia*, **33**, 239-247.
- 240 Nozaki, H., Aizawa, K. & Watanabe, M.M. 1996 Re-examination of two NIES strains labeled *Chlorogonium metamorphum* (Volvocales, Chlorophyta) from Japan. *Nova Hedwigia Beiheft*, **112**, 483-490.
- 241 Nozaki, H., Hara, Y. & Kasaki, H. 1987 Light and electron microscopy of pyrenoids and species delimitation in *Volvulina* (Chlorophyta, Volvocaceae). *J. Phycol.*, **23**, 359-364.
- 242 Nozaki, H. & Ito, M. 1994 Phylogenetic relationships within the colonial Volvocales (Chlorophyta) inferred from cladistic analysis based on morphological data. *J. Phycol.*, **30**, 353-365.
- 243 Nozaki, H., Ito, M., Sano, R., Uchida, H., Watanabe, M.M. & Kuroiwa, T. 1995 Phylogenetic relationships within the colonial Volvocales (Chlorophyta) inferred from *rbcL* gene sequence data. *J. Phycol.*, **31**, 970-979.
- 244 Nozaki, H., Ito, M., Sano, R., Uchida, H., Watanabe, M. M., Takahashi, H. & Kuroiwa, T. 1997 Phylogenetic analysis of *Yamagishiella* and *Platydorina* (Volvocaceae, Chlorophyta) based on *rbcL* gene sequences. *J. Phycol.*, **33**, 272-278.
- 245 Nozaki, H., Ito, M., Uchida, H., Watanabe, M. M. & Kuroiwa, T. 1997 Phylogenetic analysis of *Eudorina* species (Volvocaceae, Chlorophyta) based on *rbcL* gene sequences. *J. Phycol.*, **33**, 859-863.
- 246 Nozaki, H., Ito, M., Watanabe, M. M. & Kuroiwa, T. 1996 Ultrastructure of the vegetative colonies and systematic position of *Basichlamys* (Volvocales, Chlorophyta). *Eur. J. Phycol.*, **31**, 67-72.

- 247 Nozaki, H., Ito, M., Watanabe, M. M., Takano, H. & Kuroiwa, T. 1997 Phylogenetic analysis of morphological species of *Carteria* (Volvocales, Chlorophyta) based on *rbcL* gene sequences. *J. Phycol.*, **33**, 864-867.
- 248 Nozaki, H., Katagiri, M., Nakagawa, M., Aizawa, K. & Watanabe, M. M. 1995 Taxonomic re-examination of the two strains labeled "*Chlorella*" in the Microbial Culture Collection at the National Institute for Environmental Studies (NIES-Collection). *Microbiol. Cult. Coll.*, **11**, 11-18.
- 249 Nozaki, H., Kuroiwa, H. & Kuroiwa, T. 1994 Light and electron microscopic characterization of two types of pyrenoids in *Gonium* (Goniaceae, Chlorophyta). *J. Phycol.*, **30**, 279-290.
- 250 Nozaki, H., Kuroiwa, H., Mita, T. & Kuroiwa, T. 1989 *Pleodorina japonica* sp. nov. (Volvocales, Chlorophyta) with bacteria-like endosymbionts. *Phycologia*, **28**, 252-267.
- 251 Nozaki, H. & Kuroiwa, T. 1990 *Volvolina compacta* sp. nov. (Volvocaceae, Chlorophyta) from Nepal. *Phycologia*, **29**, 410-417.
- 252 Nozaki, H. & Kuroiwa, T. 1991 Morphology and sexual reproduction of *Gonium multicoccum* (Volvocales, Chlorophyta) from Nepal. *Phycologia*, **30**, 381-393.
- 253 Nozaki, H. & Kuroiwa, T. 1991 *Pandorina colemaniae* sp. nov. (Volvocaceae, Chlorophyta) from Japan. *Phycologia*, **30**, 449-457.
- 254 Nozaki, H. & Kuroiwa, T. 1992 Ultrastructure of the extracellular matrix and taxonomy of *Eudorina*, *Pleodorina* and *Yamagishiella* gen. nov. (Volvocaceae, Chlorophyta). *Phycologia*, **31**, 529-541.
- 255 Nozaki, H., Ohta, N., Morita, E. & Watanabe, M. M. 1998 Toward a natural system of species in *Chlorogonium* (Volvocales, Chlorophyta): a combined analysis of morphological and *rbcL* gene sequence data. *J. Phycol.*, **34**, 1024-1037.
- 256 Nozaki, H. & Ohtani, S. 1992 *Gonium sociale* (Volvocales, Chlorophyta) from Antarctica. *Jpn. J. Phycol.*, **40**, 267-271.
- 257 Nozaki, H., Song, L., Liu, Y., Hiroki, M. & Watanabe, M. M. 1998 Taxonomic re-examination of a chinese strain labeled '*Eudorina* sp.' (Volvocaceae, Chlorophyta) based on morphological and DNA sequence data. *Phycol. Rés.*, **46(Suppl.)**, 63-70.
- 258 Ogawa, T. & Terui, G. 1970 Studies on the growth of *Spirulina platensis*. (I) On the pure culture of *Spirulina platensis*. *J. Ferment. Technol.*, **48**, 361-367.
- 259 Ohmori, M. & Hattori, A. 1971 Heterocyst and nitrogen fixation in *Anabaena cylindrica*. In *Proceedings of the Seventh International Seaweed Symposium*, University of Tokyo Press, Tokyo, p. 598-601.
- 260 Ohmori, M. & Hattori, A. 1971 Nitrogen fixation and heterocysts in the blue-green alga *Anabaena cylindrica*. *Plant & Cell Physiol.*, **12**, 961-967.
- 261 Ohmori, M. & Hattori, A. 1972 Effect of nitrate on nitrogen-fixation by the blue-green alga *Anabaena cylindrica*. *Plant & Cell Physiol.*, **13**, 589-599.
- 262 Ohmori, M. & Hattori, A. 1974 Effect of ammonia on nitrogen fixation by the blue-green alga *Anabaena cylindrica*. *Plant & Cell Physiol.*, **15**, 131-142.
- 263 Oikawa, E., Ohnuma, T., Shimizu, A. & Ishibashi, Y. 1994 Phylogenetic studies on musty odor producing cyanobacteria using the 16S ribosomal RNA. [16S ribosômu RNA enki-hairetsu no hikaku ni yoru kabi-shû sansai ransô-ru no bunrui.] *J. Jpn. Water Works Assoc. [Suidô-Kyôkai Zasshi]*, **63(12)**, 56-63 (in Japanese without English title).
- 264 Okaichi, T., Nishio, S. & Imatomi, Y. 1982 Collection and mass culture [Shiryô no saisyû to baiyô]. In *Toxic phytoplankton - Occurrence, mode of action, and toxins [Yûdoku Purankuton - Hassei, Sayôkikô, Dokuseibun]*, Ed. by Jpn. Fish. Soc., Kôseisya-Kôseikaku, Tokyo, p. 23-34 (in Japanese without English title).
- 265 Okamura, K. 1992 Growth characteristics of *Skeletonema costatum* for algal growth potential. *Kanagawa-ken Kankyô Kagaku Sentâ Kenkyû Hôkoku*, **15**, 12-15 (in Japanese with English title).
- 266 Okino, T., Matsuda, H., Murakami, M. & Yamaguchi, K. 1993 Microginin, an angiotensin-converting enzyme inhibitor from

- the blue-green alga *Microcystis aeruginosa*. *Tetrahedron Letters*, **34**, 501-504.
- 267 Okino, T., Matsuda, H., Murakami, M. & Yamaguchi, K. 1995 New microviridins, elastase inhibitors from the blue-green alga *Microcystis aeruginosa*. *Tetrahedron*, **51**, 10679-10686.
- 268 Okino, T., Murakami, M., Haraguchi, R., Munekata, H., Matsuda, H. & Yamaguchi, K. 1993 Micropeptins A and B, plasmin and trypsin inhibitors from the blue-green alga *Microcystis aeruginosa*. *Tetrahedron Letters*, **34**, 8131-8134.
- 269 Okino, T., Sun, Q., Matsuda, H., Murakami, M. & Yamaguchi, K. 1997 Nostopeptins A and B, elastase inhibitors from the cyanobacterium *Nostoc minutum*. *J. Nat. Prod.*, **60**, 158-161.
- 270 Okuyama, H., Morita, N. & Kogame, K. 1992 Occurrence of octadecapentaenoic acid in lipids of a cold stenothermic alga, prymnesiophyte strain B. *J. Phycol.*, **28**, 465-472.
- 271 Omura, T., Umita, T., Aizawa, J., Komatsu, Y., Ishizaki, M. & Onuma, M. 1991 Accumulation of extracellular non-volatile organic acids in batch cultures of *Chlorella vulgaris* and *Microcystis aeruginosa*. *Jpn. J. Water Poll. Res.*, **14**, 615-623 (in Japanese with English summary).
- 272 Ooi, T., Kusumi, T., Kakisawa, H. & Watanabe, M. M. 1989 Structure of cyanoviridin RR, a toxin from the blue-green alga, *Microcystis viridis*. *J. Appl. Phycol.*, **1**, 31-38.
- 273 Osafune, T. 1973 Three-dimensional structures of giant mitochondria, dictyosomes and "concentric lamellar bodies" formed during the cell cycle of *Euglena gracilis* (Z) in synchronous culture. *J. Electron Microscopy*, **22**, 51-61.
- 274 Osafune, T., Mihara, S., Hase, E. & Ohkuro, I. 1975 Formation and division of giant mitochondria during the cell cycle of *Euglena gracilis* Z in synchronous culture. I. Some characteristics of changes in the morphology of mitochondria and oxygen-uptake activity of cells. *Plant & Cell Physiol.*, **16**, 313-326.
- 275 Osafune, T., Mihara, S., Hase, E. & Ohkuro, I. 1975 Formation and division of giant mitochondria during the cell cycle of *Euglena gracilis* Z in synchronous culture. II. Modes of division of giant mitochondria. *J. Electron Microscopy*, **24**, 33-39.
- 276 Osafune, T., Mihara, S., Hase, E. & Ohkuro, I. 1975 Formation and division of giant mitochondria during the cell cycle of *Euglena gracilis* Z in synchronous culture. III. Three-dimensional structures of mitochondria after division of giant forms. *J. Electron Microscopy*, **24**, 283-286.
- 277 Otsuki, A., Watanabe, M. M. & Sugahara, K. 1987 Chlorophyll pigments in methanol extracts from ten axenic cultured diatoms and three green algae as determined by reverse phase HPLC with fluorometric detection. *J. Phycol.*, **23**, 406-414.
- 278 Pedersén, M. & DaSilva, E. J. 1973 Simple brominated phenols in the bluegreen alga *Calothrix brevissima* West. *Planta*, **115**, 83-86.
- 279 Pjon, C.-J. & Fujita, Y. 1973 Immunological identification of pigment component of a photochemically active chromoprotein (ACP) isolated from the blue-green alga *Anabaena cylindrica*. *Plant & Cell Physiol.*, **14**, 201-205.
- 280 Pjon, C.-J. & Fujita, Y. 1974 The separation and characterization of the photoreactive component of a photochemically active chromoprotein (ACP) obtained from the blue-green alga *Anabaena cylindrica*. *Plant & Cell Physiol.*, **15**, 789-797.
- 281 Pjon, C.-J., Suzuki, R. & Fujita, Y. 1972 Photochemical activity fluorescence and absorption spectrum of the photochemically active chromoprotein (ACP) isolated from the blue-green alga *Anabaena cylindrica*. *Plant & Cell Physiol.*, **13**, 437-447.
- 282 Pringsheim, E. G. 1946 The biphasic or soil-water culture method for growing algae and flagellata. *J. Ecol.*, **33**, 193-204.
- 283 Provasoli, L. 1963 Growing marine seaweeds. In *Proceedings of the Fourth International Seaweed Symposium*, University of Tokyo Press, Tokyo, p. 9-17.
- 284 Provasoli, L. 1968 Media and prospects for the cultivation of marine algae. In *Cultures and Collections of Algae*, Eds. by Watanabe, A. & Hattori, A., Proc. U.S.-Japan Conf., Hakone, Sept.

- 1966., Jpn. Soc. Plant Physiol., p. 63-75.
- 285 Provasoli, L., McLaughlin, J. J. A. & Droop, M. R. 1957 The development of artificial media for marine algae. *Arch. Mikrobiol.*, **25**, 392-428.
- 286 Provasoli, L. & Pintner, I. J. 1959 Artificial media for fresh-water algae: problems and suggestions. In *The Ecology of Algae. Spec. Pub. No. 2.*, Eds. by Tryon, C. A., Jr. & Hartmann, R. T., Pymatuning Laboratory of Field Biology, University of Pittsburgh, Pittsburgh, p. 84-96.
- 287 Sakamoto, S., Yamaguchi, M., Watanabe, M. F., Watanabe, M. & Kamiya, H. 1996 Distribution and characterization of lectins from natural and cultured *Microcystis* spp. In *Harmful and Toxic algal Blooms*, Eds. by Yasumoto, T., Oshima, T. & Fukuyo, Y., Intergovernmental Oceanographic Commission of UNESCO 1996, p. 569-572.
- 288 Sako, Y., Adachi, M. & Ishida, Y. 1995 Specific monoclonal antibodies and DNA probes for the identification of the toxic dinoflagellate genus *Alexandrium*. In *Harmful Marine Algal Blooms*, Eds. by Lassus, P., et al., Lavoisier, Intercept Ltd., p. 77-82.
- 289 Sako, Y., Ishida, Y., Kadota, H. & Hata, Y. 1984 Sexual reproduction and cyst formation in the freshwater dinoflagellate *Peridinium cunningtonii*. *Bull. Jpn. Soc. Sci. Fish.*, **50**, 743-750.
- 290 Sano, T. & Kaya, K. 1995 Oscillamide Y, a chymotrypsin inhibitor from toxic *Oscillatoria agardhii*. *Tetrahedron Letters*, **36**, 5933-5936.
- 291 Sano, T. & Kaya, K. 1995 A 2-amino-2-butenoic acid (Dhb)-containing microcystin isolated from *Oscillatoria agardhii*. *Tetrahedron Letters*, **36**, 8603-8606.
- 292 Sano, T. & Kaya, K. 1996 Oscillapeptin G, a tyrosinase inhibitor from toxic *Oscillatoria agardhii*. *J. Nat. Prod.*, **59**, 90-92.
- 293 Sartoni, G. & Sonni, C. 1991 *Tribonema marinum* J. Feldmann and *Acinetospora crinita* (Carmichael) Sauvageau in the benthic mucilaginous aggregates observed along Tuscan shores during the summer of 1991. *Informatore Botanico Italiano*, **23**, 23-30 (in Italian with English summary).
- 294 Sasa, T., Suda, S., Watanabe, M. M. & Takaichi, S. 1992 A yellow marine *Chlamydomonas*: morphology and pigment composition. *Plant & Cell Physiol.*, **33**, 527-534.
- 295 Sasa, T., Takaichi, S., Hatakeyama, N. & Watanabe, M. M. 1992 A novel carotenoid ester, loroxanthin dodecenoate, from *Pyramimonas parkeae* (Prasinophyceae) and a chlorarachnio-phycean alga. *Plant & Cell Physiol.*, **33**, 921-925.
- 296 Satoh, E., Satoh, S. & Fujii, T. 1992 Synchronization of phosphorylation of a chloroplast protein with the cell division cycle in *Heterosigma akashiwo*. *Plant & Cell Physiol.*, **33**, 275-280.
- 297 Sawada, K., Handa, N., Shiraiwa, Y., Danbara, A. & Montani, S. 1996 Long-chain alkenones and alkyl alkenoates in the coastal and pelagic sediments of the northwest North Pacific, with special reference to reconstruction of *Emiliana huxleyi* and *Gephyrocapsa oceanica* ratios. *Org. Geochem.*, **24**, 751-764.
- 298 Scheibe, J. 1972 Photoreversible pigment: occurrence in a blue-green alga. *Science*, **176**, 1037-1039.
- 299 Schönhuber, W., Zarda, B., Eix, S., Rippka, R., Herdman, M., Ludwig, W. & Amann, R. 1999 *In situ* identification of cyanobacteria with horseradish peroxidase-labeled, rRNA-targeted oligonucleotide probes. *Appl. Environm. Microbiol.*, **65**, 1259-1267.
- 300 Sekiguchi, M., Ohtsuki, A., Shibata, Y. & Morita, M. 1995 Arsenic compounds accumulated in phytoplanktons of marine origin. *J. Environ. Chem. [Kankyō-Kagaku]*, **5**, 366-367 (in Japanese with English title).
- 301 Sekimoto, H. & Fujii, T. 1991 Sexual reproduction of microalgae: its physiological and biochemical study. *Cell Science*, **7**, 562-571 (in Japanese with English title).
- 302 Sekimoto, H. & Fujii, T. 1992 Analysis of gametic protoplast release in the *Closterium peracerosum -strigosum-littorale* complex (Chlorophyta). *J. Phycol.*, **28**, 615-619.
- 303 Sekimoto, H. & Fujii, T. 1993 Cell-cell interaction involved in the sexual reproduction of *Closterium* [Mikadukimo no yūsei seisyoku ni kakawaru saibōkan jōhō kōkan]. *Iden*, **47** (1),

- 43-48 (in Japanese without English title).
- 304 Sekimoto, H., Satoh, S. & Fujii, T. 1990 Biochemical and physiological properties of a protein inducing protoplast release during conjugation in the *Closterium peracerosum-strigosum-littorale* complex. *Planta*, **182**, 348-354.
- 305 Sekimoto, H., Satoh, S. & Fujii, T. 1992 Biochemical and physiological properties of a gametic protoplast-release-inducing protein in *Closterium*. *Korean J. Phycol.*, **7**, 121-129.
- 306 Sekimoto, H., Satoh, S. & Fujii, T. 1993 Analysis of binding of biotinylated protoplast-release-inducing protein that induces release of gametic protoplasts in the *Closterium peracerosum-strigosum-littorale* complex. *Planta*, **189**, 468-474.
- 307 Sekimoto, H., Sone, Y. & Fujii, T. 1994 cDNA cloning of a 42-kilodalton subunit of protoplast-release-inducing protein from *Closterium*. *Plant Physiol.*, **104**, 1095-1096.
- 308 Sekimoto, H., Sone, Y. & Fujii, T. 1994 A cDNA encoding a 19-kilodalton subunit of protoplast-release inducing protein from *Closterium*. *Plant Physiol.*, **105**, 447.
- 309 Sekimoto, H., Sone, Y. & Fujii, T. 1994 Regulation of expression of the genes for a sex pheromone by an inducer of the sex pheromone in the *Closterium peracerosum-strigosum-littorale* complex. *Planta*, **193**, 137-144.
- 310 Sekimoto, H., Sone, Y. & Fujii, T. 1995 Biochemical, physiological, and molecular analysis of sexual isolation in the species complex *Closterium peracerosum-strigosum-littorale* (Chlorophyta). *J. Phycol.*, **31**, 611-615.
- 311 Sekino, K., Kobayashi, H. & Shiraiwa, Y. 1996 Role of coccoliths in the utilization of inorganic carbon by a marine unicellular coccolithophorid, *Emiliana huxleyi*: a survey using intact cells and protoplasts. *Plant Cell Physiol.*, **37**, 123-127.
- 312 Sekino, K. & Shiraiwa, Y. 1994 Accumulation and utilization of dissolved inorganic carbon by a marine unicellular coccolithophorid, *Emiliana huxleyi*. *Plant Cell Physiol.*, **35**, 353-361.
- 313 Sekino, K. & Shiraiwa, Y. 1995 CO₂ release from the culture of calcifying and noncalcifying marine unicellular algae. *J. Mar. Biotechnol.*, **3**, 101-104.
- 314 Sekino, K. & Shiraiwa, Y. 1996 Evidence for the involvement of mitochondrial respiration in calcification in a marine coccolithophorid, *Emiliana huxleyi*. *Plant Cell Physiol.*, **37**, 1030-1033.
- 315 Shibukawa, K., Takamura, Y., Takao, E., Irie, M. & Yagi, O. 1986 Studies on the growth factors of an axenic clone of the cyanobacterium *Microcystis aeruginosa* K-3A. *Jpn. J. Limnol.*, **47**, 247-255 (in Japanese with English summary).
- 316 Shin, H. J., Matsuda, H., Murakami, M. & Yamaguchi, K. 1996 Agardhipeptins A and B, two new cyclic hepta- and octapeptide, from the cyanobacterium *Oscillatoria agardhii* (NIES-204). *Tetrahedron*, **52**, 13129-13136.
- 317 Shin, H. J., Matsuda, H., Murakami, M. & Yamaguchi, K. 1997 Aeruginosins 205A and -B, serine protease inhibitory glycopeptides from the cyanobacterium *Oscillatoria agardhii* (NIES-205). *J. Org. Chem.*, **62**, 1810-1813.
- 318 Shin, H. J., Matsuda, H., Murakami, M. & Yamaguchi, K. 1997 Anabaenopeptins E and F, two new cyclic peptides from the cyanobacterium *Oscillatoria agardhii* (NIES-204). *J. Nat. Prod.*, **60**, 139-141.
- 319 Shin, H. J., Matsuda, H., Murakami, M. & Yamaguchi, K. 1997 Circinamide, a novel papain inhibitor from the cyanobacterium *Anabaena circinalis* (NIES-41). *Tetrahedron*, **53**, 5747-5754.
- 320 Shin, H. J., Murakami, M., Matsuda, H., Ishida, K. & Yamaguchi, K. 1995 Oscillapeptin, an elastase and chymotrypsin inhibitor from the cyanobacterium *Oscillatoria agardhii* (NIES-204). *Tetrahedron Letters*, **36**, 5235-5238.
- 321 Shin, H. J., Murakami, M., Matsuda, H. & Yamaguchi, K. 1996 Microviridins D-F, serine protease inhibitors from the cyanobacterium *Oscillatoria agardhii* (NIES-204). *Tetrahedron*, **52**, 8159-8168.
- 322 Shinobu, K., Abe, S., Asano, K. & Abe, T. 1992 On the laboratory experiment of algal culturing (AGP-test) [Sōrui baiyō no shitsunai jikken (AGP shiken) ni tsuite]. *Heisei 4-nendo Tōhoku Chihō Kensetsu-kyoku Kan-nai Gijutsu Kenkyū*

- Happyôkai Shiryô*, **Kyô-28**, 151-154 (in Japanese without English title).
- 323 Shono, M., Hara, Y., Wada, M. & Fujii, T. 1996 A sodium pump in the plasma membrane of the marine alga *Heterosigma akashiwo*. *Plant Cell Physiol.*, **37**, 385-388.
- 324 Shono, M., Wada, M. & Fujii, T. 1995 Partial purification of a Na⁺-ATPase from the plasma membrane of the marine alga *Heterosigma akashiwo*. *Plant Physiol.*, **108**, 1615-1621.
- 325 Soma, Y., Imaizumi, T., Yagi, K. & Kasuga, S. 1993 Estimation of Algal Succession in lake water using HPLC analysis of pigments. *Can. J. Fish. Aquat. Sci.*, **50**, 1142-1146.
- 326 Suda, S. & Watanabe, M. M. 1995 Life cycle of *Pseudocarteria mucosa* (Korschikov) Ettl (Volvocales, Chlorophyta). *Phycologia*, **34**, 58-64.
- 327 Suda, S. & Watanabe, M. M. 1995 Notes on microalgae in Japan (2). *Hafniomonas montana* (Geitler) Ettl et Moestrup (Chlorophyceae, Dunaliellales). *Jpn. J. Phycol. (Sôru)*, **43**, 40-41 (in Japanese with English title).
- 328 Suda, S., Watanabe, M. M. & Inouye, I. 1989 Evidence for sexual reproduction in the primitive green alga *Nephroselmis olivacea* (Prasinophyceae). *J. Phycol.*, **25**, 596-600.
- 329 Suda, S., Watanabe, M. M. & Inouye, I. 1990 Morphological observations on the rare quadriflagellate *Pseudocarteria mucosa* (Chlorophyceae, Volvocales). *Phycologia*, **29**, 54-64.
- 330 Sudo, R., Tai, S., Yagi, O., Okada, M., Hosomi, M. & Yamane, A. N. 1981 Comprehensive studies on the eutrophication of fresh-water areas. Determination of algal growth potential by algal assay procedure. *Res. Rep. Natl. Inst. Environ. Stud.*, **No. 26**, 1-53 (in Japanese with English title).
- 331 Sugahara, K., Kobayashi, Y., Watanabe, M. M. & Watanabe, M. 1982 Changes in cellular contents of ATP during growth of red tide algae. *Res. Rep. Natl. Inst. Environ. Stud.*, **No. 30**, 323-327 (in Japanese with English summary).
- 332 Sugaya, Y., Yasuno, M. & Yanai, T. 1990 Effects of toxic *Microcystis viridis* and isolated toxins on goldfish. *Jpn. J. Limnol.*, **51**, 149-153.
- 333 Sugiura, N., Iwami, N., Itayama, T. & Inamori, Y. 1997 Population dynamics of grazer *Monas guttula* and prey *Microcystis viridis* using aquatic large scale microcosm. *J. Jpn. Soc. Wat. Environ.*, **20**, 332-337.
- 334 Suzuki, R. & Fujita, Y. 1972 Studies on the Hill reaction of membrane fragments of blue-green algae II. The reaction steps inactivated at high water concentration. *Plant & Cell Physiol.*, **13**, 427-436.
- 335 Suzuki, S., Tazawa, R., Miya, A., Adachi, T., Kanki, R., Toyobe, M. & Oguchi, M. 1994 Gas revitalization by microalgae. I. Studies on species and culture condition. *CELSS J.*, **7**, 23-28 (in Japanese with English summary).
- 336 Tabata, K. 1996 Cyanobacteria as teaching materials for inquiring biology. [Tankyû katsudô no sozai to shite no shiano-bakteria.] *The Heredity [Iden]*, **50**(9), 58-62 (in Japanese without English title).
- 337 Takamura, N., Kasai, F. & Watanabe, M. M. 1988 Differences in the tolerant level of benthic algae to heavy metal - The effects of Cu, Cd, and Zn on the photosynthesis. *Res. Rep. Natl. Inst. Environ. Stud.*, **No. 114**, 223-232 (in Japanese with English summary).
- 338 Takamura, N., Kasai, F. & Watanabe, M. M. 1989 Effects of Cu, Cd and Zn on photosynthesis of freshwater benthic algae. *J. Appl. Phycol.*, **1**, 39-52.
- 339 Takamura, N., Kasai, F. & Watanabe, M. M. 1990 Unique response of Cyanophyceae to copper. *J. Appl. Phycol.*, **2**, 293-296.
- 340 Takeda, H. 1988 Classification of *Chlorella* strains by cell wall sugar composition. *Phytochem.*, **27**, 3823-3826.
- 341 Tan, S. & Ho, K.-K. 1989 Purification of an acidic plastocyanin from *Microcystis aeruginosa*. *Biochim. et Biophys. Acta*, **973**, 111-117.
- 342 Tanaka, S. & Nozaki, H. 1997 Notes on microalgae in Japan (3). *Pteromonas angulosa* (Carter) Lemmermann (Chlorophyceae, Volvocales). *Jpn. J. Phycol. (Sôru)*, **45**, 183-184 (in Japanese with English summary).
- 343 Tjahjono, A. E., Hayama, Y., Kakizono, T., Terada, Y., Nishio, N. & Nagai, S. 1994

- Hyper-accumulation of astaxanthin in a green alga *Haematococcus pluvialis* at elevated temperatures. *Biotech. Letters*, **16**, 133-138.
- 344 Tjahjono, A. E., Kakizono, T., Hayama, Y. & Nagai, S. 1993 Formation and regeneration of protoplast from a unicellular green alga *Haematococcus pluvialis*. *J. Ferment. Bioengineering*, **75**, 196-200.
- 345 Tjahjono, A. E., Kakizono, T., Hayama, Y., Nishio, N. & Nagai S. 1994 Isolation of resistant mutants against carotenoid biosynthesis inhibitors for a green alga *Haematococcus pluvialis*, and their hybrid formation by protoplast fusion for breeding of higher astaxanthin producers. *J. Ferment. Bioengineering*, **77**, 352-357.
- 346 Tominaga, H., Hayashida, Y., Hosoya, Y., Kurokawa, M., Sawa, Y. & Ochiai, H. 1993 Characterization of a small cryptic plasmid, pPF1, from *Phormidium foveolarum* and vector construction. *Biosci. Biotech. Biochem.*, **57**, 1795-1799.
- 347 Tominaga, H., Soejima, K., Kawagishi, S., Ashida, H., Sawa, Y. & Ochiai, H. 1993 Structural organization of a cryptic plasmid, pMA1, from *Microcystis aeruginosa* f. *aeruginosa* Kützing. *Biosci. Biotech. Biochem.*, **57**, 1503-1507.
- 348 Tsubo, Y. 1975 Formation of colorless cells in algae. In *Advance of Phycology in Japan*, Eds. by Tokida, J. & Hirose, H., VEB Gustav Fischer Verlag, Jena, p. 180-193.
- 349 Tsubo, Y., Ueda, T. & Yokomura, E. 1971 Chloroplast autonomy revealed by nutritionally controlled *Euglena gracilis*. In *Proceedings of the Seventh International Seaweed Symposium*, University of Tokyo Press, Tokyo, p. 336-338.
- 350 Tsuji, T. & Fujita, Y. 1971 Ascorbate photo-oxidation by a photochemically active chromoprotein isolated from the blue-green alga *Anabaena cylindrica*: the effect of monochromatic illumination. *Plant & Cell Physiol.*, **12**, 807-811.
- 351 Tsuji, T. & Fujita, Y. 1972 Electron donor-specificity observed in photosystem I reactions of membrane fragments of the blue-green alga *Anabaena variabilis* and the higher plant *Spinacea oleracea*. *Plant & Cell Physiol.*, **13**, 93-99.
- 352 Tsukui, T. & Yamazaki, M. 1992 Studies on eutrophication in rivers (II). Determination of AGP in the Kanda river system. *Tokyo-to Kankyô Kagaku Kenkyûjo Nenpô 1992*, 162-166 (in Japanese with English Summary).
- 353 Tsusue, Y. & Fujita, Y. 1964 Mono- and oligo-saccharides in the blue-green alga, *Tolypothrix tenuis*. *J. Gen. Appl. Microbiol.*, **10**, 283-294.
- 354 Turmel, M., Lemieux, C., Burger, G., Lang, B. F., Otis, C., Plante, I. & Gray, M. W. 1999 The complete mitochondrial DNA sequence of *Nephroselmis olivacea* and *Pedinomonas minor*: Two radically different evolutionary patterns within green algae. *The Plant Cell*, **11**, 1717-1729.
- 355 Turmel, M., Otis, C. & Lemieux, C. 1999 The complete chloroplast DNA sequence of the green alga *Nephroselmis olivacea*: Insights into the architecture of ancestral chloroplast genomes. *Proc. Natl. Acad. Sci. U.S.A.*, **96**, 10248-10253.
- 356 Ubekôsan Co. Ltd. 1988 Effects of UR-158 on the algal mortality [UR-158 no sassôkôka]. Unpublished report in Ubekôsan Co. Ltd., 3 pp. (in Japanese without English title).
- 357 Uchida, A., Nagasaki, K., Hiroishi, S. & Ishida, Y. 1989 The application of monoclonal antibodies to an identification of *Chattonella marina* and *Chattonella antiqua*. *Bull. Jpn. Soc. Sci. Fish.*, **55**, 721-725.
- 358 Uchida, H., Kitae, K., Tomizawa, K. & Yokota, A. 1998 Comparison of the nucleotide sequence and secondary structure of the 5.8S ribosomal RNA gene of *Chlamydomonas tetragama* with those of green algae. *DNA Sequence*, **8**, 403-408.
- 359 Ueda, K. & Nonaka, M. 1992 Division of chloroplasts in a green alga, *Pediastrum duplex*. *Ann. Bot.*, **69**, 113-118.
- 360 Ukai, Y., Fujita, Y., Morimura, Y. & Watanabe, A. 1958 Studies on growth of blue green alga *Tolypothrix tenuis*. *J. Gen. Appl. Microbiol.*, **4**, 163-169.
- 361 Uye, S. & Takamatsu, K. 1990 Feeding interactions between planktonic copepods and red-tide flagellates from Japanese coastal waters.

- Mar. Ecol. Prog. Ser.*, **59**, 97-107.
- 362 Wada, M., Hara, Y., Kato, M., Yamada, M. & Fujii, T. 1987 Diurnal appearance, fine structure, and chemical composition of fatty particles in *Heterosigma akashiwo* (Raphidophyceae). *Protoplasma*, **137**, 134-139.
- 363 Wada, M., Satoh, S., Kasamo, K. & Fujii, T. 1989 Presence of a Na⁺-activated ATPase in the plasma membrane of the marine raphidophycean *Heterosigma akashiwo*. *Plant & Cell Physiol.*, **30**, 923-928.
- 364 Wada, M., Urayama, O., Satoh, S., Hara, Y., Ikawa, Y. & Fujii, T. 1992 A marine algal Na⁺-activated ATPase possesses an immunologically identical epitope to Na⁺, K⁺-ATPase. *FEBS Letters*, **309**, 272-274.
- 365 Watanabe, A. 1959 On the mass-culturing of a nitrogen-fixing blue-green alga, *Tolypothrix tenuis*. *J. Gen. Appl. Microbiol.*, **5**, 85-91.
- 366 Watanabe, A. 1960 List of algal strains in collection at the Institute of Applied Microbiology, University of Tokyo. *J. Gen. Appl. Microbiol.*, **6**, 283-292.
- 367 Watanabe, A. 1962 Effect of nitrogen-fixing blue-green alga: *Tolypothrix tenuis* on the nitrogenous fertility of paddy soil and on the crop yield of rice plant. *J. Gen. Appl. Microbiol.*, **8**, 85-91.
- 368 Watanabe, A., Hattori, A., Fujita, Y. & Kiyohara, T. 1959 Large scale culture of a blue-green alga, *Tolypothrix tenuis*, utilizing hot spring and natural gas as heat and carbon dioxide sources. *J. Gen. Appl. Microbiol.*, **5**, 51-57.
- 369 Watanabe, A., Ito, R. & Sasa, T. 1955 Microalgae as a source of nutrients for daphnids. *J. Gen. Appl. Microbiol.*, **1**, 137-141.
- 370 Watanabe, A., Ito, R. & Sasa, T. 1955 Effect of various chemicals upon the growth of daphnids. *J. Gen. Appl. Microbiol.*, **1**, 190-193.
- 371 Watanabe, A., Shiota, M., Endo, H. & Yamamoto, Y. 1969 An observation of the practical applications of nitrogen fixing blue-green algae for rice cultivation. In *The Third International Conference on the Global Impacts of Applied Microbiology*, Bombay, p. 53-64.
- 372 Watanabe, A. & Yamamoto, Y. 1967 Heterotrophic nitrogen fixation by the blue-green alga *Anabaenopsis circularis*. *Nature*, **214**, 738.
- 373 Watanabe, A. & Yamamoto, Y. 1968 Effects of antibiotics on the growth of microalgae. *Phykos*, **7**, 248-258.
- 374 Watanabe, K. I., Ehara, M., Inagaki, Y. & Ohama, T. 1998 Distinctive origins of group I introns found in the *COXI* genes of three green algae. *Gene*, **213**, 1-7.
- 375 Watanabe, M. 1977 A preliminary study of *Gonium viridistellatum* sp. nov. (Chlorophyta, Volvocaceae). *Bull. Jpn. Soc. Phycol.*, **25**, Suppl. (Mem. Iss. Yamada), 379-384.
- 376 Watanabe, M. 1978 A taxonomic study of the *Closterium calosporum* complex (1). *Bull. Natn. Sci. Mus., Ser. B (Bot.)*, **4**, 133-154, pls. 1-8.
- 377 Watanabe, M. 1979 A taxonomic study of the *Closterium calosporum* complex (2). *Bull. Natn. Sci. Mus., Ser. B (Bot.)*, **5**, 1-23, pls. 4-6.
- 378 Watanabe, M. 1982 The diurnal variations in the cell densities of *Olisthodiscus luteus* and *Skeletonema costatum*. *Res. Rep. Natl. Inst. Environ. Stud.*, No. **30**, 143-154 (in Japanese with English summary).
- 379 Watanabe, M. & Harashima, A. 1982 Bioconvection in culture of *Olisthodiscus luteus* and Rayleigh-Taylor instability. *Res. Rep. Natl. Inst. Environ. Stud.*, No. **30**, 155-173 (in Japanese with English summary).
- 380 Watanabe, M. & Harashima, A. 1982 The distribution pattern of *Olisthodiscus luteus* in convection cells. *Res. Rep. Natl. Inst. Environ. Stud.*, No. **30**, 175-190 (in Japanese with English summary).
- 381 Watanabe, M., Kohata, K. & Kunugi, M. 1987 Diel vertical migration of *Heterosigma akashiwo* under salinity and phosphate stratifications and metabolism of intracellular phosphate pools. *Res. Rep. Natl. Inst. Environ. Stud.*, No. **110**, 103-114 (in Japanese with English summary).
- 382 Watanabe, M., Kohata, K. & Kunugi, M. 1987 ³¹P nuclear magnetic resonance study of intracellular phosphate pools and polyphosphate metabolism in *Heterosigma akashiwo* (Hada) Hada (Raphidophyceae). *J. Phycol.*, **23**, 54-62.

- 383 Watanabe, M., Kohata, K. & Kunugi, M. 1988 Phosphate accumulation and metabolism by *Heterosigma akashiwo* (Raphidophyceae) during diel vertical migration in a stratified microcosm. *J. Phycol.*, **24**, 22-28.
- 384 Watanabe, M., Takamatsu, T., Kohata, K., Kunugi, M., Kawashima, M. & Koyama, M. 1989 Luxury phosphate uptake and variation of intracellular metal concentrations in *Heterosigma akashiwo* (Raphidophyceae). *J. Phycol.*, **25**, 428-436.
- 385 Watanabe, M., Watanabe, M. M., Kohata, K. & Harashima, A. 1982 The use of a controlled experimental ecosystem (Microcosm) in studies of the mechanism of red tide outbreaks. *Res. Rep. Natl. Inst. Environ. Stud.*, No. **30**, 11-26 (in Japanese with English summary).
- 386 Watanabe, M. F., Oishi, S., Watanabe, Y. & Watanabe, M. 1986 Strong probability of lethal toxicity in the blue-green alga *Microcystis viridis* Lemmermann. *J. Phycol.*, **22**, 552-556.
- 387 Watanabe, M. M. 1977 Biosystematics in *Closterium* of sexual unicellular green algae and *Calothrix* and *Spirulina* of asexual filamentous blue-green algae, with special reference to the analyses of natural populations. *Thesis D. Sci.*, Hokkaido University, 114 pp., 28 tables, 54 figs.
- 388 Watanabe, M. M. 1983 Growth characteristics of freshwater red tide alga, *Peridinium* based on axenic culture. Establishment of synthetic culture medium [Junsuibaiyôhō niyoru tansuiakashio *Peridinium* no zôsyokutokusei no kaiseki-gôseibaichi no kakuritsu]. *Res. Data Natl. Inst. Environ. Stud.*, No. **24**, 111-121 (in Japanese without English title).
- 389 Watanabe, M. M. 1989 Toxicity of algae [Sôru no dokusei]. *Jpn. J. Water Poll. Res.*, **12**, 750-756 (in Japanese without English title).
- 390 Watanabe, M. M. & Hiroki, M. (Eds.) 1997 *NIES-Collection. List of Strains, Fifth Edition, 1997, Microalgae and Protozoa*. Microbial Culture Collection, National Institute for Environmental Studies, Tsukuba, 140 pp.
- 391 Watanabe, M. M. & Ichimura, T. 1977 Fresh- and salt-water forms of *Spirulina platensis* in axenic cultures. *Bull. Jpn. Soc. Phycol.*, **25**, *Suppl. (Mem. Iss. Yamada)*, 371-377.
- 392 Watanabe, M. M. & Ichimura, T. 1978 Biosystematic studies of the *Closterium peracerosum-strigosum-littorale* complex. II. Reproductive isolation and morphological variation among several populations from the Northern Kanto area in Japan. *Bot. Mag. Tokyo*, **91**, 1-10.
- 393 Watanabe, M. M. & Ichimura, T. 1978 Biosystematic studies of the *Closterium peracerosum-strigosum-littorale* complex. III. Degrees of sexual isolation among the three population groups from the Northern Kanto area. *Bot. Mag. Tokyo*, **91**, 11-24.
- 394 Watanabe, M. M. & Ichimura, T. 1982 Biosystematic studies of the *Closterium peracerosum-strigosum-littorale* complex. IV. Hybrid breakdown between two closely related groups, Group II-A and Group II-B. *Bot. Mag. Tokyo*, **95**, 241-247.
- 395 Watanabe, M. M. & Kasai, F. (Eds.) 1985 *NIES-Collection. List of Strains, First Edition, 1985, Microalgae*. Microbial Culture Collection, National Institute for Environmental Studies, Tsukuba, 116 pp.
- 396 Watanabe, M. M. & Kasai, F. (Eds.) 1985 *NIES-Collection. List of Strains, First Edition, 1985, Microalgae* [Hozonkabu-risuto dai 1 pan bisaisôru]. Microbial Culture Collection, National Institute for Environmental Studies, Tsukuba, 114 pp. (in Japanese without English title).
- 397 Watanabe, M. M. & Kasai, F. (Eds.) 1986 *Supplement to NIES-Collection. List of Strains, First Edition, 1985, Microalgae*. Microbial Culture Collection, National Institute for Environmental Studies, Tsukuba, 25 pp.
- 398 Watanabe, M. M. & Kasai, F. (Eds.) 1987 *Second Supplement to NIES-Collection. List of Strains, First Edition, 1985, Microalgae*. Microbial Culture Collection, National Institute for Environmental Studies, Tsukuba, 25 pp.
- 399 Watanabe, M. M., Kasai, F., Hiwatari, T., Suda, S. & Nei, T. 1984 Cryopreservation of microalgae in liquid nitrogen - Survival of algal cultures after freezing. *Jpn. J. Freez. Dry.*, **30**, 23-26 (in Japanese with English title).
- 400 Watanabe, M. M., Kasai, F. & Sudo, R. (Eds.) 1988 *NIES-Collection. List of Strains, Second*

- Edition, 1988, *Microalgae and Protozoa*. Microbial Culture Collection, National Institute for Environmental Studies, Tsukuba, 148 pp.
- 401 Watanabe, M. M., Kohata, K., Nakamura, Y. & Watanabe, M. 1982 Phosphate-limited continuous culture of a red tide flagellate, *Olisthodiscus luteus*: establishment of its method and the analysis of growth kinetics. *Res. Rep. Natl. Inst. Environ. Stud.*, No. 30, 113-130 (in Japanese with English summary).
- 402 Watanabe, M. M. & Kurogi, M. 1975 Taxonomic reexamination of the two species of blue-green algae, *Calothrix scopulorum* and *C. crustacea*. *Bot. Mag. Tokyo*, 88, 111-125.
- 403 Watanabe, M. M. & Nakamura, Y. 1984 Growth characteristics of a red tide flagellate, *Heterosigma akashiwo* Hada. 1. The effects of temperature, salinity, light intensity and pH on growth. *Res. Rep. Natl. Inst. Environ. Stud.*, No. 63, 51-58 (in Japanese with English summary).
- 404 Watanabe, M. M. & Nakamura, Y. 1984 Growth characteristics of a red tide flagellate, *Heterosigma akashiwo* Hada. 2. The utilization of nutrients. *Res. Rep. Natl. Inst. Environ. Stud.*, No. 63, 59-68 (in Japanese with English summary).
- 405 Watanabe, M. M., Nakamura, Y. & Kohata, K. 1983 Diurnal vertical migration and dark uptake of nitrate and phosphate of the red tide flagellates, *Heterosigma akashiwo* Hada and *Chattonella antiqua* (Hada) Ono (Raphidophyceae). *Jpn. J. Phycol.*, 31, 161-166.
- 406 Watanabe, M. M., Nakamura, Y. & Kohata, K. 1984 Diurnal vertical migration of a red tide flagellate, *Heterosigma akashiwo* Hada, with special reference to the ecological role. *Res. Rep. Natl. Inst. Environ. Stud.*, No. 63, 69-77 (in Japanese with English summary).
- 407 Watanabe, M. M., Nakamura, Y., Mori, S. & Yamochi, S. 1982 Effects of physico-chemical factors and nutrients on the growth of *Heterosigma akashiwo* Hada from Osaka Bay, Japan. *Jpn. J. Phycol.*, 30, 279-288.
- 408 Watanabe, M. M. & Nozaki, H. (Eds.) 1994 *NIES-Collection. List of Strains, Fourth Edition, 1994, Microalgae and Protozoa*. Microbial Culture Collection, National Institute for Environmental Studies, Tsukuba, 127 pp.
- 409 Watanabe, M. M. & Satake, K. N. (Eds.) 1991 *NIES-Collection. List of Strains, Third Edition, 1991, Microalgae and Protozoa*. Microbial Culture Collection, National Institute for Environmental Studies, Tsukuba, 163 pp.
- 410 Watanabe, M. M. & Sawaguchi, T. 1995 Cryopreservation of a water-bloom forming cyanobacterium, *Microcystis aeruginosa* f. *aeruginosa*. *Phycol. Res.*, 43, 111-116.
- 411 Watanabe, M. M., Suda, S., Kasai, F. & Sawaguchi, T. 1985 Axenic cultures of the three species of *Microcystis* (Cyanophyta = Cyano-bacteria). *Bull. JFCC*, 1, 57-63.
- 412 Watanabe, M. M., Takeuchi, Y. & Takamura, N. 1987 Cu tolerance of freshwater benthic diatom, *Achnanthes minutissima*. In *Biological Monitoring of Environmental Pollution*, Eds. by Yasuno, M. & Whitton, B. A., Tokai University Press, Tokyo, p. 171-177.
- 413 Watanabe, M. M., Takeuchi, Y. & Takamura, N. 1988 Copper tolerance of benthic diatom *Achnanthes minutissima*. *Res. Rep. Natl. Inst. Environ. Stud.*, No. 114, 233-243 (in Japanese with English summary).
- 414 Watanabe, M. M., Yamochi, S., Kohata, K. & Watanabe, M. 1982 Vertical migration and diel periodicity of phosphate uptake in a red tide flagellate, *Olisthodiscus luteus*. *Res. Rep. Natl. Inst. Environ. Stud.*, No. 30, 131-141 (in Japanese with English summary).
- 415 Watanabe, S. 1981 Observations on *Urnella terrestris* Playfair (Chlorophyceae, Chlorococcales) in culture. *Phycologia*, 20, 12-15.
- 416 Watanabe, S. 1983 New and interesting green algae from soils of some Asian and Oceanian regions. *Arch. Protistenk.*, 127, 223-270.
- 417 Watanabe, Y., Ohmura, N. & Saiki, H. 1992 Microbial CO₂ fixation. 2. Isolation and determination of cultural characteristics of *Chlorella* strains which function under CO₂ enriched atmosphere. *CRIEPI Kenkyū-Hōkoku*, U92014, 1-21 (in Japanese with English summary).
- 418 Yagi, O., Hagiwara, T., Takamura, Y. & Sudo, R. 1984 Growth characteristics of axenic and unialgal *Microcystis* isolated from Lake Kasumigaura. *Jpn. J. Water Poll. Res.*, 7, 496-

- 503 (in Japanese with English summary).
- 419 Yagi, O., Hagiwara, T., Takamura, Y. & Sudo, R. 1987 Limiting nutrients of algal growth in Lake Kasumigaura. *Jpn. J. Water Poll. Res.*, **10**, 115-122 (in Japanese with English summary).
- 420 Yagi, O., Ohkubo, N., Tomioka, N. & Okada, M. 1989 Limiting nutrients of algal growth in Ushiku Marsh. *Jpn. J. Limnol.*, **50**, 139-148 (in Japanese with English summary).
- 421 Yagi, O., Okada, M. & Sudo, R. 1979 Cultivation of *Microcystis* and red-tide-organisms. *Res. Rep. Natl. Inst. Environ. Stud.*, No. 6, 223-229 (in Japanese with English summary).
- 422 Yagi, O., Sudo, R., Imaoka, T. & Teranishi, S. 1984 Algal growth potential on municipal wastewater using red tide organisms. *Res. Rep. Natl. Inst. Environ. Stud.*, No. 63, 231-242 (in Japanese with English summary).
- 423 Yamada, N., Aoyama, K., Yamada, M. & Hamamura, N. 1985 Studies on earthy-musty odor in natural water (1). Growth characteristics and 2-methylisoborneol production of *Phormidium tenue*. *Jpn. J. Water Poll. Res.*, **8**, 515-521 (in Japanese with English summary).
- 424 Yamada, N., Aoyama, K., Yamada, M. & Hamamura, N. 1986 Studies on earthy-musty odor in natural water (3). Isolation of bacteria-free *Phormidium tenue* and the effect of associated bacteria on the growth of axenic *P. tenue*. *Jpn. J. Water Poll. Res.*, **9**, 379-385 (in Japanese with English summary).
- 425 Yamaguchi, K., Murakami, M. & Okino, T. 1989 Screening of angiotensin-converting enzyme inhibitory activities in microalgae. *J. Appl. Phycol.*, **1**, 271-275.
- 426 Yamamoto, Y. 1972 The fatty acid composition of akinetes, heterocysts and vegetative cells in *Anabaena cylindrica*. *Plant & Cell Physiol.*, **13**, 913-915.
- 427 Yamamoto, Y. 1975 Effect of desiccation on the germination of akinetes of *Anabaena cylindrica*. *Plant & Cell Physiol.*, **16**, 749-752.
- 428 Yamamoto, Y. 1976 Effect of some physical and chemical factors on the germination of akinetes of *Anabaena cylindrica*. *J. Gen. Appl. Microbiol.*, **22**, 311-323.
- 429 Yamamoto, Y. 1978 Detection of algal lysing biological agents in lakes by the soft-agar overlay technique. *Jpn. J. Limnol.*, **39**, 9-14 (in Japanese with English summary).
- 430 Yamamoto, Y. 1981 Observation on the occurrence of microbial agents which cause lysis of blue-green algae in Lake Kasumigaura. *Jpn. J. Limnol.*, **42**, 20-27.
- 431 Yamamoto, Y. & Suzuki, K. 1990 Distribution and algal-lysing activity of fruiting myxobacteria in Lake Suwa. *J. Phycol.*, **26**, 457-462.
- 432 Yamatogi, T., Miyahara, J. & Tanaka, J. 1996 Preculture of noxious red tide flagellate *Chattonella antiqua* (Raphidophyceae) for AGP assay. *Bull. Nagasaki Pref. Inst. Fish.*, **22**, 7-13 (in Japanese with English summary).
- 433 Yamochi, S. 1983 Mechanisms for outbreak of *Heterosigma akashiwo* red tide in Osaka Bay, Japan. Part 1. Nutrient factors involved in controlling the growth of *Heterosigma akashiwo* Hada. *J. Oceanogr. Soc. Japan*, **39**, 310-316.
- 434 Yanagida, Y. 1992 An assay system for detecting the effects of weed killers, by using the green microalga *Scenedesmus quadricauda* [Bisyô ryokusô *Scenedesmus quadricauda* wo mochiita josôzai no eikyôhyôkahô nitsuite]. *Report of Freshwater Fisheries Experiment Station Ibaraki Prefecture*, No. 28, 124-127 (in Japanese without English title).
- 435 Yasumoto, T. 1992 Toxin of the marine microalgae [Kaiyô bisaisôru no Tokishin]. *BIO Medica*, **7**, 57-62 (in Japanese without English title).
- 436 Yoshida, T. & Ancajas, R. R. 1970 Application of the acetylene reduction method in nitrogen fixation studies. *Soil Sci. & Plant Nutr.*, **16**, 234-237.
- 437 Yoshida, Y. & Kawaguchi, K. 1983 Buoyancy and phototaxis of *Chattonella antiqua* (Hada) Ono. *Bull. Plankton Soc. Jpn.*, **30**, 11-19 (in Japanese with English summary).
- 438 Yoshizako, F., Nishimura, A. & Chubachi, M. 1992 Microbial reduction of cyclohexanone by *Chlorella pyrenoidosa* Chick. *J. Ferment.*

Bioengineering, 74, 395-397.

- 439 Yoshizako, F., Nishimura, A. & Chubachi, M. 1994 Identification of algal transformation products from alicyclic ketones. *J. Ferment. Bioengineering*, 77, 144-147.
- 440 Yoshizako, F., Nishimura, A., Chubachi, M., Horii, T. & Ueno, T. 1991 Bioconversion of cyclohexaneacetic acid to monohydroxycyclohexaneacetic acids by *Chlorella pyrenoidosa* Chick. *J. Ferment. Bioengineering*, 72, 343-346.
- 441 Yoshizako, F., Ogino, M., Nishimura, A., Chubachi, M. & Horii, T. 1995 Biotransformation of cyclic β -keto esters by *Chlorella pyrenoidosa* Chick. *J. Ferment. Bioengineering*, 79, 141-145.
- 442 Yuki-jirushi [Snow Brand] Milk Products Co.Ltd. & Showa Denko K.K. 1992 Separation and purification of useful substances from *Euglena*. [*Euglena* kara no yûyô-busshitsu no Bunri Seisei.] In *The Advanced Technology of the Separation and Purification of the Materials for Physiologically Functional Foods and Their Development* [*Kinôsei Shokuhin Sozai no Kôdo Bunri Seisei to Kaihatsu*], Ed. and published by The Japanese Research and Development Association for the High Separation System in Food Industry [*Shokuhin Sangyô Hai-separêshon Sisutemu Gijutu Kenkyû Kumiai*], Tokyo, p.363-382 (in Japanese without English title).

Committee for Evaluating Microbial Culture Strains

Makoto M. Watanabe, D. Sci.	Environmental Biology Division	<Chairman>
Kunimitsu Kaya, D. A.	Environmental Chemistry Division	<Sub-Chairman>
Osami Yagi, D. A.	Regional and Community Environment Division	
Yuhei Inamori, D. Sci.	Regional and Community Environment Division	
Noriko Takamura, Ph. D	Regional and Community Environment Division	
Hiroo Uchiyama, D. A.	Water and Soil Environment Division	
Fumie Kasai, D. Sci.	Environmental Biology Division	
Mikiya Hiroki, D. A.	Environmental Biology Division / Research Coordinator	<Secretary>
Masanobu Kawachi, D. Sci.	Environmental Biology Division	
Tomomichi Yanagita, D. Sci.	Professor Emeritus of University of Tokyo	<Adviser>
Mitsuo Chihara, D. Sci.	Director-General of the Natural History Museum and Institute, Chiba	<Adviser>
Kazuo Komagata, D. A.	Professor of Tokyo University of Agriculture	<Adviser>
Yoshinobu Shigenaka, D. Sci.	Professor Emeritus of Hiroshima University	<Adviser>
Yuzaburo Ishida, D. A.	Professor of Fukuyama University	<Adviser>
Tsuneyoshi Kuroiwa, D. Sci.	Professor of University of Tokyo	<Adviser>
Isao Inouye, D. Sci.	Professor of the University of Tsukuba	<Adviser>

January 2000
Accepted by Editorial Board
National Institute for Environmental Studies

[Research Records of National Institute for Environmental Studies R-152-2000]
ISSN 1341-3463

NIES-Collection. List of Strains
Sixth Edition
Microalgae and Protozoa
2000

Correspondence
Makoto M. Watanabe
Environmental Microbiology Laboratory
Environmental Biology Division
National Institute for Environmental Studies

1 March, 2000

Published by National Institute for Environmental Studies
16-2 Onogawa, Tsukuba, Ibaraki 305-0053, Japan,
Tel: +81-298-50-2556 Fax: +81-298-50-2587

Printed by Nissei Eblo Co.,Ltd.
Address: 3-13-11, Azuma, Tsukuba, Ibaraki 305-0031 Japan

Cover Photos

Front Cover

1	4	5	
2	3		
	6	7	8
10			9
	11	13	14
12			15

- 1 *Chamaesiphon polymorphus* (NIES-433)
- 2 *Hyalotheca dissiliens* v. *dissiliens* f. *tridentula* (NIES-294)
- 3 *Chattonella antiqua* (NIES-85)
- 4 *Staurastrum paradoxum* (NIES-528)
- 5 *Pyrocystis lunura* (NIES-609)
- 6 *Dinobryon divergens* (NIES-284)
- 7 *Phormidium jenkelianum* (NIES-506)
- 8 *Pterosperma cristatum* (NIES-626)
- 9 *Calothrix scopulorum* (NIES-268)
- 10 *Volvox aureus* (NIES-396)
- 11 *Micrasterias crux-melitensis* (NIES-152)
- 12 *Odontella aurita* (NIES-589)
- 13 *Anabaena spiroides* f. *spiroides* (NIES-77)
- 14 *Pleodorina indica* (NIES-736)
- 15 *Cryptomonas rostratiformis* (NIES-345)

Back Cover

16	18	19	20
17			
	21	22	23
24	25	26	
	29		27 31
28		30	

- 16 *Ceratium hirundinella* (NIES-376)
- 17 *Trentepohlia* sp.
- 18 *Euglena* sp.
- 19 *Cricosphaera roscoffensis* (NIES-8)
- 20 *Monoraphidium griffithii* (NIES-385)
- 21 *Cyanophora tetracyanea* (NIES-764)
- 22 *Porphyridium* sp.
- 23 *Lagerheimia ciliata* (NIES-382)
- 24 *Fibrocapsa japonica* (NIES-462)
- 25 *Carteria eugametos* (NIES-631)
- 26 *Nitzschia palea* (NIES-488)
- 27 *Chroomonas coerulea* (NIES-713)
- 28 *Microcystis aeruginosa* f. *aeruginosa* (NIES-100)
- 29 *Chlorarachnion reptans* (NIES-624)
- 30 *Nostoc minutum* (NIES-29)
- 31 *Acinetospora crinita* (NIES-548)

Photos : Kosei Yumoto (1-24, 27-31)
Shoichiro Suda (25, 26)